

10-25-00

A Box  
SD

Express Mail Label No. EL547697129US  
Date of Deposit: October 23, 2000  
File No.: 99-75

JC894 U.S. PTO  
10/23/00

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE  
APPLICATION FEE TRANSMITTAL

Commissioner for Patents  
Box Patent Application  
Washington, D.C. 20231

Sir:

JC914 U.S. PTO  
09/695369  
10/23/00

Transmitted herewith for filing under 37 CFR §1.53(b) is the

- ☒ patent application ☐ continuation patent application  
☐ divisional patent application ☐ continuation-in-part patent application

of Applicants: Wenfeng Xu, Catherine E. Lofton-Day, Randall Henne, Scott R. Presnell, Yue Yao,  
Julia E. Novak, Donald C. Foster, David P. Yee

Title: UMLR POLYPEPTIDES

- ☒ 119 pages of specification ☐ \_\_\_\_\_ sheets of drawings  
☒ 31 pages of sequence listing  
☐ An assignment of the invention to \_\_\_\_\_  
☒ 3 sheets of ☐ signed ☒ unsigned Declaration and Power of Attorney  
☒ **ASCII Computer Disk Sequence pursuant to 37 C.F.R. 1.821(f). It is believed that the content of the paper sequence listing and the computer readable sequence listing are the same.**

CALCULATION OF APPLICATION FEE


Claim Type	No. Filed	Less	Extra	Extra Rate	Fee
Total	38	-20	18	\$18.00	\$324.00
Independent	9	-3	6	\$80.00	\$480.00
Basic Fee					\$710.00
Multiple Dependency Fee					
If Applicable (\$270.00)					\$000.00
<b>Total Filing Fee</b>					<b>\$1,514.00</b>

- ☐ Priority of application Serial No. \_\_\_\_\_ filed on \_\_\_\_\_ in \_\_\_\_\_ is claimed under 35 U.S.C. 119. A certified copy thereof is submitted herewith.
- ☒ The benefit of applications Serial No's. 60/160,880 (filed on October 22, 1999); 60/163,215 (filed on November 2, 1999); 60/218,769 (filed on July 17, 2000); 60/222,221 (filed on August 1, 2000), all in the U.S. Patent and Trademark Office, is claimed under 35 U.S.C. 120 or 119(e) 1.

Please charge ZymoGenetics, Inc., Deposit Account No. 26-0290 as follows:

- ☒ Filing fee, estimated to be \$1,514.00  
☐ Assignment recording fee  
☒ Any additional fees associated with this paper or during the pendency of this application.  
☐ The issue fee set in 37 C.F.R. 1.18 at or before mailing of the Notice of Allowance, pursuant to 37 C.F.R. 1.311(b).

A copy of this sheet is enclosed.

Respectfully submitted,  
  
Robyn Adams  
Registration No. 44,495

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

**EXPRESS MAIL CERTIFICATE**

Box Patent Application  
Commissioner for Patents  
Washington, DC 20231



Re: U.S. Patent Application for  
UMLR POLYPEPTIDES

Applicants: Wenfeng Xu, Catherine E. Lofton-Day, Randall Henne, Scott R.  
Presnell, Yue Yao, Julia E. Novak, Donald C. Foster, David P. Yee

Sir:

Express Mail Label No. EL547697129US

Date of Deposit October 23, 2000

I hereby certify that the following attached paper(s) or fee

1. Return Post card
2. Application Fee Transmittal (in duplicate)
3. Patent Application (119 pages)
4. Unexecuted Declaration and Power of Attorney
5. Sequence Listing (31 pages)
6. ASCII Computer Disk Sequence pursuant to 37 CFR 1.821(e).

are being deposited with the United States Postal Service "Express Mail Post Office to Addressee" under 37 C.F.R. 1.10 on the date indicated above, addressed to the Commissioner for Patents, Washington, DC 20231.

  
Amy Toman

ZymoGenetics, Inc  
1201 Eastlake Avenue East  
Seattle, WA 98102  
(206) 442-6600

File Number : 99-75  
Filing Date: October 23, 2000  
Express Mail Label No. EL547697129US

UNITED STATES PATENT APPLICATION

OF

Wenfeng Xu, Catherine E. Lofton-Day, Randall Henne, Scott R.

Presnell, Yue Yao, Julia E. Novak, Donald C. Foster, David P.

Yee

FOR

UMLR POLYPEPTIDES

Description

## UMLR POLYPEPTIDES

5

## REFERENCE TO RELATED APPLICATIONS

This application is related to Provisional Applications 60/160, 880 filed on October 22, 1999; 60/163, 215, filed on November 2, 1999; 60/218,769, filed July 17,  
10 2000; and 60/222,221, filed on August 1, 2000. Under 35 U.S.C. § 119(e)(1), this application claims benefit of said Provisional Applications.

## BACKGROUND OF THE INVENTION

Cellular interactions which occur during an immune response are  
15 regulated by members of several families of cell surface receptors, including the tumor necrosis factor receptor (TNFR) family. The TNFR family consists of a number of integral membrane glycoprotein receptors many of which, in conjunction with their respective ligands, regulate interactions between different hematopoietic cell lineages (Smith et al., The TNF Receptor Superfamily of Cellular and Viral Proteins: Activation,  
20 Costimulation and Death, 76:959-62, 1994; Cosman, Stem Cells 12:440-55, 1994).

The TNF receptor family is composed of a number of type I integral membrane glycoproteins which exhibit sequence homology, particularly with respect to cysteine-rich repeats in their extracellular domains. The TNF receptor family includes p75 NGFR (Johnson et al., Cell 47:545-54, 1989), p55 TNFR-I (Loetscher et al., Cell  
25 61:351-59, 1990), p75 TNFR-II (Schall et al., Cell 61:361-70, 1990), TNFR-RP/TNFR-III (Crowe et al. Science 264:707-10, 1994), CD27 (Camerini et al., J. Immunol. 147:3165-69, 1991), CD30 (Falini et al., Blood 85:1-14, 1995), CD40 (Clark and Lane, Annu. Rev. Immunol. 9:97-127, 1991), 4-1BB (Kwon and Weissman, Proc. Natl. Acad. Sci. USA 86:1963-67, 1989; Schwarz et al., Gene 134:295-298, 1993), OX40 (Mallet et al., EMBO J. 9:1063-68, 1990), FAS/APO-1 (Itoh et al., Cell 66:233-43, 1991), DR3  
30 (Chinnaiyan et al., Science 274:990-92, 1996) also known as WSL-1 (Kitson et al.,

Nature 384:327-75, 1996), DR4 (Pan et al., Science 276:111-13, 1997), DR5 (Pan et al., Science 277:815-8, 1997; Sheridan et al., Science 277:818-21, 1997), osteoprotegerin (OPG) (Simonet et al., Cell 89:309-19, 1997; Human Genome Science, WIPO Publication WO96/28546), CAR1, found in chickens (Brojatsch et al., Cell 87:845-55, 5 1996), TRID or DcR1 (Pan et al., Science 277:815-8, 1997; Sheridan et al., Science 277:818-21, 1997) plus several viral open reading frames encoding TNFR-related molecules. NGFR, TNFR-I, CD30, CD40, 4-1BB, DR3 and OX40 are mainly restricted to cells of the lymphoid/hematopoietic system. TNFR-I, TNFR-II, TNFR-III and DR4 are found in most human tissues.

10               Members of the TNF receptor family are characterized by a multi-domain structure comprising an extracellular region, a transmembrane domain, a linker region between the extracellular ligand-binding region and the transmembrane domain and a cytoplasmic domain, which in several members of this family (TNFR 1, Fas, DR3, DR4, DR5, CAR1 and low affinity NGFR) contains a death domain associated with apoptosis.

15   One member, TRID or DcR1 (Pan et al., Science 277:815-8, 1997; Sheridan et al., ibid.) has a hydrophobic N-terminus with cysteine rich repeats #2 and #3 followed by five tandem repeats of 15 amino acid residues which concludes with a transmembrane domain. The extracellular ligand-binding region is characterized by the presence of one to six cysteine-rich motifs each containing about six cysteines and approximately 40

20   amino acids, although variation in the size and number of these motifs occurs among members of this family. The cysteine-rich regions provide the motif for binding to shared structures in the ligands. The highest degree of homology among the TNFR family members is within this extracellular cysteine-rich region. Among human TNFRs the average homology is in the range of 25% to 30%. Between the last cysteine-rich

25   repeat and the transmembrane domain is a small spacer region of between 8 to 70 amino acid residues. Cell surface TNF receptors are anchored in the cell membrane by a transmembrane domain characterized by a sequence of hydrophobic amino acid residues. On the opposite end of the protein from the extracellular ligand-binding region and separated from it by the transmembrane domain is the cytoplasmic domain. The

30   cytoplasmic domains of TNFR family members are small, from 46 to 221 amino acid residues, which suggests possible differences in the signaling mechanisms among family

members. In the TNF receptor for example, activation is triggered by the aggregation of cytoplasmic domains of three receptors when their corresponding extracellular domains bind to trimeric ligand.

One member of the TNF receptor family, osteoprotegerin (Simonet et al., *ibid.*), is unique in that it is a secreted protein. Soluble forms of other TNF receptors have been described for TNFR-I, TNFR-II, low-affinity NGFR, FAS, CD27, CD30, CD40 and 4-1BB, but these were generated either by cleaving from the cell membrane or secreted by alternatively spliced mRNA. OPG inhibits osteoclast maturation and it is thought that it might serve to regulate bone density by modulating osteoclast differentiation from hematopoietic precursors. OPG provided protection from normal osteoclast remodeling and ovariectomy-associated bone loss.

Ligands for these receptors have been identified, and with one exception (NGF) belong to the TNF ligand family. The members of the TNF ligand family share approximately 20% sequence identity in the extracellular ligand-binding regions, and exist mainly as type II membrane glycoproteins, biologically active as trimeric or multimeric complexes. This group includes TNF, LT-a, LT-b (Browning et al., *Cell* 72:847-56, 1993), CD27L (Goodwin et al., *Cell* 73:447-56, 1993), CD30L (Smith et al., *Cell* 73:1349-60, 1993), CD40L (Armitage et al., *Nature* 357:80-82, 1992), 4-1BBL (Goodwin et al., *Eur. J. Immunol.* 23:2631-41, 1993), OX40L (Godfrey et al., *J. Exp. Med.* 180:757-62, 1994), TRAIL or apo-2 (Wiley et al., *Immunity* 3:673-82, 1995), TNFg (Human Genome Sciences, WIPO Publication WO96/14328) and FasL (Cosman, *ibid.*; Lotz et al., *J. Leuko. Biol.* 60:1-7, 1996). Soluble ligand forms have been identified for TNF, LT-a and FasL. It is not known whether a specific protease cleaves each ligand, releasing it from the membrane, or whether one protease serves the same function for all TNF ligand family members. TACE (TNF-alpha converting enzyme) has been shown to cleave TNF (Moss et al., *Nature* 385:733-36, 1997; Black et al., *Nature* 385:729-33, 1997). No other such enzymes are known.

The X-ray crystallographic structures have been resolved for human TNF (Jones et al., *Nature* 338:225-28, 1989), LT-a (Eck et al., *J. Biol. Chem.* 267:2119-122, 1992) and the LT-a/TNFR complex (Banner et al., *Cell* 73:431-45, 1993). This complex features three receptor molecules bound symmetrically to one LT-a trimer. A model of

trimeric ligand binding through receptor oligomerization has been proposed to initiate signal transduction pathways. The identification of biological activity of several TNF members has been facilitated through use of monoclonal antibodies specific for the corresponding receptor. These monoclonal antibodies tend to be stimulatory when  
 5 immobilized and antagonistic in soluble form. This is further evidence that receptor crosslinking is a prerequisite for signal transduction in this receptor family. Importantly, the use of receptor-specific monoclonal antibodies or soluble receptors in the form of multimeric Ig fusion proteins has been useful in determining biological function *in vitro* and *in vivo* for several family members. Soluble receptor-Ig fusion proteins have been  
 10 used successfully in the cloning of the cell surface ligands corresponding to the CD40, CD30, CD27, 4-1BB and Fas receptors.

In general, the members of the tumor necrosis factor ligand family mediate interactions between different hematopoietic cells, such as T cell/B cell, T cell/monocyte and T cell/T cell interactions. The result of this two-way communication  
 15 can be stimulatory or inhibitory, depending on the target cell or the activation state. These TNF proteins are involved in regulation of cell proliferation, activation and differentiation, including control of cell survival or death by apoptosis or cytotoxicity. One member of this family, OX-40, is restricted to T cells where it acts as a costimulatory receptor. However, among the TNFR family members there are  
 20 differences in distribution, kinetics of induction and requirements for induction, which support a defined role for each of the ligands in T cell-mediated immune responses.

The demonstrated *in vitro* and *in vivo* activities of these TNF receptor ligand family members illustrate the enormous clinical potential of, and need for, other TNF receptors, TNF ligands, TNFR agonists, and TNFR antagonists. The present  
 25 invention addresses this need by providing a novel TNF receptor and related compositions and methods.

#### SUMMARY OF THE INVENTION

Within one aspect, the invention provides an isolated polypeptide  
 30 comprising residues 2 to 129 of SEQ ID NO:2. Within an embodiment, the isolated polypeptide comprises residues 2 to X of SEQ ID NO:2; wherein X is an integer between

130 and 136. Within another embodiment, the isolated polypeptide comprises residues 1 to 129 of SEQ ID NO:2. Within another embodiment, the isolated polypeptide comprises residues 1 to X of SEQ ID NO:2, wherein X is an integer between 130 and 136. Within another embodiment, the isolated polypeptide is selected from the group consisting of: a) polypeptides comprising residues 1 to 161 of SEQ ID NO:2; and b) polypeptides comprising residues 1 to 269 of SEQ ID NO:2.

Within another aspect, the invention provides an isolated polypeptide selected from the group consisting of: a) polypeptides comprising residues 1 to 297 of SEQ ID NO:27; b) polypeptides comprising residues 1 to 267 of SEQ ID NO:29; and c) polypeptides comprising residues 1 to 299 of SEQ ID NO:35.

Within another aspect, the invention provides an isolated polypeptide comprising residues 1 to 173 of SEQ ID NO:38.

Within another aspect, is provided an isolated polypeptide selected from the group consisting of: a) polypeptide molecules comprising residues 1 to 147 of SEQ ID NO:38; b) polypeptide molecules comprising residues 1 to 154 of SEQ ID NO:38; c) polypeptide molecules comprising residues 1 to 163 of SEQ ID NO:38; and d) polypeptide molecules comprising residues 1 to 165 of SEQ ID NO:38.

Within another aspect, the invention provides an isolated polynucleotide that encodes the polypeptide comprising residues 2 to 129 of SEQ ID NO:2. Within an embodiment, the isolated polynucleotide encodes a polypeptide that comprises residues 2 to X of SEQ ID NO:2; wherein X is an integer between 130 and 136. Within another embodiment, the isolated polynucleotide encodes a polypeptide that comprises residues 1 to 129 of SEQ ID NO:2. Within another embodiment, the isolated polynucleotide encodes a polypeptide that comprises residues 1 to X of SEQ ID NO:2, wherein X is an integer between 130 and 136. Within another embodiment, the isolated polypeptide encodes a polypeptide that is selected from the group consisting of: a) polypeptides comprising residues 1 to 161 of SEQ ID NO:2; and b) polypeptides comprising residues 1 to 269 of SEQ ID NO:2.

Within another aspect, the invention provides an expression vector comprising the following operably linked elements: a) a transcription promoter; b) a DNA segment wherein the DNA segment is a polynucleotide molecule encoding the



polypeptide molecule of comprising residues 1 to 129 of SEQ ID NO:2 ; and a transcription terminator. Within an embodiment, the DNA segment contains an affinity tag. Within another embodiment, the invention provides a cultured cell into which has been introduced the expression vector, wherein said cell expresses the polypeptide encoded by the DNA segment. Within another embodiment, the invention provides a method of producing a polypeptide comprising culturing the cell, whereby said cell expresses the polypeptide encoded by the DNA segment; and recovering the polypeptide. Within another embodiment, is provided the polypeptide produced by the cell.

Within another aspect, the invention provides a method of detecting lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma comprising, contacting said lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma with a polynucleotide consisting of the polynucleotide sequence as shown in SEQ ID NOs:1, 31, 32, 33, or 36, wherein the polynucleotide hybridizes to the nucleic acid in the lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma.

Within an embodiment the polynucleotide is selected from the group consisting of: a polynucleotide consisting of at least 16 contiguous nucleotides as shown in SEQ ID NO:1; a polynucleotide consisting of from 17 to 25 contiguous nucleotides as shown in SEQ ID NO:1; a polynucleotide consisting of 40 contiguous nucleotides as shown in SEQ ID NO:1; a polynucleotide consisting of 60 contiguous nucleotides as shown in SEQ ID NO:1; a polynucleotide consisting of at least 16 contiguous nucleotides as shown in SEQ ID NO:31; a polynucleotide consisting of from 17 to 25 contiguous nucleotides as shown in SEQ ID NO:31; a polynucleotide consisting of 40 contiguous nucleotides as shown in SEQ ID NO:31; a polynucleotide consisting of 60 contiguous nucleotides as shown in SEQ ID NO:31; a polynucleotide consisting of at least 16 contiguous nucleotides as shown in SEQ ID NO:33; a polynucleotide consisting of from 17 to 25 contiguous nucleotides as shown in SEQ ID NO:33; a polynucleotide consisting of 40 contiguous nucleotides as shown in SEQ ID NO:33; a polynucleotide consisting of 60 contiguous nucleotides as shown in SEQ ID NO:33; a polynucleotide consisting of at least 16 contiguous nucleotides as shown in SEQ ID NO:36; a polynucleotide consisting of from 17 to 25 contiguous nucleotides as shown in SEQ ID NO:36; a polynucleotide consisting of 40 contiguous nucleotides as shown in SEQ ID NO:36; a polynucleotide

consisting of 60 contiguous nucleotides as shown in SEQ ID NO:36 a polynucleotide encoding a polypeptide consisting of at least 16 contiguous nucleotides as shown in SEQ ID NO:38; a polynucleotide encoding a polypeptide consisting of from 17 to 25 contiguous nucleotides as shown in SEQ ID NO:38; a polynucleotide encoding a polypeptide consisting of 40 contiguous nucleotides as shown in SEQ ID NO:38; a polynucleotide encoding a polypeptide consisting of 60 contiguous nucleotides as shown in SEQ ID NO:38; and a polynucleotide encoding a polypeptide, wherein the polypeptide comprises residues 139 to 297 of SEQ ID NO:27, or residues 139 to 267 of SEQ ID NO:29.

10                Within another embodiment is method of detecting lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma comprising, contacting said lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma with an antibody to the polypeptide comprising residues 2 to X wherein x is an integer from 130 to 136 . Within an embodiment, the antibody is generated to a polypeptide selected from

15 the group consisting of: a polypeptide comprising residues 4 to 13 of SEQ ID NO:2; a polypeptide comprising residues 17 to 39 of SEQ ID NO:2; a polypeptide comprising residues 44 to 60 of SEQ ID NO:2; a polypeptide comprising residues 69 to 104 of SEQ ID NO:2; a polypeptide comprising residues 106 to 115 of SEQ ID NO:2; a polypeptide comprising residues 128 to 136 of SEQ ID NO:2; a polypeptide comprising residues 165 to 174

20 of SEQ ID NO:2; a polypeptide comprising residues 179 to 204 of SEQ ID NO:2; a polypeptide comprising residues 245 to 260 of SEQ ID NO:2; a polypeptide comprising residues 1 to 13 of SEQ ID NO:2; a polypeptide comprising residues 21 to 38 of SEQ ID NO:2; a polypeptide comprising residues 44 to 59 of SEQ ID NO:2; a polypeptide comprising residues 88 to 99 of SEQ ID NO:2; a polypeptide comprising residues 165 to

25 173 of SEQ ID NO:2; a polypeptide comprising residues 194 to 205 of SEQ ID NO:2; and a polypeptide comprising residues 213 to 221 of SEQ ID NO:2. Within another embodiment the method of detecting lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma according to claim 19 wherein the antibody is generated to a polypeptide selected from the group consisting of: a polypeptide comprising residues 181

30 to 210 of SEQ ID NO:35; a polypeptide comprising residues 219 to 226 of SEQ ID NO:35; a polypeptide comprising residues 218 to 236 of SEQ ID NO:35; a polypeptide

comprising residues 162 to 169 of SEQ ID NO:38; a polypeptide comprising residues 149 to 173 of SEQ ID NO:38; a polypeptide comprising residues 194 to 236 of SEQ ID NO:35; a polypeptide comprising residues 150 to 171 of SEQ ID NO:38; a polypeptide comprising residues 230 to 235 of SEQ ID NO:35; and a polypeptide comprising residues 151 to 158 of SEQ ID NO:38. Within another embodiment, the method of detecting lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma according to claim 19 wherein the antibody is generated to a polypeptide selected from the group consisting of: a polypeptide consisting of residues 4 to 13 of SEQ ID NO:2; a polypeptide consisting of residues 17 to 39 of SEQ ID NO:2; a polypeptide consisting of residues 44 to 60 of SEQ ID NO:2; a polypeptide consisting of residues 69 to 104 of SEQ ID NO:2; a polypeptide consisting of residues 106 to 115 of SEQ ID NO:2; a polypeptide consisting of residues 128 to 136 of SEQ ID NO:2; a polypeptide consisting of residues 165 to 174 of SEQ ID NO:2; a polypeptide consisting of residues 179 to 204 of SEQ ID NO:2; a polypeptide consisting of residues 245 to 260 of SEQ ID NO:2; a polypeptide consisting of residues 1 to 13 of SEQ ID NO:2; a polypeptide consisting of residues 21 to 38 of SEQ ID NO:2; a polypeptide consisting of residues 44 to 59 of SEQ ID NO:2; a polypeptide consisting of residues 88 to 99 of SEQ ID NO:2; a polypeptide consisting of residues 165 to 173 of SEQ ID NO:2; a polypeptide consisting of residues 194 to 205 of SEQ ID NO:2; and a polypeptide consisting of residues 213 to 221 of SEQ ID NO:2. Within another embodiment, the method of detecting lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma wherein the antibody is generated to a polypeptide selected from the group consisting of: a polypeptide consisting of residues 181 to 210 of SEQ ID NO:35; a polypeptide consisting of residues 219 to 226 of SEQ ID NO:35; a polypeptide consisting of residues 218 to 236 of SEQ ID NO:35; a polypeptide consisting of residues 162 to 169 of SEQ ID NO:38; a polypeptide consisting of residues 149 to 173 of SEQ ID NO:38; a polypeptide consisting of residues 194 to 236 of SEQ ID NO:35; a polypeptide consisting of residues 150 to 171 of SEQ ID NO:38; a polypeptide consisting of residues 230 to 235 of SEQ ID NO:35; and a polypeptide consisting of residues 151 to 158 of SEQ ID NO:38.

Within another aspect of the invention, is provided a method of inhibiting the quantity of lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or

lymphoma cells expressing a polypeptide selected from the group consisting of: an isolated polypeptide as shown in SEQ ID NO:2 ; an isolated polypeptide as shown in SEQ ID NO:27; an isolated polypeptide as shown in SEQ ID NO:29; and an isolated polypeptide as shown in SEQ ID NO:35; comprising administering to the cells an isolated polypeptide wherein the isolated polypeptide consists of residues 1 to X of SEQ ID NO:2, wherein X is an integer between 129 and 136. Within another aspect the method uses an isolated polypeptide comprising residues 1 to 173 fo SEQ ID NO:38.

Within another aspect, the invention provides a method of modulating apoptosis of lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma cells comprising administering to the cells an isolated polypeptide wherein the isolated polypeptide comprising residues 2 to X of SEQ ID NO:2, wherein X is an integer from 129 to 136. Within an embodiment the method of modulating apoptosis of lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma cells, uses the isolated polypeptide comprising residues 1 to X, wherein X is an integer from 129 to 136. Within another aspect the method uses an isolated polypeptide comprising residues 1 to 173 of SEQ ID NO:38.

Within another aspect, the invention provides an isolated polypeptide consisting of residues 2 to 129 of SEQ ID NO:2. Within an embodiment, the isolated polypeptide consists of residues 2 to X of SEQ ID NO:2; wherein X is an integer between 130 and 136. Within another embodiment, the isolated consists of residues 1 to 129 of SEQ ID NO:2. Within another embodiment, the isolated polypeptide consists of residues 1 to X of SEQ ID NO:2, wherein X is an integer between 130 and 136. Within another embodiment, the isolated polypeptide is selected from the group consisting of: polypeptides consisting of residues 1 to 161 of SEQ ID NO:2; and polypeptides consisting of residues 1 to 269 of SEQ ID NO:2. Within another embodiment, the isolated polypeptide selected from the group consisting of: polypeptides consisting of residues 1 to 297 of SEQ ID NO:27; polypeptides consisting of residues 1 to 267 of SEQ ID NO:29; and polypeptides consisting of residues 1 to 299 of SEQ ID NO:35.

Within another aspect, the invention provides an isolated polypeptide consisting of residues 1 to 173 of SEQ ID NO:38.

Within another aspect, the invention provides an isolated polypeptide selected from the group consisting of: polypeptide molecules consisting of residues 1 to 147 of SEQ ID NO:38; polypeptide molecules consisting of residues 1 to 154 of SEQ ID NO:38; polypeptide molecules consisting of residues 1 to 163 of SEQ ID NO:38; and  
 5 polypeptide molecules consisting of residues 1 to 165 of SEQ ID NO:38. Within an embodiment the isolated polynucleotide encodes a polypeptide consisting of residues 2 to 129 of SEQ ID NO:2 .

Within one aspect, the invention provides an isolated polypeptide molecule comprising residues 1 to 129 of SEQ ID NO:2. Within an embodiment the  
 10 isolated polypeptide molecule comprises residues 1 to 136 of SEQ ID NO:2. Within another embodiment the isolated polypeptide molecule comprises residues 1 to 269 of SEQ ID NO:2.

Within another aspect the invention provides an isolated polypeptide molecule selected from the group consisting of: a polypeptide molecule comprising  
 15 residues 2 to 42 of SEQ ID NO:2; a polypeptide molecule comprising residues 43 to 84 of SEQ ID NO:2; a polypeptide molecule comprising residues 85 to 129 of SEQ ID NO:2; a polypeptide molecule comprising residues 130 to 136 of SEQ ID NO:2; a polypeptide molecule comprising residues 137 to 161 of SEQ ID NO:2; and a polypeptide molecule comprising residues 162 to 269 of SEQ ID NO:2.

20 Within another aspect the invention provides a isolated polynucleotide molecule encoding a polypeptide molecule wherein the polypeptide molecule comprises residues 1 to 120 of SEQ ID NO:2.

Within another aspect, the invention provides an isolated polynucleotide molecule encoding a polypeptide molecule wherein the polypeptide molecule comprises  
 25 residues 1 to 136 of SEQ ID NO:2.

Within another aspect of the invention is provided an isolated polynucleotide molecule encoding a polypeptide molecule wherein the polypeptide molecule comprises residues 1 to 269 of SEQ ID NO:2.

30 Within another aspect, an isolated polynucleotide molecule encoding a polypeptide molecule wherein the polypeptide molecule is selected from the group consisting of: a polypeptide molecule comprising residues 2 to 42 of SEQ ID NO:2; a

polypeptide molecule comprising residues 43 to 84 of SEQ ID NO:2; a polypeptide molecule comprising residues 85 to 129 of SEQ ID NO:2; a polypeptide molecule comprising residues 130 to 136 of SEQ ID NO:2; a polypeptide molecule comprising residues 137 to 161 of SEQ ID NO:2; and a polypeptide molecule comprising residues 162 to 269 of SEQ ID NO:2 is provided.

Within another aspect, the invention provides an expression vector comprising the following operably linked elements: a transcription promoter; a DNA segment wherein the DNA segment is a polynucleotide molecule encoding the polypeptide molecule of claim 1; and a transcription terminator. Within an embodiment, the expression vector according contains an affinity tag. Within another embodiment, is provided a cultured cell into which has been introduced the expression vector according, and the cell expresses the polypeptide encoded by the DNA segment. Within a further embodiment, the invention provides a method of producing a polypeptide comprising culturing the cell, whereby said cell expresses the polypeptide encoded by the DNA segment; and recovering the polypeptide.

Within another aspect, the invention provides a method of producing an antibody comprising the following steps in order: inoculating an animal with a polypeptide selected from the group consisting of: a polypeptide molecule consisting of residues 2 to 42 of SEQ ID NO:2; a polypeptide molecule consisting of residues 43 to 84 of SEQ ID NO:2; a polypeptide molecule consisting of residues 85 to 129 of SEQ ID NO:2; a polypeptide molecule consisting of residues 130 to 136 of SEQ ID NO:2; a polypeptide molecule consisting of residues 137 to 161 of SEQ ID NO:2; a polypeptide molecule consisting of residues 162 to 269 of SEQ ID NO:2; a polypeptide molecule consisting of residues 1 to 269 of SEQ ID NO:2; a polypeptide molecule consisting of residues 1 to 129 of SEQ ID NO:2; a polypeptide molecule consisting of residues 1 to 136 of SEQ ID NO:2; and a polypeptide molecule consisting of residues 137 to 269 of SEQ ID NO:2 wherein the polypeptide elicits an immune response in the animal to produce the antibody; and isolating the antibody from the animal. Within an embodiment, the antibody produced by the method binds to a residues 1 to 269 of SEQ ID NO:2. Within an embodiment, is provided the antibody of claim 14, wherein the

antibody is a monoclonal antibody. Within an embodiment, the antibody specifically binds to the polypeptide.

Within another aspect, the invention provides a method of producing an antibody comprising the following steps in order: inoculating an animal with an epitope bearing portion of a polypeptide wherein the epitope bearing portion is selected from the group consisting of: a polypeptide molecule consisting of residues 4 to 13 of SEQ ID NO:2; a polypeptide molecule consisting of residues 17 to 39 of SEQ ID NO:2; a polypeptide molecule consisting of residues 44 to 60 of SEQ ID NO:2; a polypeptide molecule consisting of residues 69 to 104 of SEQ ID NO:2; a polypeptide molecule consisting of residues 106 to 115 of SEQ ID NO:2; a polypeptide molecule consisting of residues 128 to 136 of SEQ ID NO:2; a polypeptide molecule consisting of residues 165 to 174 of SEQ ID NO:2; a polypeptide molecule consisting of residues 179 to 204 of SEQ ID NO:2; a polypeptide molecule consisting of residues 245 to 260 of SEQ ID NO:2; a polypeptide molecule consisting of residues 1 to 13 of SEQ ID NO:2; a polypeptide molecule consisting of residues 21 to 38 of SEQ ID NO:2; a polypeptide molecule consisting of residues 44 to 59 of SEQ ID NO:2; a polypeptide molecule consisting of residues 88 to 99 of SEQ ID NO:2; a polypeptide molecule consisting of residues 165 to 173 of SEQ ID NO:2; a polypeptide molecule consisting of residues 213 to 221 of SEQ ID NO:2 wherein the polypeptide elicits an immune response in the animal to produce the antibody; and isolating the antibody from the animal. Within an embodiment, the antibody produced by the method binds to a residues 1 to 269 of SEQ ID NO:2. Within an embodiment, the antibody is a monoclonal antibody. Within an embodiment, the antibody specifically binds to the polypeptide.

Within another aspect, the invention provides a method of forming a reversible peptide-receptor complex comprising; providing a receptor wherein the receptor comprises residues 1 to 129 of SEQ ID NO:2; and contacting the receptor with a peptide; whereby the receptor binds the peptide. Within an embodiment, is provided a method wherein the peptide is selected from the group consisting of: TNF peptides; antibodies; and other binding proteins.

Within another aspect, the invention provides an isolated polypeptide molecule selected from the groups consisting of: a polypeptide molecule comprising

residues 1 to 297 of SEQ ID NO:27; and a polypeptide molecule comprising residues 1 to 267 of SEQ ID NO:29. Within an embodiment, the invention provides a polynucleotide encoding the isolated polypeptide.

Within another aspect, the invention provides a method of detecting carcinoma in lung tissue or breast tissue comprising, contacting said carcinoma with a polynucleotide consisting of the polynucleotide sequence as shown in SEQ ID NO:1, wherein the polynucleotide hybridizes to the nucleic acid in the carcinoma tissue. Within an embodiment, the polynucleotide is selected from the group consisting of: a polynucleotide consisting of at least 16 contiguous nucleotides as shown in SEQ ID NO:1; a polynucleotide consisting of from 17 to 25 contiguous nucleotides as shown in SEQ ID NO:1; a polynucleotide consisting of 40 contiguous nucleotides as shown in SEQ ID NO:1; a polynucleotide consisting of 60 contiguous nucleotides as shown in SEQ ID NO:1; a polynucleotide selected from the group consisting of: the polynucleotide as shown in SEQ ID NO:10; the polynucleotide as shown in SEQ ID NO:11; the polynucleotide as shown in SEQ ID NO:12; the polynucleotide as shown in SEQ ID NO:13; the polynucleotide as shown in SEQ ID NO:23; the polynucleotide as shown in SEQ ID NO:24; the polynucleotide as shown in SEQ ID NO:25; and the polynucleotide as shown in SEQ ID NO:26; a polynucleotide encoding a polypeptide, wherein the polypeptide comprises residues 139 to 297 of SEQ ID NO:27, or residues 139 to 267 of SEQ ID NO:29.

Within another aspect, the invention provides a method of detecting carcinoma in lung tissue or breast tissue comprising, contacting said carcinoma with an antibody to the polypeptide as shown in SEQ ID NO:2. Within an embodiment, the antibody is generated to a polypeptide selected from the group consisting of: a polypeptide comprising residues 4 to 13 of SEQ ID NO:2; a polypeptide comprising residues 17 to 39 of SEQ ID NO:2; a polypeptide comprising residues 44 to 60 of SEQ ID NO:2; a polypeptide comprising residues 69 to 104 of SEQ ID NO:2; a polypeptide comprising residues 106 to 115 of SEQ ID NO:2; a polypeptide comprising 128 to 136 of SEQ ID NO:2; a polypeptide comprising residues 165 to 174 of SEQ ID NO:2; a polypeptide comprising residues 179 to 204 of SEQ ID NO:2; a polypeptide comprising residues 245 to 260 of SEQ ID NO:2; a polypeptide comprising residues 1 to 13 of SEQ



ID NO:2; a polypeptide comprising residues 21 to 38 of SEQ ID NO:2; a polypeptide comprising residues 44 to 59 of SEQ ID NO:2; a polypeptide comprising residues 88 to 99 of SEQ ID NO:2; a polypeptide comprising residues 165 to 173 of SEQ ID NO:2; a polypeptide comprising residues 194 to 205 of SEQ ID NO:2; and a polypeptide comprising residues 213 to 221 of SEQ ID NO:2. Within another embodiment, the antibody is generated to a polypeptide selected from the group consisting of: a polypeptide consisting of residues 4 to 13 of SEQ ID NO:2; a polypeptide consisting of residues 17 to 39 of SEQ ID NO:2; a polypeptide consisting of residues 44 to 60 of SEQ ID NO:2; a polypeptide consisting of residues 69 to 104 of SEQ ID NO:2; a polypeptide consisting of residues 106 to 115 of SEQ ID NO:2; a polypeptide consisting of residues 128 to 136 of SEQ ID NO:2; a polypeptide consisting of residues 165 to 174 of SEQ ID NO:2; a polypeptide consisting of residues 179 to 204 of SEQ ID NO:2; a polypeptide consisting of residues 245 to 260 of SEQ ID NO:2; a polypeptide consisting of residues 1 to 13 of SEQ ID NO:2; a polypeptide consisting of residues 21 to 38 of SEQ ID NO:2; a polypeptide consisting of residues 44 to 59 of SEQ ID NO:2; a polypeptide consisting of residues 88 to 99 of SEQ ID NO:2; a polypeptide consisting of residues 165 to 173 of SEQ ID NO:2; a polypeptide consisting of residues 194 to 205 of SEQ ID NO:2; and p) a polypeptide consisting of residues 213 to 221 of SEQ ID NO:2.

Within another aspect of the invention is provided a method of detecting inflamed tissue in lung carcinoma or breast carcinoma comprising, contacting said carcinoma with a polynucleotide consisting of the polynucleotide sequence as shown in SEQ ID NO:1, wherein the polynucleotide hybridizes to the nucleic acid in the inflamed tissue. Within an embodiment, the polynucleotide is selected from the group consisting of: a polynucleotide consisting of at least 16 contiguous nucleotides as shown in SEQ ID NO:1; a polynucleotide consisting of from 17 to 25 contiguous nucleotides as shown in SEQ ID NO:1; a polynucleotide consisting of 40 contiguous nucleotides as shown in SEQ ID NO:1; a polynucleotide consisting of 60 contiguous nucleotides as shown in SEQ ID NO:1; a polynucleotide selected from the group consisting of: the polynucleotide as shown in SEQ ID NO:10; the polynucleotide as shown in SEQ ID NO:11; the polynucleotide as shown in SEQ ID NO:12; the polynucleotide as shown in SEQ ID NO:13; the polynucleotide as shown in SEQ ID NO:23; the polynucleotide as

shown in SEQ ID NO:24; the polynucleotide as shown in SEQ ID NO:25; and the polynucleotide as shown in SEQ ID NO:26; a polynucleotide encoding a polypeptide, wherein the polypeptide comprises residues 139 to 297 of SEQ ID NO:27, or residues 139 to 267 of SEQ ID NO:29.

- 5                    Within another aspect, the invention provides a method of detecting inflamed tissue in lung carcinoma or breast carcinoma comprising, contacting said carcinoma with an antibody to the polypeptide as shown in SEQ ID NO:2. Within an embodiment, the antibody is generated to a polypeptide selected from the group consisting of: a polypeptide comprising residues 4 to 13 of SEQ ID NO:2; a polypeptide
- 10 comprising residues 17 to 39 of SEQ ID NO:2; a polypeptide comprising residues 44 to 60 of SEQ ID NO:2; a polypeptide comprising residues 69 to 104 of SEQ ID NO:2; a polypeptide comprising residues 106 to 115 of SEQ ID NO:2; a polypeptide comprising residues 128 to 136 of SEQ ID NO:2; a polypeptide comprising residues 165 to 174 of SEQ ID NO:2; a polypeptide comprising residues 179 to 204 of SEQ ID NO:2; a polypeptide
- 15 comprising residues 245 to 260 of SEQ ID NO:2; a polypeptide comprising residues 1 to 13 of SEQ ID NO:2; a polypeptide comprising residues 21 to 38 of SEQ ID NO:2; a polypeptide comprising residues 44 to 59 of SEQ ID NO:2; a polypeptide comprising residues 88 to 99 of SEQ ID NO:2; a polypeptide comprising residues 165 to 173 of SEQ ID NO:2; a polypeptide comprising residues 194 to 205 of SEQ ID NO:2; and a
- 20 polypeptide comprising residues 213 to 221 of SEQ ID NO:2. Within another aspect, the antibody is generated to a polypeptide selected from the group consisting of: a polypeptide consisting of residues 4 to 13 of SEQ ID NO:2; a polypeptide consisting of residues 17 to 39 of SEQ ID NO:2; a polypeptide consisting of residues 44 to 60 of SEQ ID NO:2; a polypeptide consisting of residues 69 to 104 of SEQ ID NO:2; a polypeptide
- 25 consisting of residues 106 to 115 of SEQ ID NO:2; a polypeptide consisting of residues 128 to 136 of SEQ ID NO:2; a polypeptide consisting of residues 165 to 174 of SEQ ID NO:2; a polypeptide consisting of residues 179 to 204 of SEQ ID NO:2; a polypeptide consisting of residues 245 to 260 of SEQ ID NO:2; a polypeptide consisting of residues 1 to 13 of SEQ ID NO:2; a polypeptide consisting of residues 21 to 38 of SEQ ID NO:2;
- 30 a polypeptide consisting of residues 44 to 59 of SEQ ID NO:2; a polypeptide consisting of residues 88 to 99 of SEQ ID NO:2; a polypeptide consisting of residues 165 to 173 of

SEQ ID NO:2; a polypeptide consisting of residues 194 to 205 of SEQ ID NO:2; and a polypeptide consisting of residues 213 to 221 of SEQ ID NO:2.

Within another aspect, the invention provides a method of modulating the quantity of cells expressing a polypeptide selected from the group consisting of: a  
5 polypeptide comprising the amino acid sequence of SEQ ID NO:1; a polypeptide comprising the amino acid sequence of SEQ ID NO:27; and a polypeptide comprising the amino acid sequence of SEQ ID NO:29, comprising contacting the cells with an antibody generated to the polypeptide as shown in the amino acid sequence of SEQ ID NO:2.

10 Within another aspect, the invention provides a method of modulating the quantity of cells expressing a polypeptide selected from the group consisting of: a polypeptide comprising the amino acid sequence of SEQ ID NO:1; a polypeptide comprising the amino acid sequence of SEQ ID NO:27; and a polypeptide comprising the amino acid sequence of SEQ ID NO:29, comprising administering to the cells an  
15 isolated polypeptide wherein the isolated polypeptide consists of residues 1 to X of SEQ ID NO:2, wherein X is an integer between 129 and 136.

Within another aspect, the invention provides a method of modulating receptor activation in cells comprising administering to the cells an isolated polypeptide, wherein the isolated polypeptide comprises an amino acid as shown in SEQ ID NO:2  
20 from residue 1 to residue X, wherein X is an integer from 129 to 136. Within an embodiment, the cells express a polypeptide sequence selected from the group consisting of: a polypeptide comprising the amino acid sequence of SEQ ID NO:1; a polypeptide comprising the amino acid sequence of SEQ ID NO:27; and a polypeptide comprising the amino acid sequence of SEQ ID NO:29. Within another embodiment, the isolated  
25 polypeptide consists of an amino acid as shown in SEQ ID NO:2 from residue 1 to residue X.

Within another aspect of the invention is provided a method of modulating proliferation of carcinoma cells comprising administering to the cells an isolated polypeptide wherein the isolated polypeptide comprises an amino acid as shown  
30 in SEQ ID NO:2 from residue 1 to X, wherein X is an integer from 129 to 136. Within an embodiment, the carcinoma cells are lung carcinoma or breast carcinoma cells.

Within another embodiment, the isolated polypeptide consists of an amino acid as shown in SEQ ID NO:2 from residue 1 to X.

Within another aspect, the invention provides a method of modulating proliferation of inflammatory cells comprising administering to the cells an isolated polypeptide wherein the isolated polypeptide comprises an amino acid as shown in SEQ ID NO:2 from residue 1 to X, wherein X is an integer from 129 to 136. Within an embodiment, the isolated polypeptide consists of an amino acid as shown in SEQ ID NO:2 from residue 1 to X.

Within another aspect, the invention provides a method of modulating apoptosis of carcinoma cells comprising administering to the cells an isolated polypeptide wherein the isolated polypeptide comprises an amino acid as shown in SEQ ID NO:2 from residue 1 to X, wherein X is an integer from 129 to 136. Within an embodiment, the carcinoma cells are lung carcinoma or breast carcinoma cells. Within another embodiment, the isolated polypeptide consists of an amino acid as shown in SEQ ID NO:2 from residue 1 to X.

Within another aspect, the invention provides a method of modulating apoptosis of inflammatory cells comprising administering to the cells an isolated polypeptide wherein the isolated polypeptide comprises an amino acid as shown in SEQ ID NO:2 from residue 1 to X, wherein X is an integer from 129 to 136. Within an embodiment, the isolated polypeptide consists of an amino acid as shown in SEQ ID NO:2 from residue 1 to X.

These and other aspects of the invention will become evident upon reference to the following detailed description of the invention and attached drawings.

## DETAILED DESCRIPTION OF THE INVENTION

Prior to setting forth the invention in detail, it may be helpful to the understanding thereof to define the following terms:

The term “affinity tag” is used herein to denote a polypeptide segment that can be attached to a second polypeptide to provide for purification of the second polypeptide or provide sites for attachment of the second polypeptide to a substrate. In

principal, any peptide or protein for which an antibody or other specific binding agent is available can be used as an affinity tag. Affinity tags include a poly-histidine tract, protein A (Nilsson et al., *EMBO J.* 4:1075, 1985; Nilsson et al., *Methods Enzymol.* 198:3, 1991), glutathione S transferase (Smith and Johnson, *Gene* 67:31, 1988), Glu-Glu  
5 affinity tag (Grussenmeyer et al., *Proc. Natl. Acad. Sci. USA* 82:7952-4, 1985) (SEQ ID NO:7), substance P, Flag™ peptide (Hopp et al., *Biotechnology* 6:1204-1210, 1988), streptavidin binding peptide, maltose binding protein (Guan et al., *Gene* 67:21-30, 1987), cellulose binding protein, thioredoxin, ubiquitin, T7 polymerase, or other antigenic epitope or binding domain. See, in general, Ford et al., *Protein Expression and*  
10 *Purification* 2: 95-107, 1991. DNAs encoding affinity tags and other reagents are available from commercial suppliers (e.g., Pharmacia Biotech, Piscataway, NJ; New England Biolabs, Beverly, MA; Eastman Kodak, New Haven, CT).

The terms “amino-terminal” and “carboxyl-terminal” are used herein to denote positions within polypeptides. Where the context allows, these terms are used  
15 with reference to a particular sequence or portion of a polypeptide to denote proximity or relative position. For example, a certain sequence positioned carboxyl-terminal to a reference sequence within a polypeptide is located proximal to the carboxyl terminus of the reference sequence, but is not necessarily at the carboxyl terminus of the complete polypeptide.

20 The term “complements of a polynucleotide molecule” is a polynucleotide molecule having a complementary base sequence and reverse orientation as compared to a reference sequence. For example, the sequence 5' ATGCACGGG 3' is complementary to 5' CCCGTGCAT 3'.

The term “corresponding to”, when applied to positions of amino acid  
25 residues in sequences, means corresponding positions in a plurality of sequences when the sequences are optimally aligned.

The term “degenerate nucleotide sequence” denotes a sequence of nucleotides that includes one or more degenerate codons (as compared to a reference polynucleotide molecule that encodes a polypeptide). Degenerate codons contain  
30 different triplets of nucleotides, but encode the same amino acid residue (i.e., GAU and GAC triplets each encode Asp).

The term "expression vector" is used to denote a DNA molecule, linear or circular, that comprises a segment encoding a polypeptide of interest operably linked to additional segments that provide for its transcription. Such additional segments include promoter and terminator sequences, and may also include one or more origins of replication, one or more selectable markers, an enhancer, a polyadenylation signal, etc. Expression vectors are generally derived from plasmid or viral DNA, or may contain elements of both.

The term "isolated", when applied to a polynucleotide, denotes that the polynucleotide has been removed from its natural genetic milieu and is thus free of other extraneous or unwanted coding sequences, and is in a form suitable for use within genetically engineered protein production systems. Such isolated molecules are those that are separated from their natural environment and include cDNA and genomic clones. Isolated DNA molecules of the present invention are free of other genes with which they are ordinarily associated, but may include naturally occurring 5' and 3' untranslated regions such as promoters and terminators. The identification of associated regions will be evident to one of ordinary skill in the art (see for example, Dynan and Tijan, Nature 316:774-78, 1985).

An "isolated" polypeptide or protein is a polypeptide or protein that is found in a condition other than its native environment, such as apart from blood and animal tissue. In a preferred form, the isolated polypeptide is substantially free of other polypeptides, particularly other polypeptides of animal origin. It is preferred to provide the polypeptides in a highly purified form, i.e. greater than 95% pure, more preferably greater than 99% pure. When used in this context, the term "isolated" does not exclude the presence of the same polypeptide in alternative physical forms, such as dimers or alternatively glycosylated or derivatized forms.

"Operably linked" means that two or more entities are joined together such that they function in concert for their intended purposes. When referring to DNA segments, the phrase indicates, for example, that coding sequences are joined in the correct reading frame, and transcription initiates in the promoter and proceeds through the coding segment(s) to the terminator. When referring to polypeptides, "operably linked" includes both covalently (e.g., by disulfide bonding) and non-covalently (e.g., by

hydrogen bonding, hydrophobic interactions, or salt-bridge interactions) linked sequences, wherein the desired function(s) of the sequences are retained.

The term "ortholog" or "species homolog", denotes a polypeptide or protein obtained from one species that is the functional counterpart of a polypeptide or protein from a different species. Sequence differences among orthologs are the result of speciation.

A "polynucleotide" is a single- or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. Polynucleotides include RNA and DNA, and may be isolated from natural sources, synthesized *in vitro*, or prepared from a combination of natural and synthetic molecules. Sizes of polynucleotides are expressed as base pairs (abbreviated "bp"), nucleotides ("nt"), or kilobases ("kb"). Where the context allows, the latter two terms may describe polynucleotides that are single-stranded or double-stranded. When the term is applied to double-stranded molecules it is used to denote overall length and will be understood to be equivalent to the term "base pairs". It will be recognized by those skilled in the art that the two strands of a double-stranded polynucleotide may differ slightly in length and that the ends thereof may be staggered as a result of enzymatic cleavage; thus all nucleotides within a double-stranded polynucleotide molecule may not be paired. Such unpaired ends will in general not exceed 20 nt in length.

A "polypeptide" is a polymer of amino acid residues joined by peptide bonds, whether produced naturally or synthetically. Polypeptides of less than about 10 amino acid residues are commonly referred to as "peptides".

A "peptide-receptor complex" is formed when a peptide, or ligand, binds to a receptor resulting in a change in the properties of the receptor. This change can result in an initiation of sequences of reactions leading to a change in cellular function, or the inability of the receptor to bind additional peptides. The forming of a peptide-receptor complex can be reversible.

The term "promoter" is used herein for its art-recognized meaning to denote a portion of a gene containing DNA sequences that provide for the binding of RNA polymerase and initiation of transcription. Promoter sequences are commonly, but not always, found in the 5' non-coding regions of genes.

A "protein" is a macromolecule comprising one or more polypeptide chains. A protein may also comprise non-peptidic components, such as carbohydrate groups. Carbohydrates and other non-peptidic substituents may be added to a protein by the cell in which the protein is produced, and will vary with the type of cell. Proteins are defined herein in terms of their amino acid backbone structures; substituents such as carbohydrate groups are generally not specified, but may be present nonetheless.

The term "receptor" denotes a cell-associated protein that binds to a bioactive molecule (i.e., a ligand) and mediates the effect of the ligand on the cell. Membrane-bound receptors are characterized by a multi-domain or multi-peptide structure comprising an extracellular ligand-binding domain and an intracellular effector domain that is typically involved in signal transduction. Binding of ligand to receptor results in a conformational change in the receptor that causes an interaction between the effector domain and other molecule(s) in the cell. This interaction in turn leads to an alteration in the metabolism of the cell. Metabolic events that are linked to receptor-ligand interactions include gene transcription, phosphorylation, dephosphorylation, increases in cyclic AMP production, mobilization of cellular calcium, mobilization of membrane lipids, cell adhesion, hydrolysis of inositol lipids and hydrolysis of phospholipids. In general, receptors can be membrane bound, cytosolic or nuclear; monomeric (e.g., thyroid stimulating hormone receptor, beta-adrenergic receptor) or multimeric (e.g., PDGF receptor, growth hormone receptor, IL-3 receptor, GM-CSF receptor, G-CSF receptor, erythropoietin receptor and IL-6 receptor).

The term "secretory signal sequence" denotes a DNA sequence that encodes a polypeptide (a "secretory peptide") that, as a component of a larger polypeptide, directs the larger polypeptide through a secretory pathway of a cell in which it is synthesized. The larger polypeptide is commonly cleaved to remove the secretory peptide during transit through the secretory pathway.

A "segment" is a portion of a larger molecule (e.g., polynucleotide or polypeptide) having specified attributes. For example, a DNA segment encoding a specified polypeptide is a portion of a longer DNA molecule, such as a plasmid or plasmid fragment, that, when read from the 5' to the 3' direction, encodes the sequence of amino acids of the specified polypeptide.



A "soluble receptor" is a receptor polypeptide that is not bound to a cell membrane. Soluble receptors are most commonly ligand-binding receptor polypeptides that lack transmembrane and cytoplasmic domains. Soluble receptors can comprise additional amino acid residues, such as affinity tags that provide for purification of the polypeptide or provide sites for attachment of the polypeptide to a substrate. Many cell-surface receptors have naturally occurring, soluble counterparts that are produced by proteolysis or translated from alternatively spliced mRNAs. Receptor polypeptides are said to be substantially free of transmembrane and intracellular polypeptide segments when they lack sufficient portions of these segments to provide membrane anchoring or signal transduction, respectively.

The term "splice variant" is used herein to denote alternative forms of RNA transcribed from a gene. Splice variation arises naturally through use of alternative splicing sites within a transcribed RNA molecule, or less commonly between separately transcribed RNA molecules, and may result in several mRNAs transcribed from the same gene. Splice variants may encode polypeptides having altered amino acid sequence. The term splice variant is also used herein to denote a protein encoded by a splice variant of an mRNA transcribed from a gene.

Molecular weights and lengths of polymers determined by imprecise analytical methods (e.g., gel electrophoresis) will be understood to be approximate values. When such a value is expressed as "about" X or "approximately" X, the stated value of X will be understood to be accurate to  $\pm 10\%$ .

All references cited herein are incorporated by reference in their entirety.

The present invention is based in part upon the discovery of a novel DNA sequences (SEQ ID NOs:1) and corresponding polypeptide sequences (SEQ ID NOs:2) which have homology to members of the tumor necrosis factor receptor family. The receptor has been designated (Uterine Myometrium Leiomyoma Receptor) and is also referred to herein as ztnfr11. Novel UMLR receptor-encoding polynucleotides and polypeptides of the present invention were identified by the discovery of a novel cDNA having homology to tumor necrosis factor receptor family members. Sequence analysis of a deduced amino acid sequence indicates the presence of 2 complete extracellular, cysteine-rich pseudo-repeats (# 1, comprises residues 2 to 42, nucleotides 107 to 229; #2

comprises residues 43 to 84 of SEQ ID NO:2, nucleotides 230 to 355 of SEQ ID NO:1), and one partial cysteine-rich pseudo-repeat (#3, comprises residues 85 to approximately 129 of SEQ ID NO:2, nucleotides 356 to approximately 490 of SEQ ID NO:1). Other members of the TNFR superfamily are also known to have partial cysteine-rich repeats.

- 5 These include APO4 and TRAIL. The membrane-bound form further consists of a linker region (approximately residues 130 to 136 of SEQ ID NO:2, nucleotides 491 to 511 of SEQ ID NO:1), a transmembrane domain (residues 137 to 161 of SEQ ID NO:2, nucleotides 512 to 586 of SEQ ID NO:1) and a cytoplasmic, or signaling, domain (residues 162 to 269 of SEQ ID NO:2, nucleotides 587 to 910 of SEQ ID NO:1). The
- 10 mature membrane-bound UMLR protein comprises amino acid residue 1 to 269 of SEQ ID NO:2, nucleotides 104 to 913 of SEQ ID NO:1. Those skilled in the art will recognize that these domain boundaries are approximate, and are based on alignments with known proteins and predictions of protein folding. These features indicate that the receptors encoded by the DNA sequences of SEQ ID NO:1 is a member of the TNF
- 15 receptor family.

Structurally, the TNF receptor family is characterized by an extracellular portion composed of several modules called, historically, "cysteine-rich pseudo-repeats". A prototypical family member has four of these pseudo-repeats, each about 29-43 residues long, one right after the other. A typical pseudo-repeat has 6 cysteine residues.

20 They are called pseudo-repeats because, although they appear to originate from a common ancestral module, they may not repeat exactly. The crystal structure of the p55 TNF receptor revealed that each pseudo-repeat corresponds to one folding domain, and that all four pseudo-repeats fold into the same tertiary structure, held together internally by disulfide bonds.

- 25 Four motifs for the pseudo-repeat sequences are shown below. Within in each motif, X represents an amino acid residue, the numbers contained within curly brackets are multipliers for the preceding residue and residues in brackets are optional. Alternative sequences are indicated with asterisks.

Pseudo-repeat motif #1 (SEQ ID NO:4)

X-C-X{10-14}-C-C-X-X-C-X{5-9}-C-X{6-8}-C-X

Pseudo-repeat motif #2 (SEQ ID NO:5)

5 X-C-X{13-15}-C-X-X-C-X{2-3}-C-X{8-11}-C-X{7}-C

Pseudo-repeat motif #3 (SEQ ID NO:6) and alternative motif (SEQ ID NO:7)

X-C-X{5-6}-X-X{4-9}-C-X-X-C-X{2-7}-C-X{8-9}-C-X{7}-C-(X)

10 \* \*\*\*\*\*  
C X{10-16}

Some TNF receptors, such as OX-40, osteoprotegerin and CD-27, do not contain a complete pseudo-repeat #3 as defined in SEQ ID NOs:6 and 7.

15 Pseudo-repeat motif #4 (SEQ ID NO:8) and alternative motif (SEQ ID NO:9)

X-C-X{10-14}-C-X-X-C-X-X-C-X{4-10}-C-X{3-7}-C-X

\* \*  
X X

20 The pseudo-repeat # 2 (residues 43 to 84 of SEQ ID NO:2) is encompassed by motif # 4 (SEQ ID NO:8). Pseudo-repeats #1(residues 2 to 42 of SEQ ID NO:2) and #3 (residues 85 to approximately 107 of SEQ ID NO:2) most closely resemble motif #4 (SEQ ID NO:8) .

25 Within UMLR, the region corresponding to pseudo-repeat #1 has conserved cysteine residues at amino acid residues 3, 15, 18, 21, 31 and 41 of SEQ ID NO:2. The region corresponding to pseudo-repeat #2 has conserved cysteine residues at amino acid residues 44, 58, 61, 64, 75 and 83 of SEQ ID NO:2. The region corresponding to pseudo-repeat #3 has conserved cysteine residues at amino acid residues 86, 104, 107 and 118 of SEQ ID NO:2.

The cycteine-rich pseudo-repeats are the known ligand binding sites for TNF ligands. Thus, the amino acid residues of UMLR most likely to be involved in ligand binding are likely to be found within residues 1 to 129 of SEQ ID NO:2.

The cysteine-rich domains (pseudo-repeats) of the extracellular ligand binding region of UMLR are similar to several other members of the TNF receptor family. One member of this family is human APO4. See Chaudhary, WO 99/11791. Human APO4 is expressed in the developing fetus as well as adult, and has been characterized as having apoptotic activity, and the ability to activate the JNK signaling pathway. Additionally, mouse APO4 has two isoforms: one with a short cytoplasmic tail, and the other with a longer cytoplasmic tail. These isoforms are the result of alternative splicing. Additional isoforms of the present invention have been identified as well and include one isoform with an insertion as illustrated by the polypeptide sequence as shown in SEQ ID NO:27. As the insertion in this isoform is in the cytoplasmic domain, the polypeptide as shown in SEQ ID NO:27 may be involved in cell signaling.

The other isoform contains this insertion as well as a deletion as illustrated by the polypeptide sequence as shown in SEQ ID NO:29. The deletion in the isoform represented by SEQ ID NO: 29 is in the transmembrane and cytoplasmic domains, and as such may be used as a soluble form of the receptor. An additional sequence identified is SEQ ID NO:38.

OPG is a soluble tumor necrosis factor receptor, which reduces its ligand's concentration in circulation, and regulates osteoclast differentiation and activation (Lacey, D.L., et al., Cell 93: 165-176, 1998). UMLR can be a membrane-bound signaling receptor (residues 1 to 269 of SEQ ID NO:2; residues 1 to 297 of SEQ ID NO:27; or residues 1 to 299 of SEQ ID NO:35) or a soluble receptor (residues 1 to approximately 129 of SEQ ID NO:2) which regulate the amount of circulating ligand. The linker region of UMLR (residues 130 to 136 of SEQ ID NO:2) may not be necessary for binding of the receptor to the ligand. Thus, portions of it may be deleted from a construct for the soluble receptor. Therefore, a construct for a soluble receptor can be as short as residues 1 to 129 of SEQ ID NO:2, or as long as residues 1 to 136 of SEQ ID NO:2. An additional sequences for a soluble ztnfr11 sequences are shown in SEQ ID NO:38.

UMLR can act in a manner analogous to hAPO4 but with an independent ligand. Therefore the ligand for UMLR can function in development, in cellular interactions of the uterus, inflammation, and/ or abnormal cell growth. However it is likely to be homologous to TNF or NGF, or perhaps have a completely different structure from known TNF ligands.

Analysis of the tissue distribution of UMLR RNA was performed by Northern blot techniques. Two transcripts of approximately 3.5 kb and approximately 4.7 kb were detected corresponding to UMLR in melanoma and tumorous lung cell lines. Additional analyses show UMLR RNA in osteosarcoma cell lines and a lymphoma cell line by RT-PCR, and in breast carcinoma and cutaneous lymphoma tissue by *in situ* hybridization. As UMLR polypeptides are expressed in osteosarcoma cell lines, which are derived from osteoblasts, molecules of the present invention may be involved in osteoblast differentiation and activation.

The polynucleotides of the present invention were originally discovered in a uterus, myometrium with leiomyoma library. Expression in tumor cells is consistent with other members of the TNFR family that are associated with growth regulation, differentiation and tumorigenesis.

Broad tissue distribution is also not unknown in the TNF receptor family. Several members, TNFR-I, TNFR-II, TNFR-III and DR4, are found in most human tissues. OPG was detected in lung, heart, kidney, placenta, and to a lesser degree, in hematopoietic and immune organs (Simmonet et al., *ibid.*).

Chromosomal localization of UMLR to Xq11-q12 was determined using radiation hybrid chimeras.

The present invention provides polynucleotide molecules, including DNA and RNA molecules, that encode the UMLR polypeptides disclosed herein. Those skilled in the art will readily recognize that, in view of the degeneracy of the genetic code, considerable sequence variation is possible among these polynucleotide molecules. SEQ ID NO:3 is a degenerate DNA sequence that encompasses all DNAs that encode the UMLR polypeptide of SEQ ID NO:2. SEQ ID NO:28 is a degenerate DNA sequence that encompasses all DNAs that encode the UMLR polypeptide of SEQ ID NO:27. SEQ ID NO:30 is a degenerate DNA sequence that encompasses all DNAs that encode the

UMLR polypeptide of SEQ ID NO:29. SEQ ID NO:39 is a degenerate DNA sequence that encompasses all DNAs that encode the UMLR polypeptide of SEQ ID NO:38. Those skilled in the art will recognize that the degenerate sequence of SEQ ID NOs:3, 28, 30, and 39 also provide all RNA sequences encoding SEQ ID NOs:2, 27, 29, and 38, respectively, by substituting U for T. Thus, UMLR polypeptide-encoding polynucleotides comprising nucleotide 1 to nucleotide 1162 of SEQ ID NO:1, nucleotide 1 to nucleotide 891 of SEQ ID NO:28, nucleotide 1 to nucleotide 801 of SEQ ID NO:30, nucleotide 1 to nucleotide 519 of SEQ ID NO:39, and their RNA equivalents are contemplated by the present invention. Table 1 sets forth the one-letter codes used within SEQ ID Nos:3, 28, 30, and 39 to denote degenerate nucleotide positions. "Resolutions" are the nucleotides denoted by a code letter. "Complement" indicates the code for the complementary nucleotide(s). For example, the code Y denotes either C or T, and its complement R denotes A or G, A being complementary to T, and G being complementary to C.

TABLE 1

Nucleotide	Resolution	Nucleotide	Complement
A	A	T	T
C	C	G	G
G	G	C	C
T	T	A	A
R	A G	Y	C T
Y	C T	R	A G
M	A C	K	G T
K	G T	M	A C
S	C G	S	C G
W	A T	W	A T
H	A C T	D	A G T
B	C G T	V	A C G
V	A C G	B	C G T
D	A G T	H	A C T
N	A C G T	N	A C G T

The degenerate codons used in SEQ ID NOs:3, 28, 30, and 39

5 encompassing all possible codons for a given amino acid, are set forth in Table 2.

TABLE 2

Amino Acid	One Letter Code	Codons	Degenerate Codon
Cys	C	TGC TGT	TGY
Ser	S	AGC AGT TCA TCC TCG TCT	WSN
Thr	T	ACA ACC ACG ACT	CAN
Pro	P	CCA CCC CCG CCT	CCN
Ala	A	GCA GCC GCG GCT	GCN
Gly	G	GGA GGC GGG GGT	GGN
Asn	N	AAC AAT	AA Y
Asp	D	GAC GAT	GAY
Glu	E	GAA GAG	GAR
Gln	Q	CAA CAG	CAR
His	H	CAC CAT	CAY
Arg	R	AGA AGG CGA CGC CGG CGT	MGN
Lys	K	AAA AAG	AAR
Met	M	ATG	ATG
Ile	I	ATA ATC ATT	ATH
Leu	L	CTA CTC CTG CTT TTA TTG	YTN
Val	V	GTA GTC GTG GTT	GTN
Phe	F	TTC TTT	TTY
Tyr	Y	TAC TAT	TAY
Trp	W	TGG	TGG
Ter	.	TAA TAG TGA	TRR
Asn Asp	B		RAY
Glu Gln	Z		SAR
Any	X		NNN



One of ordinary skill in the art will appreciate that some ambiguity is introduced in determining a degenerate codon, representative of all possible codons encoding each amino acid. For example, the degenerate codon for serine (WSN) can, in some circumstances, encode arginine (AGR), and the degenerate codon for arginine (MGN) can, in some circumstances, encode serine (AGY). A similar relationship exists between codons encoding phenylalanine and leucine. Thus, some polynucleotides encompassed by the degenerate sequence may encode variant amino acid sequences, but one of ordinary skill in the art can easily identify such variant sequences by reference to the amino acid sequences of SEQ ID NOs:2, 27, 29, 35, and 38. Variant sequences can be readily tested for functionality as described herein.

The present invention further provides polynucleotide molecules, including DNA and RNA molecules, encoding UMLR proteins. The polynucleotides of the present invention include the sense strand; the anti-sense strand; and the DNA as double-stranded, having both the sense and anti-sense strand annealed together by their respective hydrogen bonds. Representative DNA sequences encoding UMLR proteins are set forth in SEQ ID NOs:1, 3, 22, 28, 30, 31, 32, 33, 36, and 39. DNA sequences encoding other UMLR proteins can be readily generated by those of ordinary skill in the art based on the genetic code.

One of ordinary skill in the art will also appreciate that different species can exhibit "preferential codon usage." Preferential codons for a particular species can be introduced into the polynucleotides of the present invention by a variety of methods known in the art. Introduction of preferential codon sequences into recombinant DNA can, for example, enhance production of the protein by making protein translation more efficient within a particular cell type or species. Therefore, the degenerate codon sequences disclosed in SEQ ID NOs:3, 28, 30, and 39 serve as templates for optimizing expression of polynucleotides in various cell types and species commonly used in the art and disclosed herein. Sequences containing preferential codons can be tested and optimized for expression in various species, and tested for functionality as disclosed herein.

Within preferred embodiments of the invention the isolated polynucleotides will hybridize to similar sized regions of SEQ ID NOs:1, 3, 22, 28, 30,

31, 32, 33, 36, and 39, or a sequence complementary thereto under stringent conditions. Polynucleotide hybridization is well known in the art and widely used for many applications, see for example, Sambrook et al., Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, NY, 1989; Ausubel et al., eds., Current  
 5 Protocols in Molecular Biology, John Wiley and Sons, Inc., NY, 1987; Berger and Kimmel, eds., Guide to Molecular Cloning Techniques, Methods in Enzymology, volume 152, 1987 and Wetmur, Crit. Rev. Biochem. Mol. Biol. 26:227-59, 1990. Polynucleotide hybridization exploits the ability of single stranded complementary sequences to form a double helix hybrid. Such hybrids include DNA-DNA, RNA-RNA  
 10 and DNA-RNA.

As an illustration, a nucleic acid molecule encoding a variant UMLR polypeptide can be hybridized with a nucleic acid molecule having the nucleotide sequence of SEQ ID NOs:1, 3, 22, 28, 30, 31, 32, 33, 36, and 39 (or their complements) at 65°C overnight in ExpressHyb™ Hybridization Solution (CLONTECH Laboratories,  
 15 Inc., Palo Alto, CA). One of skill in the art can devise variations of these hybridization conditions.

Following hybridization, the nucleic acid molecules can be washed to remove non-hybridized nucleic acid molecules under stringent conditions, or under highly stringent conditions. Typical stringent washing conditions include washing in a  
 20 solution of 0.5x - 2x SSC with 0.1% sodium dodecyl sulfate (SDS) at 55 - 65°C. That is, nucleic acid molecules encoding a variant UMLR polypeptide hybridize with a nucleic acid molecule having the nucleotide sequences of SEQ ID NOs:1, 3, 22, 28, 30, 31, 32, 33, 35, 36, and 39 (or their complements) under stringent washing conditions, in which the wash stringency is equivalent to 0.1x - 2x SSC with 0.1% SDS at 55 - 65°C,  
 25 including 0.1x SSC with 0.1% SDS at 55°C, or 2xSSC with 0.1% SDS at 65°C. One of skill in the art can readily devise equivalent conditions, for example, by substituting SSPE for SSC in the wash solution.

The present invention also contemplates UMLR variant nucleic acid molecules that can be identified using two criteria: a determination of the similarity  
 30 between the encoded polypeptides with the amino acid sequences of SEQ ID NOs:2, 27, 29, 34, 35, 37, and 38 (as described below), and a hybridization assay, as described

above. Such UMLR variants include nucleic acid molecules that hybridize with a nucleic acid molecule having the nucleotide sequence of SEQ ID NOs:1, 3, 22, 28, 30, 31, 32, 33, 36, and 39 (or their complements) under stringent washing conditions, in which the wash stringency is equivalent to 0.1x - 2x SSC with 0.1% SDS at 55 - 65°C, and (2) that encode a polypeptide having at least 80%, preferably 90%, more preferably, 95% or greater than 95% sequence identity to the amino acid sequence of SEQ ID NOs:2, 27, 29, 34, 35, 37, and 38. Alternatively, UMLR variants can be characterized as nucleic acid molecules (1) that hybridize with a nucleic acid molecule having the nucleotide sequence of SEQ ID NOs:1, 3, 22, 28, 30, 31, 32, 33, 35, 36, and 39 (or their complements) under highly stringent washing conditions, in which the wash stringency is equivalent to 0.1x - 0.2x SSC with 0.1% SDS at 50 - 65°C, and (2) that encode a polypeptide having at least 80%, preferably 90%, more preferably 95% or greater than 95% sequence identity to the amino acid sequence of SEQ ID NOs:2, 27, 29, 34, 35, 37, and 38.

The highly conserved amino acids in the cysteine-rich pseudo-repeat domains of UMLR can be used as a tool to identify new family members. For instance, reverse transcription-polymerase chain reaction (RT-PCR) can be used to amplify sequences encoding the conserved cysteine-rich domain from RNA obtained from a variety of tissue sources or cell lines. In particular, highly degenerate primers designed from the UMLR sequences are useful for this purpose.

As previously noted, the isolated polynucleotides of the present invention include DNA and RNA. Methods for preparing DNA and RNA are well known in the art. In general, RNA is isolated from a tissue or cell that produces large amounts of UMLR RNA. Such tissues and cells are identified by Northern blotting (Thomas, Proc. Natl. Acad. Sci. USA 77:5201, 1980) and in situ hybridization, and include tissues of the uterus, lung carcinoma and breast carcinoma, as well as cell lines of melanoma, osteosarcoma and lymphoma. Cells lines which express ztnfr11 polynucleotides and polypeptides are available from ATCC (Manassas, VA) and include, but are not limited to ATCC numbers HTB-64, CRL-1585, HTB-68, and CRL-1424.

Total RNA can be prepared using guanidine isothiocyanate extraction followed by isolation by centrifugation in a CsCl gradient (Chirgwin et al., Biochemistry

18:52-94, 1979). Poly (A)<sup>+</sup> RNA is prepared from total RNA using the method of Aviv and Leder (Proc. Natl. Acad. Sci. USA 69:1408-12, 1972). Complementary DNA (cDNA) is prepared from poly(A)<sup>+</sup> RNA using known methods. In the alternative, genomic DNA can be isolated. Polynucleotides encoding UMLR polypeptides are then  
 5 identified and isolated by, for example, hybridization or PCR.

A full-length clone encoding UMLR can be obtained by conventional cloning procedures. Complementary DNA (cDNA) clones are preferred, although for some applications (e.g., expression in transgenic animals) it may be preferable to use a genomic clone, or to modify a cDNA clone to include at least one genomic intron.  
 10 Methods for preparing cDNA and genomic clones are well known and within the level of ordinary skill in the art, and include the use of the sequence disclosed herein, or parts thereof, for probing or priming a library. Expression libraries can be probed with antibodies to UMLR or other specific binding partners.

The invention also provides isolated and purified UMLR polynucleotide  
 15 probes. Such polynucleotide probes can be RNA or DNA. DNA can be either cDNA or genomic DNA. Polynucleotide probes are single or double-stranded DNA or RNA, generally synthetic oligonucleotides, but may be generated from cloned cDNA or genomic sequences and will generally comprise at least 16 nucleotides, more often from 17 nucleotides to 25 or more nucleotides, sometimes 40 to 60 nucleotides, and in some  
 20 instances a substantial portion, domain or even the entire UMLR gene or cDNA. The synthetic oligonucleotides of the present invention have at least 80% identity to a representative UMLR DNA sequence (SEQ ID NOs:1, 3, 22, 28, 30, 31, 32, 33, 36, and 39) or their complements. The invention also provides oligonucleotide probes or primers comprising at least 14 contiguous nucleotides of a polynucleotide of SEQ ID NOs:1, 3,  
 25 22, 28, 30, 31, 32, 33, 36, and 39 or a sequence complementary to SEQ ID NOs:1, 3, 22, 28, 30, 31, 32, 33, 35, 36, and 39.

Preferred regions from which to construct probes include the 5' and/or 3' coding sequences, ligand binding regions, and signal sequences, and the like. Techniques for developing polynucleotide probes and hybridization techniques are  
 30 known in the art, see for example, Ausubel et al., eds., Current Protocols in Molecular Biology, John Wiley and Sons, Inc., NY, 1991. For use as probes, the molecules can be

labeled to provide a detectable signal, such as with an enzyme, biotin, a radionuclide, fluorophore, chemiluminescer, paramagnetic particle and the like, which are commercially available from many sources, such as Molecular Probes, Inc., Eugene, OR, and Amersham Corp., Arlington Heights, IL, using techniques that are well known in the art. Such probes can also be used in hybridizations to detect the presence or quantify the amount of UMLR gene or mRNA transcript in a sample. UMLR polynucleotide probes could be used to hybridize to DNA or RNA targets for diagnostic purposes, using such techniques such as fluorescent *in situ* hybridization (FISH) or immunohistochemistry. Polynucleotide probes can be used to identify genes encoding UMLR-like proteins. For example, UMLR polynucleotides can be used as primers and/or templates in PCR reactions to identify other novel members of the TNFR family. Such probes can also be used to screen libraries for related sequences encoding novel tumor necrosis factor receptors. Such screening would be carried out under conditions of low stringency which would allow identification of sequences which are substantially homologous, but not requiring complete homology to the probe sequence. Such methods and conditions are well known in the art, see, for example, Sambrook et al., Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, NY, 1989. Such low stringency conditions could include hybridization temperatures less than 42°C, formamide concentrations of less than 50% and moderate to low concentrations of salt. Libraries may be made of genomic DNA or cDNA. Polynucleotide probes are also useful for Southern, Northern, or slot blots, colony and plaque hybridization and *in situ* hybridization. Mixtures of different UMLR polynucleotide probes can be prepared which would increase sensitivity or the detection of low copy number targets, in screening systems.

In addition, such polynucleotide probes could be used to hybridize to counterpart sequences on individual chromosomes. Chromosomal identification and/or mapping of the UMLR gene could provide useful information about gene function and disease association. Many mapping techniques are available to one skilled in the art, for example, mapping somatic cell hybrids, and fluorescence *in situ* hybridization (FISH). A preferred method is radiation hybrid mapping. Radiation hybrid mapping is a somatic cell genetic technique developed for constructing high-resolution, contiguous maps of

mammalian chromosomes (Cox et al., Science 250:245-50, 1990). Partial or full knowledge of a gene's sequence allows the designing of PCR primers suitable for use with chromosomal radiation hybrid mapping panels. Commercially available radiation hybrid mapping panels which cover the entire human genome, such as the Stanford G3

5 RH Panel and the GeneBridge 4 RH Panel (Research Genetics, Inc., Huntsville, AL), are available. These panels enable rapid, PCR based, chromosomal localizations and ordering of genes, sequence-tagged sites (STSs), and other non-polymorphic- and polymorphic markers within a region of interest. This includes establishing directly proportional physical distances between newly discovered genes of interest and

10 previously mapped markers. The precise knowledge of a gene's position can be useful in a number of ways including: 1) determining if a sequence is part of an existing contig and obtaining additional surrounding genetic sequences in various forms such as YAC-, BAC- or cDNA clones, 2) providing a possible candidate gene for an inheritable disease which shows linkage to the same chromosomal region, and 3) for cross-referencing

15 model organisms such as mouse which may be beneficial in helping to determine what function a particular gene might have.

UMLR polynucleotide sequences disclosed herein can also be used as probes or primers to clone 5' non-coding regions of a UMLR gene. In view of the tissue-specific expression observed for UMLR by Northern blotting, this gene region is

20 expected to provide for specific expression in tissues of the uterus, lung carcinoma and breast carcinoma, as well as in cell lines of melanoma, osteosarcoma and lymphoma. Promoter elements from a UMLR gene could thus be used to direct the tissue-specific expression of heterologous genes in, for example, transgenic animals or patients treated with gene therapy. Cloning of 5' flanking sequences also facilitates production of

25 UMLR proteins by "gene activation" as disclosed in U.S. Patent No. 5,641,670. Briefly, expression of an endogenous UMLR gene in a cell is altered by introducing into the UMLR locus a DNA construct comprising at least a targeting sequence, a regulatory sequence, an exon, and an unpaired splice donor site. The targeting sequence is a UMLR 5' non-coding sequence that permits homologous recombination of the construct with the

30 endogenous UMLR locus, whereby the sequences within the construct become operably linked with the endogenous UMLR coding sequence. In this way, an endogenous UMLR

promoter can be replaced or supplemented with other regulatory sequences to provide enhanced, tissue-specific, or otherwise regulated expression.

The polynucleotides of the present invention can also be synthesized using DNA synthesizers. Currently the method of choice is the phosphoramidite method.

- 5 If chemically synthesized double stranded DNA is required for an application such as the synthesis of a gene or a gene fragment, then each complementary strand is made separately. The production of short genes (60 to 80 bp) is technically straightforward and can be accomplished by synthesizing the complementary strands and then annealing them. For the production of longer genes (>300 bp), however, special strategies must be
- 10 invoked, because the coupling efficiency of each cycle during chemical DNA synthesis is seldom 100%. To overcome this problem, synthetic genes (double-stranded) are assembled in modular form from single-stranded fragments that are from 20 to 100 nucleotides in length. See Glick and Pasternak, *Molecular Biotechnology, Principles and Applications of Recombinant DNA*, (ASM Press, Washington, D.C. 1994); Itakura *et al.*,
- 15 Annu. Rev. Biochem. 53: 323-356 (1984) and Climie *et al.*, Proc. Natl. Acad. Sci. USA 87:633-7, 1990.

- The present invention further provides counterpart polypeptides and polynucleotides from other species (orthologs). These species include, but are not limited to mammalian, avian, amphibian, reptile, fish, insect and other vertebrate and
- 20 invertebrate species. Of particular interest are UMLR polypeptides from other mammalian species, including murine, porcine, ovine, bovine, canine, feline, equine, and other primate polypeptides. Orthologs of human UMLR can be cloned using information and compositions provided by the present invention in combination with conventional cloning techniques. For example, a cDNA can be cloned using mRNA obtained from a
  - 25 tissue or cell type that expresses UMLR as disclosed herein. Such tissue would include, for example, tissues of the uterus, lung carcinoma, breast carcinoma, as well as in cell lines of melanoma, osteosarcoma, and lymphoma. Suitable sources of mRNA can be identified by probing Northern blots with probes designed from the sequences disclosed herein. A library is then prepared from mRNA of a positive tissue or cell line. A UMLR
  - 30 -encoding cDNA can then be isolated by a variety of methods, such as by probing with a complete or partial human cDNA or with one or more sets of degenerate probes based on

the disclosed sequences. A cDNA can also be cloned using the polymerase chain reaction, or PCR (Mullis, U.S. Patent No. 4,683,202), using primers designed from the representative human UMLR sequences disclosed herein. Within an additional method, the cDNA library can be used to transform or transfect host cells, and expression of the cDNA of interest can be detected with an antibody to UMLR polypeptide. Similar techniques can also be applied to the isolation of genomic clones.

Those skilled in the art will recognize that the sequences disclosed in SEQ ID NO:1 represents a single allele of human UMLR and that allelic variation and alternative splicing are expected to occur. Allelic variants of this sequence can be cloned by probing cDNA or genomic libraries from different individuals according to standard procedures. Allelic variants of the DNA sequences shown in SEQ ID NO:1, including those containing silent mutations and those in which mutations result in amino acid sequence changes, are within the scope of the present invention, as are proteins which are allelic variants of SEQ ID NO:2. cDNAs generated from alternatively spliced mRNAs, which retain the properties of the UMLR polypeptide are included within the scope of the present invention, as are polypeptides encoded by such cDNAs and mRNAs. Allelic variants and splice variants of these sequences can be cloned by probing cDNA or genomic libraries from different individuals or tissues according to standard procedures known in the art.

The present invention also provides isolated UMLR polypeptides that are substantially similar to the polypeptides of SEQ ID NOs:2, 27, 29, 34, 35, 37, and 38 and their orthologs. Such polypeptides will more preferably be at least 90% identical, and more preferably 95% or more identical to SEQ ID NOs:2, 27, 29, 34, 35, 37, and 38 and their orthologs. Percent sequence identity is determined by conventional methods. See, for example, Altschul et al., Bull. Math. Bio. 48: 603-16, 1986 and Henikoff and Henikoff, Proc. Natl. Acad. Sci. USA 89:10915-9, 1992. Briefly, two amino acid sequences are aligned to optimize the alignment scores using a gap opening penalty of 10, a gap extension penalty of 1, and the "blosum 62" scoring matrix of Henikoff and Henikoff (*ibid.*) as shown in Table 3 (amino acids are indicated by the standard one-letter codes). The percent identity is then calculated as:



Total number of identical matches

- x 100

[length of the longer sequence plus the number of gaps introduced into the longer sequence in order to align the two sequences]

5

Year	1954	1955	1956	1957	1958	1959	1960	1961	1962	1963	1964	1965	1966	1967	1968	1969	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100																										
Population	1,000,000	1,050,000	1,100,000	1,150,000	1,200,000	1,250,000	1,300,000	1,350,000	1,400,000	1,450,000	1,500,000	1,550,000	1,600,000	1,650,000	1,700,000	1,750,000	1,800,000	1,850,000	1,900,000	1,950,000	2,000,000	2,050,000	2,100,000	2,150,000	2,200,000	2,250,000	2,300,000	2,350,000	2,400,000	2,450,000	2,500,000	2,550,000	2,600,000	2,650,000	2,700,000	2,750,000	2,800,000	2,850,000	2,900,000	2,950,000	3,000,000	3,050,000	3,100,000	3,150,000	3,200,000	3,250,000	3,300,000	3,350,000	3,400,000	3,450,000	3,500,000	3,550,000	3,600,000	3,650,000	3,700,000	3,750,000	3,800,000	3,850,000	3,900,000	3,950,000	4,000,000	4,050,000	4,100,000	4,150,000	4,200,000	4,250,000	4,300,000	4,350,000	4,400,000	4,450,000	4,500,000	4,550,000	4,600,000	4,650,000	4,700,000	4,750,000	4,800,000	4,850,000	4,900,000	4,950,000	5,000,000	5,050,000	5,100,000	5,150,000	5,200,000	5,250,000	5,300,000	5,350,000	5,400,000	5,450,000	5,500,000	5,550,000	5,600,000	5,650,000	5,700,000	5,750,000	5,800,000	5,850,000	5,900,000	5,950,000	6,000,000	6,050,000	6,100,000	6,150,000	6,200,000	6,250,000	6,300,000	6,350,000	6,400,000	6,450,000	6,500,000	6,550,000	6,600,000	6,650,000	6,700,000	6,750,000	6,800,000	6,850,000	6,900,000	6,950,000	7,000,000	7,050,000	7,100,000	7,150,000	7,200,000	7,250,000	7,300,000	7,350,000	7,400,000	7,450,000	7,500,000	7,550,000	7,600,000	7,650,000	7,700,000	7,750,000	7,800,000	7,850,000	7,900,000	7,950,000	8,000,000	8,050,000	8,100,000	8,150,000	8,200,000	8,250,000	8,300,000	8,350,000	8,400,000	8,450,000	8,500,000	8,550,000	8,600,000	8,650,000	8,700,000	8,750,000	8,800,000	8,850,000	8,900,000	8,950,000	9,000,000	9,050,000	9,100,000	9,150,000	9,200,000	9,250,000	9,300,000	9,350,000	9,400,000	9,450,000	9,500,000	9,550,000	9,60



Sequence identity of polynucleotide molecules is determined by similar methods using a ratio as disclosed above.

Those skilled in the art appreciate that there are many established algorithms available to align two amino acid sequences. The “FASTA” similarity search  
 5 algorithm of Pearson and Lipman is a suitable protein alignment method for examining the level of identity shared by an amino acid sequence disclosed herein and the amino acid sequence of a putative variant UMLR . The FASTA algorithm is described by Pearson and Lipman, Proc. Nat’l Acad. Sci. USA 85:2444 (1988), and by Pearson, Meth. Enzymol. 183:63 (1990).

10 Briefly, FASTA first characterizes sequence similarity by identifying regions shared by the query sequence (*e.g.*, SEQ ID NOs:2, 27, 29, 34, 35, 37, and 38) and a test sequence that have either the highest density of identities (if the ktup variable is 1) or pairs of identities (if ktup=2), without considering conservative amino acid substitutions, insertions, or deletions. The ten regions with the highest density of  
 15 identities are then rescored by comparing the similarity of all paired amino acids using an amino acid substitution matrix, and the ends of the regions are “trimmed” to include only those residues that contribute to the highest score. If there are several regions with scores greater than the “cutoff” value (calculated by a predetermined formula based upon the length of the sequence and the ktup value), then the trimmed initial regions are  
 20 examined to determine whether the regions can be joined to form an approximate alignment with gaps. Finally, the highest scoring regions of the two amino acid sequences are aligned using a modification of the Needleman-Wunsch-Sellers algorithm (Needleman and Wunsch, J. Mol. Biol. 48:444 (1970); Sellers, SIAM J. Appl. Math. 26:787 (1974)), which allows for amino acid insertions and deletions. Illustrative  
 25 parameters for FASTA analysis are: ktup=1, gap opening penalty=10, gap extension penalty=1, and substitution matrix=BLOSUM62. These parameters can be introduced into a FASTA program by modifying the scoring matrix file (“SMATRIX”), as explained in Appendix 2 of Pearson, Meth. Enzymol. 183:63 (1990).

FASTA can also be used to determine the sequence identity of nucleic  
 30 acid molecules using a ratio as disclosed above. For nucleotide sequence comparisons, the ktup value can range between one to six, preferably from four to six.

The present invention includes nucleic acid molecules that encode a polypeptide having one or more conservative amino acid changes, compared with the amino acid sequences of SEQ ID NOs:2, 27, 29, 34, 35, 37, and 38. The BLOSUM62 table is an amino acid substitution matrix derived from about 2,000 local multiple  
5 alignments of protein sequence segments, representing highly conserved regions of more than 500 groups of related proteins (Henikoff and Henikoff, Proc. Nat'l Acad. Sci. USA 89:10915 (1992)). Accordingly, the BLOSUM62 substitution frequencies can be used to define conservative amino acid substitutions that may be introduced into the amino acid sequences of the present invention. As used herein, the language "conservative amino  
10 acid substitution" refers to a substitution represented by a BLOSUM62 value of greater than -1. For example, an amino acid substitution is conservative if the substitution is characterized by a BLOSUM62 value of 0, 1, 2, or 3. Preferred conservative amino acid substitutions are characterized by a BLOSUM62 value of at least 1 (e.g., 1, 2 or 3), while more preferred conservative amino acid substitutions are characterized by a BLOSUM62  
15 value of at least 2 (e.g., 2 or 3).

Conservative amino acid changes in an UMLR gene can be introduced by substituting nucleotides for the nucleotides recited in SEQ ID NOs:1, 3, 22, 28, 30, 31, 32, 33, 35, 36, and 39. Such "conservative amino acid" variants can be obtained, for example, by oligonucleotide-directed mutagenesis, linker-scanning mutagenesis,  
20 mutagenesis using the polymerase chain reaction, and the like (see Ausubel (1995) at pages 8-10 to 8-22; and McPherson (ed.), *Directed Mutagenesis: A Practical Approach* (IRL Press 1991)). The ability of such variants to modulate cell-cell interactions, apoptosis, and inflammation can be determined using a standard method, such as the assay described herein. Alternatively, a variant UMLR polypeptide can be identified by  
25 the ability to specifically bind anti-UMLR antibodies.

Essential amino acids in the polypeptides of the present invention can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, Science 244: 1081-5, 1989; Bass et al., Proc. Natl. Acad. Sci. USA 88:4498-502, 1991). In the latter technique, single  
30 alanine mutations are introduced at every residue in the molecule, and the resultant mutant molecules are tested for biological activity as disclosed below to identify amino

acid residues that are critical to the activity of the molecule. See also, Hilton et al., J. Biol. Chem. 271:4699-708, 1996. Sites of receptor-ligand interaction can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction or photoaffinity labeling, in  
 5 conjunction with mutation of putative contact site amino acids. See, for example, de Vos et al., Science 255:306-12, 1992; Smith et al., J. Mol. Biol. 224:899-904, 1992; Wlodaver et al., FEBS Lett. 309:59-64, 1992. The identities of essential amino acids can also be inferred from analysis of homologies with related TNFR-like molecules.

Multiple amino acid substitutions can be made and tested using known  
 10 methods of mutagenesis and screening, such as those disclosed by Reidhaar-Olson and Sauer (Science 241:53-7, 1988) or Bowie and Sauer (Proc. Natl. Acad. Sci. USA 86:2152-6, 1989). Briefly, these authors disclose methods for simultaneously randomizing two or more positions in a polypeptide, selecting for functional polypeptide, and then sequencing the mutagenized polypeptides to determine the spectrum of  
 15 allowable substitutions at each position. Other methods that can be used include phage display (e.g., Lowman et al., Biochem. 30:10832-7, 1991; Ladner et al., U.S. Patent No. 5,223,409; Huse, WIPO Publication WO 92/06204) and region-directed mutagenesis (Derbyshire et al., Gene 46:145, 1986; Ner et al., DNA 7:127, 1988).

Variants of the disclosed UMLR DNA and polypeptide sequences can be  
 20 generated through DNA shuffling, as disclosed by Stemmer, Nature 370:389-91, 1994, Stemmer, Proc. Natl. Acad. Sci. USA 91:10747-51, 1994 and WIPO Publication WO 97/20078. Briefly, variant DNAs are generated by *in vitro* homologous recombination by random fragmentation of a parent DNA followed by reassembly using PCR, resulting in randomly introduced point mutations. This technique can be modified by using a  
 25 family of parent DNAs, such as allelic variants or DNAs from different species, to introduce additional variability into the process. Selection or screening for the desired activity, followed by additional iterations of mutagenesis and assay provides for rapid "evolution" of sequences by selecting for desirable mutations while simultaneously selecting against detrimental changes.

30 Mutagenesis methods as disclosed herein can be combined with high-throughput, automated screening methods to detect activity of cloned, mutagenized

polypeptides in host cells. Mutagenized DNA molecules that encode active polypeptides (e.g., ligand binding activity) can be recovered from the host cells and rapidly sequenced using modern equipment. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide of interest, and can be  
5 applied to polypeptides of unknown structure.

Regardless of the particular nucleotide sequence of a variant UMLR gene, the gene encodes a polypeptide that is characterized by its cell-cell interaction activity, including but not limited to apoptosis, and tumorigensis, or by the ability to bind specifically to an anti-UMLR antibody. More specifically, variant UMLR genes encode  
10 polypeptides which exhibit at least 50%, and preferably, greater than 70, 80, or 90%, of the activity of polypeptide encoded by the human UMLR gene described herein.

Variant UMLR polypeptides or substantially homologous UMLR polypeptides are characterized as having one or more amino acid substitutions, deletions or additions. These changes are preferably of a minor nature, that is conservative amino  
15 acid substitutions and other substitutions that do not significantly affect the folding or activity of the polypeptide; small deletions, typically of one to about 30 amino acids; and amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue, a small linker peptide of up to about 20-25 residues, or an affinity tag. The present invention thus includes polypeptides of from 40 to 2000 amino acid residues that  
20 comprise a sequence that is at least 85%, preferably at least 90%, and more preferably 95% or more identical to the corresponding region of SEQ ID NOs:2, 27, 29, 34, 35, 37, and 38. Polypeptides comprising affinity tags can further comprise a proteolytic cleavage site between the UMLR polypeptide and the affinity tag. Preferred such sites include thrombin cleavage sites and factor Xa cleavage sites.

25 For any UMLR polypeptide, including variants and fusion proteins, one of ordinary skill in the art can readily generate a fully degenerate polynucleotide sequence encoding that variant using the information set forth in Tables 1 and 2 above. Moreover, those of skill in the art can use standard software to devise UMLR variants based upon the nucleotide and amino acid sequences described herein. Accordingly, the  
30 present invention includes a computer-readable medium encoded with a data structure that provides at least one of the following sequences: SEQ ID NOs:1, 2, 3, 22, 27, 28,

29, 30, 31, 32, 33, 34, 35, 36, 37, 38, and 39. Suitable forms of computer-readable media include magnetic media and optically-readable media. Examples of magnetic media include a hard or fixed drive, a random access memory (RAM) chip, a floppy disk, digital linear tape (DLT), a disk cache, and a ZIP disk. Optically readable media are exemplified by compact discs (*e.g.*, CD-read only memory (ROM), CD-rewritable (RW), and CD-recordable), and digital versatile/video discs (DVD) (*e.g.*, DVD-ROM, DVD-RAM, and DVD+RW).

The present invention further provides a variety of other polypeptide fusions and related multimeric proteins comprising one or more polypeptide fusions. For example, the cysteine-rich pseudo-repeat, or cytoplasmic polypeptide domains can be prepared as a fusion to a dimerizing protein, as disclosed in U.S. Patents Nos. 5,155,027 and 5,567,584. Preferred dimerizing proteins in this regard include other cysteine-rich pseudo-repeat or cytoplasmic polypeptide domains, or polypeptides comprising other members of the TNF receptor family of proteins, such as, for example, APO4, TNFR-I, TNFR-II, and TNFR-III. Additionally, chimeras can be prepared with the extracellular portion of cytokine receptors. For example, the extracellular domain of erythropoietin can be fused to the linker, transmembrane, and cytoplasmic domain of UMLR polypeptides to produce dimerization. These polypeptide domain fusions, can be expressed in genetically engineered cells to produce a variety of multimeric TNFR-like analogs.

Fusion proteins can be prepared by methods known to those skilled in the art by preparing each component of the fusion protein and chemically conjugating them. Alternatively, a polynucleotide encoding both components of the fusion protein in the proper reading frame can be generated using known techniques and expressed by the methods described herein. For example, part or all of a domain(s) conferring a biological function may be swapped between UMLR of the present invention with the functionally equivalent domain(s) from another family member, such as APO4, TNFR-I, TNFR-II, and TNFR-III. Such domains include, but are not limited to, conserved motifs such as the cysteine-rich pseudo-repeat, transmembrane, and cytoplasmic domains. Such fusion proteins would be expected to have a biological functional profile that is the same or similar to polypeptides of the present invention or other known TNFR family proteins

(e.g. APO4, TNFR-I, TNFR-II, and TNFR-III), depending on the fusion constructed. Moreover, such fusion proteins may exhibit other properties as disclosed herein.

Moreover, using methods described in the art, polypeptide fusions, or hybrid UMLR proteins, are constructed using regions or domains of the inventive UMLR  
 5 in combination with those of other TNFR molecules. (e.g. APO4, TNFR-I, TNFR-II, and TNFR-III), or heterologous proteins (Sambrook et al., ibid., Altschul et al., ibid., Picard, Cur. Opin. Biology, 5:511-5, 1994, and references therein). These methods allow the determination of the biological importance of larger domains or regions in a polypeptide of interest. Such hybrids may alter reaction kinetics, binding, constrict or expand the  
 10 substrate specificity, or alter tissue and cellular localization of a polypeptide, and can be applied to polypeptides of unknown structure.

Auxiliary domains can be fused to UMLR polypeptides to target them to specific cells, tissues, or macromolecules (e.g., tissues of the uterus, lung carcinoma and breast carcinoma, as well as in cells of melanoma, osteosarcoma, and lymphoma). For  
 15 example, a protease domain could be targeted to a predetermined cell type (uterus, melanoma, osteosarcoma, breast carcinoma, and lymphoma, and in tumorous lung) by fusing it to a the cysteine-rich pseudo-repeat domains (i.e., residues 1 to 129 of SEQ ID NO:2, or residues 1 to 173 of SEQ ID NO:38), or a portion thereof which has been shown to bind the cognate ligand for UMLR. In this way, polypeptides, polypeptide  
 20 fragments and proteins can be targeted for therapeutic or diagnostic purposes. Such the cysteine-rich pseudo-repeat domains, or portions thereof can be fused to two or more moieties, such as an affinity tag for purification and a targeting domains. Polypeptide fusions can also comprise one or more cleavage sites, particularly between domains. See, Tuan et al., Connective Tissue Research 34:1-9, 1996.

25 Polypeptide fusions of the present invention will generally contain not more than about 1,500 amino acid residues, preferably not more than about 1,200 residues, more preferably not more than about 1,000 residues, and will in many cases be considerably smaller. For example, residues of UMLR polypeptide can be fused to E. coli  $\beta$ -galactosidase (1,021 residues; see Casadaban et al., J. Bacteriol. 143:971-980,  
 30 1980), a 10-residue spacer, and a 4-residue factor Xa cleavage site. In a second example,



residues of UMLR polypeptide can be fused to maltose binding protein (approximately 370 residues), a 4-residue cleavage site, and a 6-residue polyhistidine tag.

UMLR polypeptides or fragments thereof may also be prepared through chemical synthesis. UMLR polypeptides may be monomers or multimers; glycosylated or non-glycosylated; pegylated or non-pegylated; and may or may not include an initial methionine amino acid residue.

The invention also provides soluble UMLR receptors, used to form fusion or chimeric proteins with human Ig, as His-tagged proteins, or FLAG<sup>TM</sup>-tagged proteins. One such construct is comprises residues 1 to 129 of SEQ ID NO:2, fused to human Ig.

UMLR or UMLR-Ig chimeric proteins are used, for example, to identify the UMLR ligands, including the natural ligand, as well as agonists and antagonists of the natural ligand. Another construct comprises residues 1 to 173 of SEQ ID NO: 38 fused to the human Ig. Additional constructs comprise residues 1 to 130 of SEQ ID NO:2, residues 1 to 131 of SEQ ID NO:2, residues 1 to 132 of SEQ ID NO:2, residues 1 to 133 of SEQ ID NO:2, residues 1 to 134 of SEQ ID NO:2, residues 1 to 135 of SEQ ID NO:2, residues 1 to 136 of SEQ ID NO:2 fused to the human Ig. Additional constructs consist of residues 2 to 130 of SEQ ID NO:2, residues 2 to 131 of SEQ ID NO:2, residues 2 to 132 of SEQ ID NO:2, residues 2 to 133 of SEQ ID NO:2, residues 2 to 134 of SEQ ID NO:2, residues 2 to 135 of SEQ ID NO:2, residues 2 to 136 of SEQ ID NO:2, residues 1 to 173 of SEQ ID NO:38, residues 1 to 147 of SEQ ID NO:38, residues 1 to 154 of SEQ ID NO:38, and residues 1 to 165 of SEQ ID NO:38, fused to the human Ig. Using labeled soluble UMLR, cells expressing the ligand are identified by fluorescence immunocytometry or immunohistochemistry. The soluble fusion proteins or soluble Ig fusion protein is useful in studying the distribution of the ligand on tissues or specific cell lineages, and to provide insight into receptor/ligand biology.

In an alternative approach, a soluble UMLR receptor extracellular ligand-binding region can be expressed as a chimera with immunoglobulin heavy chain constant regions, typically an F<sub>C</sub> fragment, which contains two constant region domains and a hinge region, but lacks the variable region. Such fusions are typically secreted as multimeric molecules, wherein the Fc portions are disulfide bonded to each other and two receptor polypeptides are arrayed in close proximity to each other. Fusions of this

type can be used to affinity purify the cognate ligand from solution, as an *in vitro* assay tool, to block signals *in vitro* by specifically titrating out ligand, and as antagonists *in vivo* by administering them to block ligand stimulation. To purify ligand, a UMLR-Ig fusion protein (chimera) is added to a sample containing the ligand under conditions that

5 facilitate receptor-ligand binding (typically near-physiological temperature, pH, and ionic strength). The chimera-ligand complex is then separated by the mixture using protein A, which is immobilized on a solid support (e.g., insoluble resin beads). The ligand is then eluted using conventional chemical techniques, such as with a salt or pH gradient. In the alternative, the chimera itself can be bound to a solid support, with binding and elution

10 carried out as above. For use in assays, the chimeras are bound to a support via the F<sub>C</sub> region and used in an ELISA format.

Ligand binding studies have been performed on a number of known tumor necrosis factors including TNF $\alpha$ , TNF $\beta$ , THANK (Mukhopadhyay et al., J. Biol. Chem. 274:15978-81, 1999), FasL, RankL, Trail, and Tweak. None of these ligands has

15 been shown to bind to ztnfr11.

To direct the export of a UMLR polypeptide from the host cell, the UMLR DNA may be linked to a second DNA segment encoding a secretory peptide, such as a t-PA secretory peptide or a UMLR secretory peptide. To facilitate purification of the secreted polypeptide, a C-terminal extension, such as a poly-histidine tag,

20 substance P, Flag peptide (Hopp et al., Bio/Technology 6:1204-1210, 1988; available from Eastman Kodak Co., New Haven, CT), maltose binding protein, or another polypeptide or protein for which an antibody or other specific binding agent is available, can be fused to the UMLR polypeptide.

The present invention also includes "functional fragments" of UMLR

25 polypeptides and nucleic acid molecules encoding such functional fragments. Routine deletion analyses of nucleic acid molecules can be performed to obtain functional fragments of a nucleic acid molecule that encodes an UMLR polypeptide. As an illustration, DNA molecules having the nucleotide sequence of SEQ ID NOs:1, 22, 32, 33, and 36, can be digested with *Bal*31 nuclease to obtain a series of nested deletions.

30 The fragments are then inserted into expression vectors in proper reading frame, and the expressed polypeptides are isolated and tested for cell-cell interactions, or for the ability

to bind anti-UMLR antibodies. One alternative to exonuclease digestion is to use oligonucleotide-directed mutagenesis to introduce deletions or stop codons to specify production of a desired fragment. Alternatively, particular fragments of an UMLR gene can be synthesized using the polymerase chain reaction.

5                   Standard methods for identifying functional domains are well-known to those of skill in the art. For example, studies on the truncation at either or both termini of interferons have been summarized by Horisberger and Di Marco, Pharmac. Ther. 66:507 (1995). Moreover, standard techniques for functional analysis of proteins are described by, for example, Treuter *et al.*, Molec. Gen. Genet. 240:113 (1993), Content *et al.*,  
10 *“Expression and preliminary deletion analysis of the 42 kDa 2-5A synthetase induced by human interferon,”* in *Biological Interferon Systems, Proceedings of ISIR-TNO Meeting on Interferon Systems*, Cantell (ed.), pages 65-72 (Nijhoff 1987), Herschman, “The EGF Receptor,” in *Control of Animal Cell Proliferation, Vol. 1*, Boynton *et al.*, (eds.) pages 169-199 (Academic Press 1985), Coumailleau *et al.*, J. Biol. Chem.  
15 270:29270 (1995); Fukunaga *et al.*, J. Biol. Chem. 270:25291 (1995); Yamaguchi *et al.*, Biochem. Pharmacol. 50:1295 (1995), and Meisel *et al.*, Plant Molec. Biol. 30:1 (1996).

The present invention also contemplates functional fragments of an UMLR gene that has amino acid changes, compared with the amino acid sequences of SEQ ID NOs:2, 27, 29, 35, and 38. A variant UMLR gene can be identified on the basis  
20 of structure by determining the level of identity with nucleotide and amino acid sequences of SEQ ID NOs:1, 2, 3, 22, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, and 39 as discussed above. An alternative approach to identifying a variant gene on the basis of structure is to determine whether a nucleic acid molecule encoding a potential variant UMLR gene can hybridize to a nucleic acid molecule having the nucleotide sequence of  
25 SEQ ID NO:1, as discussed above.

Using the methods discussed herein, one of ordinary skill in the art can identify and/or prepare a variety of polypeptide fragments or variants of SEQ ID NO:2 or that retain the ligand-binding, or intracellular signaling activity of the wild-type UMLR protein. Such polypeptides may include additional amino acids from, for example,  
30 cysteine-rich pseudo-repeats, a linker domain, a transmembrane and cytoplasmic domains, including amino acids responsible for intracellular signaling; fusion domains;

affinity tags; and the like. Similarly, the cysteine -rich pseudo repeats (i.e., residues 2 to 42 of SEQ ID NO:2, residues 43 to 84 of SEQ ID NO:2, and/or residues 85 to 129 of SEQ ID NO:2) can be substituted for the cysteine-rich pseudo repeats from other TNFR family member to increase or decrease ligand binding, or specificity.

5                Within the polypeptides of the present invention are polypeptides that comprise an epitope-bearing portion of a protein as shown in SEQ ID NO:2. An "epitope" is a region of a protein to which an antibody can bind. See, for example, Geysen et al., *Proc. Natl. Acad. Sci. USA* 81:3998-4002, 1984. Epitopes can be linear or conformational, the latter being composed of discontinuous regions of the protein that  
10 form an epitope upon folding of the protein. Linear epitopes are generally at least 6 amino acid residues in length. Relatively short synthetic peptides that mimic part of a protein sequence are routinely capable of eliciting an antiserum that reacts with the partially mimicked protein. See, Sutcliffe et al., *Science* 219:660-666, 1983. Antibodies that recognize short, linear epitopes are particularly useful in analytic and diagnostic  
15 applications that employ denatured protein, such as Western blotting (Tobin, *Proc. Natl. Acad. Sci. USA* 76:4350-4356, 1979), or in the analysis of fixed cells or tissue samples. Antibodies to linear epitopes are also useful for detecting fragments of UMLR, such as might occur in body fluids or cell culture media.

                  Antigenic, epitope-bearing polypeptides of the present invention are  
20 useful for raising antibodies, including monoclonal antibodies, that specifically bind to a UMLR protein. Antigenic, epitope-bearing polypeptides contain a sequence of at least six, preferably at least nine, more preferably from 15 to about 30 contiguous amino acid residues of a UMLR protein (e.g., SEQ ID NO:2). Polypeptides comprising a larger portion of a UMLR protein, i.e. from 30 to 50 residues up to the entire sequence, are  
25 included. It is preferred that the amino acid sequence of the epitope-bearing polypeptide is selected to provide substantial solubility in aqueous solvents, that is the sequence includes relatively hydrophilic residues, and hydrophobic residues are substantially avoided. Preferred such regions include the cysteine-rich pseudo-repeats, the linker domain, the transmembrane domain, or the cytoplasmic domain UMLR and fragments  
30 thereof. Specific, preferred polypeptides in this regard include those comprising residues 2 to 42 of SEQ ID NO:2; residues 43 to 84 of SEQ ID NO:2; residues 85 to 129 of SEQ

ID NO:2; residues 130 to 136 of SEQ ID NO:2; residues 137 to 161 of SEQ ID NO:2; residues 162 to 269 of SEQ ID NO:2; residues 1 to 269 of SEQ ID NO:2; residues 1 to 129 of SEQ ID NO:2; residues 1 to 136 of SEQ ID NO:2; residues 137 to 269 of SEQ ID NO:2; residues 85 to 173 of SEQ ID NO:38; residues 85 to 163 of SEQ ID NO:38; residues 85 to 165 of SEQ ID NO:38; residues 85 to 147 of SEQ ID NO:38; and residues 85 to 154 of SEQ ID NO:38.

The present invention also provides polypeptide fragments or peptides comprising an epitope-bearing portion of an UMLR polypeptide described herein. Such fragments or peptides may comprise an "immunogenic epitope," which is a part of a protein that elicits an antibody response when the entire protein is used as an immunogen. Immunogenic epitope-bearing peptides can be identified using standard methods (see, for example, Geysen *et al.*, Proc. Nat'l Acad. Sci. USA 81:3998 (1983)).

In contrast, polypeptide fragments or peptides may comprise an "antigenic epitope," which is a region of a protein molecule to which an antibody can specifically bind. Certain epitopes consist of a linear or contiguous stretch of amino acids, and the antigenicity of such an epitope is not disrupted by denaturing agents. It is known in the art that relatively short synthetic peptides that can mimic epitopes of a protein can be used to stimulate the production of antibodies against the protein (see, for example, Sutcliffe *et al.*, Science 219:660 (1983)). Accordingly, antigenic epitope-bearing peptides and polypeptides of the present invention are useful to raise antibodies that bind with the polypeptides described herein.

Antigenic epitope-bearing peptides and polypeptides contain at least four to ten amino acids, preferably at least ten to fifteen amino acids, more preferably 15 to 30 amino acids of SEQ ID NOs:2, 27, 29, 34, 35, 37, and 38. Such epitope-bearing peptides and polypeptides can be produced by fragmenting an UMLR polypeptide, or by chemical peptide synthesis, as described herein. Moreover, epitopes can be selected by phage display of random peptide libraries (see, for example, Lane and Stephen, Curr. Opin. Immunol. 5:268 (1993), and Cortese *et al.*, Curr. Opin. Biotechnol. 7:616 (1996)). Standard methods for identifying epitopes and producing antibodies from small peptides that comprise an epitope are described, for example, by Mole, "Epitope Mapping," in *Methods in Molecular Biology*, Vol. 10, Manson (ed.), pages 105-116 (The Humana

Press, Inc. 1992), Price, "Production and Characterization of Synthetic Peptide-Derived Antibodies," in *Monoclonal Antibodies: Production, Engineering, and Clinical Application*, Ritter and Ladyman (eds.), pages 60-84 (Cambridge University Press 1995), and Coligan *et al.* (eds.), *Current Protocols in Immunology*, pages 9.3.1 - 9.3.5 and  
 5 pages 9.4.1 - 9.4.11 (John Wiley & Sons 1997).

UMLR polypeptides can also be used to prepare antibodies that specifically bind to UMLR epitopes, peptides or polypeptides. The UMLR polypeptide or a fragment thereof serves as an antigen (immunogen) to inoculate an animal and elicit an immune response. One of skill in the art would recognize that antigenic, epitope-  
 10 bearing polypeptides contain a sequence of at least 6, preferably at least 9, and more preferably at least 15 to about 30 contiguous amino acid residues of a UMLR polypeptide (e.g., SEQ ID NOs:2, 27, 29, 34, 35, 37, and 38). Polypeptides comprising a larger portion of a UMLR polypeptide, i.e., from 30 to 10 residues up to the entire length of the amino acid sequence are included. Antigens or immunogenic epitopes can also  
 15 include attached tags, adjuvants and carriers, as described herein. Suitable antigens include the UMLR polypeptides encoded by SEQ ID NO:2 from amino acid number 1 to amino acid number 269; polypeptides encoded by SEQ ID NO:27 from amino acid number 1 to amino acid number 297; polypeptides encoded by SEQ ID NO:35 from amino acid number 1 to amino acid number 299; polypeptides encoded by SEQ ID  
 20 NO:38 from amino acid number 1 to amino acid number 173, or a contiguous 9 to 269 amino acid fragment thereof.

As an illustration, potential antigenic sites in UMLR were identified using the Jameson-Wolf method, Jameson and Wolf, *CABIOS* 4:181, (1988), as implemented by the PROTEAN program (version 3.14) of LASERGENE (DNASTAR; Madison, WI).  
 25 Default parameters were used in this analysis.

Suitable antigens from this analysis include residue 4 to residue 13 of SEQ ID NO:2; residue 17 to residue 39 of SEQ ID NO:2; residue 44 to residue 60 of SEQ ID NO:2; residue 69 to residue 104 of SEQ ID NO:2; residue 106 to residue 115 of SEQ ID NO:2; residue 128 to residue 136 of SEQ ID NO:2; residue 165 to residue 174  
 30 of SEQ ID NO:2 ; residue 179 to residue 204 of SEQ ID NO:2; and residue 245 to residue 260 of SEQ ID NO:2. Suitable antigens from this analysis of SEQ ID NOs:35

and 38 include residue 181 to residue 210 of SEQ ID NO:35; residue 219 to residue 226 of SEQ ID NO:35; residue 218 to residue 236 of SEQ ID NO:35; and residue 162 to residue 169 of SEQ ID NO:38; and residue 149 to residue 173 of SEQ ID NO:38. Hydrophilic peptides, such as those predicted by one of skill in the art from a hydrophobicity plot are also immunogenic. UMLR hydrophilic peptides include peptides comprising amino acid sequences selected from the group consisting of: residues 1 to 13 of SEQ ID NO:2; residues 21 to 38 of SEQ ID NO:2; residues 44 to 59 of SEQ ID NO:2; residues 88 to 99 of SEQ ID NO:2; residues 165 to 173 of SEQ ID NO:2; residues 194 to 205 of SEQ ID NO:2; residues 213 to 221 of SEQ ID NO:2; residue 194 to residue 236 of SEQ ID NO:35; and residue 150 to residue 171 of SEQ ID NO:38. Regions of the polypeptide, which are likely to be on the surface of the folded protein, are also likely to be antigenic. These regions include: residue 230 to residue 235 of SEQ ID NO:35; and residue 151 to residue 158 of SEQ ID NO:38. Antibodies from an immune response generated by inoculation of an animal with these antigens can be isolated and purified as described herein. Methods for preparing and isolating polyclonal and monoclonal antibodies are well known in the art. See, for example, Current Protocols in Immunology, Cooligan, et al. (eds.), National Institutes of Health, John Wiley and Sons, Inc., 1995; Sambrook et al., Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, NY, 1989; and Hurrell, J. G. R., Ed., Monoclonal Hybridoma Antibodies: Techniques and Applications, CRC Press, Inc., Boca Raton, FL, 1982.

As would be evident to one of ordinary skill in the art, polyclonal antibodies can be generated from inoculating a variety of warm-blooded animals such as horses, cows, goats, sheep, dogs, chickens, rabbits, mice, and rats with a UMLR polypeptide or a fragment thereof. The immunogenicity of a UMLR polypeptide may be increased through the use of an adjuvant, such as alum (aluminum hydroxide) or Freund's complete or incomplete adjuvant. Polypeptides useful for immunization also include fusion polypeptides, such as fusions of UMLR or a portion thereof with an immunoglobulin polypeptide or with maltose binding protein. The polypeptide immunogen may be a full-length molecule or a portion thereof. If the polypeptide portion is "hapten-like", such portion may be advantageously joined or linked to a

macromolecular carrier (such as keyhole limpet hemocyanin (KLH), bovine serum albumin (BSA) or tetanus toxoid) for immunization.

As used herein, the term "antibodies" includes polyclonal antibodies, affinity-purified polyclonal antibodies, monoclonal antibodies, and antigen-binding fragments, such as F(ab')<sub>2</sub> and Fab proteolytic fragments. Genetically engineered intact antibodies or fragments, such as chimeric antibodies, Fv fragments, single chain antibodies and the like, as well as synthetic antigen-binding peptides and polypeptides, are also included. Non-human antibodies may be humanized by grafting non-human CDRs onto human framework and constant regions, or by incorporating the entire non-human variable domains (optionally "cloaking" them with a human-like surface by replacement of exposed residues, wherein the result is a "veneered" antibody). In some instances, humanized antibodies may retain non-human residues within the human variable region framework domains to enhance proper binding characteristics. Through humanizing antibodies, biological half-life may be increased, and the potential for adverse immune reactions upon administration to humans is reduced.

Alternative techniques for generating or selecting antibodies useful herein include *in vitro* exposure of lymphocytes to UMLR protein or peptide, and selection of antibody display libraries in phage or similar vectors (for instance, through use of immobilized or labeled UMLR protein or peptide). Genes encoding polypeptides having potential UMLR polypeptide binding domains can be obtained by screening random peptide libraries displayed on phage (phage display) or on bacteria, such as *E. coli*. Nucleotide sequences encoding the polypeptides can be obtained in a number of ways, such as through random mutagenesis and random polynucleotide synthesis. These random peptide display libraries can be used to screen for peptides which interact with a known target which can be a protein or polypeptide, such as a ligand or receptor, a biological or synthetic macromolecule, or organic or inorganic substances. Techniques for creating and screening such random peptide display libraries are known in the art (Ladner et al., US Patent NO. 5,223,409; Ladner et al., US Patent NO. 4,946,778; Ladner et al., US Patent NO. 5,403,484 and Ladner et al., US Patent NO. 5,571,698) and random peptide display libraries and kits for screening such libraries are available commercially, for instance from CLONTECH Laboratories, Inc., (Palo Alto, CA), Invitrogen Inc. (San



Diego, CA), New England Biolabs, Inc. (Beverly, MA) and Pharmacia LKB Biotechnology Inc. (Piscataway, NJ). Random peptide display libraries can be screened using the UMLR sequences disclosed herein to identify proteins which bind to UMLR. These “binding proteins” which interact with UMLR polypeptides can be used for

5 tagging cells; for isolating homolog polypeptides by affinity purification; they can be directly or indirectly conjugated to drugs, toxins, radionuclides and the like. These binding proteins can also be used in analytical methods such as for screening expression libraries and neutralizing activity. The binding proteins can also be used for diagnostic

10 assays for determining circulating levels of polypeptides; for detecting or quantitating soluble polypeptides as marker of underlying pathology or disease. These binding proteins can also act as UMLR “antagonists” to block UMLR binding and signal transduction *in vitro* and *in vivo*. These anti-UMLR binding proteins would be useful for modulating, for example, apoptosis, myogenesis, immunologic recognition, tumor formation, and cell-cell interactions in general.

15 As used herein, the term “binding proteins” additionally includes antibodies to UMLR polypeptides, the cognate ligand of UMLR polypeptides, proteins useful for purification of UMLR polypeptides, and proteins associated with the cytoplasmic domain (residues 137 to 269 of SEQ ID NO:2; residues 137 to 297 of SEQ ID NO:27; residues 137 to 299 of SEQ ID NO:35, and residues 137 to 173 of SEQ ID

20 NO:38). Such cytoplasmic domain associated peptides, also called cytoplasmic mediators, function in intracellular signaling of UMLR polypeptides. See Bazzoni, F. et al., J. of Inflammation 45:221-238, 1995. These cytoplasmic mediators include, but are not limited to TRAP-1, TRADD, RIP, TRAF-1, TRAF-2, LAP-1, FADD/MORT-1, and CAP-1.

25 Antibodies are determined to be specifically binding if they exhibit a threshold level of binding activity (to a UMLR polypeptide, peptide or epitope) of at least 10-fold greater than the binding affinity to a control (non-UMLR ) polypeptide. The binding affinity of an antibody can be readily determined by one of ordinary skill in the art, for example, by Scatchard analysis (Scatchard, G., Ann. NY Acad. Sci. 51: 660-

30 672, 1949).

A variety of assays known to those skilled in the art can be utilized to detect antibodies which specifically bind to UMLR proteins or peptides. Exemplary assays are described in detail in Antibodies: A Laboratory Manual, Harlow and Lane (Eds.), Cold Spring Harbor Laboratory Press, 1988. Representative examples of such

5 assays include: concurrent immunoelectrophoresis, radioimmunoassay, radioimmuno-precipitation, enzyme-linked immunosorbent assay (ELISA), dot blot or Western blot assay, inhibition or competition assay, and sandwich assay. In addition, antibodies can be screened for binding to wild-type versus mutant UMLR protein or polypeptide.

Antibodies to UMLR may be used for immunohistochemical tagging cells

10 that express UMLR; for isolating UMLR by affinity purification; for diagnostic assays for determining circulating levels of UMLR polypeptides; for detecting or quantitating soluble UMLR as marker of underlying pathology or disease; in analytical methods employing FACS; for screening expression libraries; for generating anti-idiotypic antibodies; and as neutralizing antibodies or as antagonists to block UMLR *in vitro* and

15 *in vivo*. Suitable direct tags or labels include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent markers, chemiluminescent markers, magnetic particles and the like; indirect tags or labels may feature use of biotin-avidin or other complement/anti-complement pairs as intermediates. Antibodies herein may also be directly or indirectly conjugated to drugs, toxins, radionuclides and the like, and these

20 conjugates used for *in vivo* diagnostic or therapeutic applications. Moreover, antibodies to UMLR or fragments thereof may be used *in vitro* to detect denatured UMLR or fragments thereof in assays, for example, Western Blots or other assays known in the art.

The soluble UMLR is useful in studying the distribution of ligands on tissues or specific cell lineages, and to provide insight into receptor/ligand biology.

25 Using labeled UMLR, cells expressing the ligand are identified by fluorescence immunocytometry or immunocytochemistry. Application may also be made of the specificity of TNF receptors for their ligands.

Antibodies can be made to soluble UMLR polypeptides which are His or FLAG<sup>TM</sup> tagged. Alternatively, such polypeptides form a fusion protein with Human Ig.

30 In particular, antiserum containing polypeptide antibodies to His-tagged, or FLAG<sup>TM</sup>-tagged soluble UMLR can be used in analysis of tissue distribution of UMLR by

immunohistochemistry on human or primate tissue. These soluble UMLR polypeptides can also be used to immunize mice in order to produce monoclonal antibodies to a soluble human UMLR polypeptide. Monoclonal antibodies to a soluble human UMLR polypeptide can also be used to mimic ligand/receptor coupling, resulting in activation or inactivation of the ligand/receptor pair. For instance, it has been demonstrated that cross-linking anti-soluble CD40 monoclonal antibodies provides a stimulatory signal to B cells that have been sub-optimally activated with anti-IgM or LPS, and results in proliferation and immunoglobulin production. These same monoclonal antibodies act as antagonists when used in solution by blocking activation of the receptor. Monoclonal antibodies to UMLR can be used to determine the distribution, regulation and biological interaction of the UMLR /UMLR -ligand pair on specific cell lineages identified by tissue distribution studies.

Soluble receptors or antibodies to the receptor can also be directly or indirectly conjugated to drugs, toxins, radionuclides and the like, and these conjugates used for *in vivo* diagnostic or therapeutic applications. For instance, polypeptides or antibodies of the present invention can be used to identify or treat tissues or organs that express a corresponding anti-complementary molecule (ligand or antigen, respectively, for instance). More specifically, UMLR polypeptides or anti-UMLR antibodies, or bioactive fragments or portions thereof, can be coupled to detectable or cytotoxic molecules and delivered to a mammal having cells, tissues or organs that express the anti-complementary molecule.

Suitable detectable molecules may be directly or indirectly attached to the polypeptide or antibody, and include radionuclides, enzymes, substrates, cofactors, inhibitors, fluorescent markers, chemiluminescent markers, magnetic particles and the like. Suitable cytotoxic molecules may be directly or indirectly attached to the polypeptide or antibody, and include bacterial or plant toxins (for instance, diphtheria toxin, *Pseudomonas* exotoxin, ricin, abrin and the like), as well as therapeutic radionuclides, such as iodine-131, rhenium-188 or yttrium-90 (either directly attached to the polypeptide or antibody, or indirectly attached through means of a chelating moiety, for instance). Polypeptides or antibodies may also be conjugated to cytotoxic drugs, such as adriamycin. For indirect attachment of a detectable or cytotoxic molecule, the

detectable or cytotoxic molecule can be conjugated with a member of a complementary/anticomplementary pair, where the other member is bound to the polypeptide or antibody portion. For these purposes, biotin/streptavidin is an exemplary complementary/anticomplementary pair.

5 In another embodiment, polypeptide-toxin fusion proteins or antibody-toxin fusion proteins can be used for targeted cell or tissue inhibition or ablation (for instance, to treat cancer or inflammatory cells or tissues). Alternatively, a fusion protein including only the cysteine-rich pseudo-repeats may be suitable for directing a detectable molecule, a cytotoxic molecule or a complementary molecule to a cell or tissue type of  
10 interest. Similarly, the corresponding ligand to UMLR can be conjugated to a detectable or cytotoxic molecule and provide a generic targeting vehicle for cell/tissue-specific delivery of generic anti-complementary-detectable/ cytotoxic molecule conjugates.

As would be evident to one of skill in the art, certain molecular markers can aid in identifying the existence and prognosis of disease. Such markers include, for  
15 example, p53, C-Ki-ras, and c-erbB-2. See, in general, Schneider, P. et al. Br. J. Cancer 83(4): 473-479, 2000; and Watatani, M., et al., Surg. Today 30(6):516-522, 2000. One of routine skill in the art will also know that such molecular markers can also be used as a target for treatment with or without combination therapy with other agents, including chemotherapy. An example of a treatment approach to a membrane bound receptor is  
20 the use of trastuzumab to treat breast cancer wherein the cancerous tissue is overexpressing HER2. See, for example, Baselga, J., et al., Semin. Oncol. 26 (4 Suppl 12): 78-83, 1999; Ravdin, P., Semin. Oncol. 26 (4 Suppl 12): 117-23, 1999; Shak, S., Semin. Oncol. 26 (4 Suppl 12): 71-7, 1999; and Stebbing, J., et al., Cancer Treat. Rev. 26(4): 287-290, 2000.

25 As a protein that shows upregulation in some tumor cells, such as, for example, melanoma, lung carcinoma, breast carcinoma, osteosarcoma, and lymphoma, polynucleotides and polypeptides of the present invention, fragments thereof, and binding proteins thereto (including antibodies, and ligands) can be used in a multitude of ways to detect and/or diagnose such diseases. For example, polynucleotide probes,  
30 (including DNA, RNA, and peptide-nucleic acid) can be used as a diagnostic marker to determine if such disease tissues are present. Hybridization techniques are taught

elsewhere in this application and are widely known by one of skill in the art. Such polynucleotides, fragments thereof, and fusions thereto, can also be used to incorporate the proper polynucleotide sequence into a tissue, cell line or organism defective in the proper gene, as is also taught elsewhere in this application. As a means of treating such disease, said polynucleotides, fragments thereof, and fusions thereto, can also be administered in the presence of an agent that allows the DNA to traverse the cell membrane and act as a tag for cell ablation to a therapeutic agent with an appropriate binding partner. In this manner cells which contain ztnfr11 polynucleotides can be inhibited or destroyed.

10           Ztnfr11 polypeptides, fragments thereof, and fusions thereto, can be used both diagnostically and therapeutically. Such ztnfr11 polypeptides, including soluble polypeptides, can be used as a marker for identifying tumor cells, such as, for example, melanoma, lung carcinoma, breast carcinoma, osteosarcoma, and lymphoma. As a soluble polypeptide, this diagnosis can be determined by measuring ztnfr11 polypeptides, fragments thereof, and fusions thereto in body fluids, including, but not limited to, blood, plasma, saliva, urine, lavage fluid and biopsy fluid. As a membrane bound polypeptide ztnfr11 polypeptides, fragments thereof, and fusions thereto, can be measured in tissue biopsies (i.e., excised from the body) as well as locally (i.e., epithelial surfaces) using imaging and/ or visualization. The targeting of such disease tissues will be helpful in treatment options. For example, if the spread of disease is limited, such visualization will aid a surgeon in resection of disease tissue. Similarly, the presence of membrane bound ztnfr11 can be used as a target for a ztnfr11 binding protein (including its ligand or antibodies) that has been fused or conjugated to an inhibitory or ablative agent.

25           In another embodiment, UMLR -cytokine fusion proteins or antibody-cytokine fusion proteins can be used for enhancing *in vivo* killing of target tissues (for example, uterus, as well as in melanoma, osteosarcoma, breast carcinoma, and lymphoma, and in tumors of the lung), if the UMLR polypeptide or anti-UMLR antibody targets hyperproliferative tissues from these organs. (See, generally, Hornick et al., Blood 89:4437-47, 1997). They described fusion proteins that enable targeting of a cytokine to a desired site of action, thereby providing an elevated local concentration of cytokine. Suitable UMLR polypeptides or anti-UMLR antibodies target an undesirable

cell or tissue (i.e., a tumor or a leukemia), and the fused cytokine mediates improved target cell lysis by effector cells. Suitable cytokines for this purpose include interleukin 2 and granulocyte-macrophage colony-stimulating factor (GM-CSF), for instance.

UMLR polynucleotides and/or polypeptides may be useful for regulating the maturation of TNF ligand-bearing cells, such as T cells, B cells, lymphocytes, peripheral blood mononuclear cells, polymorphonuclear leukocytes, fibroblasts and hematopoietic cells. UMLR polypeptides will also find use in mediating metabolic or physiological processes *in vivo*. The effects of a compound on proliferation and differentiation can be measured *in vitro* using cultured cells. Bioassays and ELISAs are available to measure cellular response to UMLR, in particular are those which measure changes in cytokine production as a measure of cellular response (see for example, Current Protocols in Immunology ed. John E. Coligan et al., NIH, 1996). Assays to measure other cellular responses, including antibody isotype, monocyte activation, NK cell formation, antigen presenting cell function, apoptosis are known in the art.

The UMLR polypeptides of the present invention, including full-length polypeptides, biologically active fragments, and fusion polypeptides, can be produced in genetically engineered host cells according to conventional techniques. Suitable host cells are those cell types that can be transformed or transfected with exogenous DNA and grown in culture, and include bacteria, fungal cells, and cultured higher eukaryotic cells. Eukaryotic cells, particularly cultured cells of multicellular organisms, are preferred. Techniques for manipulating cloned DNA molecules and introducing exogenous DNA into a variety of host cells are disclosed by Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989, and Ausubel et al., eds., Current Protocols in Molecular Biology, John Wiley and Sons, Inc., NY, 1987.

In general, a DNA sequence encoding a UMLR polypeptide is operably linked to other genetic elements required for its expression, generally including a transcription promoter and terminator, within an expression vector. The vector will also commonly contain one or more selectable markers and one or more origins of replication, although those skilled in the art will recognize that within certain systems selectable markers may be provided on separate vectors, and replication of the

exogenous DNA may be provided by integration into the host cell genome. Selection of promoters, terminators, selectable markers, vectors and other elements is a matter of routine design within the level of ordinary skill in the art. Many such elements are described in the literature and are available through commercial suppliers.

5 To direct a UMLR polypeptide into the secretory pathway of a host cell, a secretory signal sequence (also known as a leader sequence, prepro sequence or pre sequence) can be provided in the expression vector. The secretory signal sequence may be derived from another secreted protein (e.g., APO4, or t-PA) or synthesized *de novo*. The secretory signal sequence is operably linked to the UMLR DNA sequence, i.e., the  
 10 two sequences are joined in the correct reading frame and positioned to direct the newly synthesized polypeptide into the secretory pathway of the host cell. Secretory signal sequences are commonly positioned 5' to the DNA sequence encoding the polypeptide of interest, although certain secretory signal sequences may be positioned elsewhere in the DNA sequence of interest (see, e.g., Welch et al., U.S. Patent No. 5,037,743; Holland et  
 15 al., U.S. Patent No. 5,143,830).

The cytoplasmic domain of UMLR can be substituted by a heterologous sequence providing a different cytoplasmic domain. In this case, the fusion product can be secreted, and the cysteine-rich pseudo-repeat domain of UMLR can direct the new  
 20 cytoplasmic domain to a specific tissue described above. This substituted cytoplasmic domain can be chosen from the cytoplasmic domain represented by the TNFR protein families. Similarly, the cysteine-rich pseudo-repeat domain of UMLR protein can be substituted by a heterologous sequence providing a different cysteine-rich pseudo-repeat domain. Again, the fusion product can be secreted and the substituted cysteine-rich  
 25 pseudo-repeat domain can target the cytoplasmic domain of UMLR to a specific tissue. The substituted cysteine-rich pseudo-repeat domain can be chosen from the cysteine-rich pseudo-repeat domains of the TNFR protein families. In these cases, the fusion products can be soluble or membrane-bound proteins.

Cultured mammalian cells are suitable hosts within the present invention. Methods for introducing exogenous DNA into mammalian host cells include calcium  
 30 phosphate-mediated transfection (Wigler et al., Cell 14:725, 1978; Corsaro and Pearson, Somatic Cell Genetics 7:603, 1981; Graham and Van der Eb, Virology 52:456, 1973),

electroporation (Neumann et al., EMBO J. 1:841-5, 1982), DEAE-dextran mediated transfection (Ausubel et al., ibid.), and liposome-mediated transfection (Hawley-Nelson et al., Focus 15:73, 1993; Ciccarone et al., Focus 15:80, 1993, and viral vectors (Miller and Rosman, BioTechniques 7:980-90, 1989; Wang and Finer, Nature Med. 2:714-6, 5 1996). The production of recombinant polypeptides in cultured mammalian cells is disclosed, for example, by Levinson et al., U.S. Patent No. 4,713,339; Hagen et al., U.S. Patent No. 4,784,950; Palmiter et al., U.S. Patent No. 4,579,821; and Ringold, U.S. Patent No. 4,656,134. Suitable cultured mammalian cells include the COS-1 (ATCC No. CRL 1650), COS-7 (ATCC No. CRL 1651), BHK (ATCC No. CRL 1632), BHK 10 570 (ATCC No. CRL 10314), 293 (ATCC No. CRL 1573; Graham et al., J. Gen. Virol. 36:59-72, 1977) and Chinese hamster ovary (e.g. CHO-K1; ATCC No. CCL 61) cell lines. Additional suitable cell lines are known in the art and available from public depositories such as the American Type Culture Collection, Rockville, Maryland. In general, strong transcription promoters are preferred, such as promoters from SV-40 or 15 cytomegalovirus. See, e.g., U.S. Patent No. 4,956,288. Other suitable promoters include those from metallothionein genes (U.S. Patent Nos. 4,579,821 and 4,601,978) and the adenovirus major late promoter.

Drug selection is generally used to select for cultured mammalian cells into which foreign DNA has been inserted. Such cells are commonly referred to as 20 "transfectants". Cells that have been cultured in the presence of the selective agent and are able to pass the gene of interest to their progeny are referred to as "stable transfectants." A preferred selectable marker is a gene encoding resistance to the antibiotic neomycin. Selection is carried out in the presence of a neomycin-type drug, such as G-418 or the like. Selection systems can also be used to increase the expression 25 level of the gene of interest, a process referred to as "amplification." Amplification is carried out by culturing transfectants in the presence of a low level of the selective agent and then increasing the amount of selective agent to select for cells that produce high levels of the products of the introduced genes. A preferred amplifiable selectable marker is dihydrofolate reductase, which confers resistance to methotrexate. Other drug 30 resistance genes (e.g., hygromycin resistance, multi-drug resistance, puromycin acetyltransferase) can also be used. Alternative markers that introduce an altered



phenotype, such as green fluorescent protein, or cell surface proteins, such as CD4, CD8, Class I MHC, or placental alkaline phosphatase, may be used to sort transfected cells from untransfected cells by such means as FACS sorting or magnetic bead separation technology.

- 5 Other higher eukaryotic cells can also be used as hosts, including plant cells, insect cells and avian cells. The use of *Agrobacterium rhizogenes* as a vector for expressing genes in plant cells has been reviewed by Sinkar et al., J. Biosci. (Bangalore) 11:47-58, 1987. Transformation of insect cells and production of foreign polypeptides therein is disclosed by Guarino et al., U.S. Patent No. 5,162,222 and WIPO publication
- 10 WO 94/06463. Insect cells can be infected with recombinant baculovirus, commonly derived from *Autographa californica nuclear polyhedrosis virus* (AcNPV). See, King, L.A. and Possee, R.D., The Baculovirus Expression System: A Laboratory Guide, London, Chapman & Hall; O'Reilly, D.R. et al., Baculovirus Expression Vectors: A Laboratory Manual, New York, Oxford University Press., 1994; and, Richardson, C. D.,
- 15 Ed., Baculovirus Expression Protocols. Methods in Molecular Biology, Totowa, NJ, Humana Press, 1995. A second method of making recombinant UMLR baculovirus utilizes a transposon-based system described by Luckow (Luckow, V.A, et al., J Virol 67:4566-79, 1993). This system, which utilizes transfer vectors, is sold in the Bac-to-Bac™ kit (Life Technologies, Rockville, MD). This system utilizes a transfer vector,
- 20 pFastBac1™ (Life Technologies) containing a Tn7 transposon to move the DNA encoding the UMLR polypeptide into a baculovirus genome maintained in *E. coli* as a large plasmid called a "bacmid." The pFastBac1™ transfer vector utilizes the AcNPV polyhedrin promoter to drive the expression of the gene of interest, in this case UMLR . However, pFastBac1™ can be modified to a considerable degree. The polyhedrin
- 25 promoter can be removed and substituted with the baculovirus basic protein promoter (also known as *Pcor*, p6.9 or MP promoter) which is expressed earlier in the baculovirus infection, and has been shown to be advantageous for expressing secreted proteins. See, Hill-Perkins, M.S. and Possee, R.D., J. Gen. Virol. 71:971-6, 1990; Bonning, B.C. et al., J. Gen. Virol. 75:1551-6, 1994; and, Chazenbalk, G.D., and Rapoport, B., J. Biol Chem
- 30 270:1543-9, 1995. In such transfer vector constructs, a short or long version of the basic protein promoter can be used. Moreover, transfer vectors can be constructed which

replace the native UMLR secretory signal sequences with secretory signal sequences derived from insect proteins. For example, a secretory signal sequence from Ecdysteroid Glucosyltransferase (EGT), honey bee Melittin (Invitrogen, Carlsbad, CA), or baculovirus gp67 (PharMingen, San Diego, CA) can be used in constructs to replace the native UMLR secretory signal sequence. In addition, transfer vectors can include an in-frame fusion with DNA encoding an epitope tag at the C- or N-terminus of the expressed UMLR polypeptide, for example, a Glu-Glu epitope tag (Grussenmeyer, T. et al., Proc. Natl. Acad. Sci. 82:7952-4, 1985). Using a technique known in the art, a transfer vector containing UMLR is transformed into *E. coli*, and screened for bacmids which contain an interrupted lacZ gene indicative of recombinant baculovirus. The bacmid DNA containing the recombinant baculovirus genome is isolated, using common techniques, and used to transfect *Spodoptera frugiperda* cells, e.g. Sf9 cells. Recombinant virus that expresses UMLR is subsequently produced. Recombinant viral stocks are made by methods commonly used the art.

The recombinant virus is used to infect host cells, typically a cell line derived from the fall armyworm, *Spodoptera frugiperda*. See, in general, Glick and Pasternak, Molecular Biotechnology: Principles and Applications of Recombinant DNA, ASM Press, Washington, D.C., 1994. Another suitable cell line is the High FiveO™ cell line (Invitrogen) derived from *Trichoplusia ni* (U.S. Patent #5,300,435). Commercially available serum-free media are used to grow and maintain the cells. Suitable media are Sf900 II™ (Life Technologies) or ESF 921™ (Expression Systems) for the Sf9 cells; and Ex-cello405™ (JRH Biosciences, Lenexa, KS) or Express FiveO™ (Life Technologies) for the *T. ni* cells. The cells are grown up from an inoculation density of approximately  $2-5 \times 10^5$  cells to a density of  $1-2 \times 10^6$  cells at which time a recombinant viral stock is added at a multiplicity of infection (MOI) of 0.1 to 10, more typically near 3. Procedures used are generally described in available laboratory manuals (King, L. A. and Possee, R.D., ibid.; O'Reilly, D.R. et al., ibid.; Richardson, C. D., ibid.). Subsequent purification of the UMLR polypeptide from the supernatant can be achieved using methods described herein.

Fungal cells, including yeast cells, can also be used within the present invention. Yeast species of particular interest in this regard include *Saccharomyces*

*cerevisiae*, *Pichia pastoris*, and *Pichia methanolica*. Methods for transforming *S. cerevisiae* cells with exogenous DNA and producing recombinant polypeptides therefrom are disclosed by, for example, Kawasaki, U.S. Patent No. 4,599,311; Kawasaki et al., U.S. Patent No. 4,931,373; Brake, U.S. Patent No. 4,870,008; Welch et al., U.S. Patent No. 5,037,743; and Murray et al., U.S. Patent No. 4,845,075. Transformed cells are selected by phenotype determined by the selectable marker, commonly drug resistance or the ability to grow in the absence of a particular nutrient (e.g., leucine). A preferred vector system for use in *Saccharomyces cerevisiae* is the *POT1* vector system disclosed by Kawasaki et al. (U.S. Patent No. 4,931,373), which allows transformed cells to be selected by growth in glucose-containing media. Suitable promoters and terminators for use in yeast include those from glycolytic enzyme genes (see, e.g., Kawasaki, U.S. Patent No. 4,599,311; Kingsman et al., U.S. Patent No. 4,615,974; and Bitter, U.S. Patent No. 4,977,092) and alcohol dehydrogenase genes. See also U.S. Patents Nos. 4,990,446; 5,063,154; 5,139,936 and 4,661,454. Transformation systems for other yeasts, including *Hansenula polymorpha*, *Schizosaccharomyces pombe*, *Kluyveromyces lactis*, *Kluyveromyces fragilis*, *Ustilago maydis*, *Pichia pastoris*, *Pichia methanolica*, *Pichia guilliermondii* and *Candida maltosa* are known in the art. See, for example, Gleeson et al., *J. Gen. Microbiol.* 132:3459-65, 1986 and Cregg, U.S. Patent No. 4,882,279. *Aspergillus* cells may be utilized according to the methods of McKnight et al., U.S. Patent No. 4,935,349. Methods for transforming *Acremonium chrysogenum* are disclosed by Sumino et al., U.S. Patent No. 5,162,228. Methods for transforming *Neurospora* are disclosed by Lambowitz, U.S. Patent No. 4,486,533. The use of *Pichia methanolica* as host for the production of recombinant proteins is disclosed in U.S. patents 5,716,808, 5,736,383, 5,854,039, and 5,888,768.

Prokaryotic host cells, including strains of the bacteria *Escherichia coli*, *Bacillus* and other genera are also useful host cells within the present invention. Techniques for transforming these hosts and expressing foreign DNA sequences cloned therein are well known in the art (see, e.g., Sambrook et al., *ibid.*). When expressing a UMLR polypeptide in bacteria such as *E. coli*, the polypeptide may be retained in the cytoplasm, typically as insoluble granules, or may be directed to the periplasmic space by a bacterial secretion sequence. In the former case, the cells are lysed, and the granules

are recovered and denatured using, for example, guanidine isothiocyanate or urea. The denatured polypeptide can then be refolded and dimerized by diluting the denaturant, such as by dialysis against a solution of urea and a combination of reduced and oxidized glutathione, followed by dialysis against a buffered saline solution. In the latter case, the polypeptide can be recovered from the periplasmic space in a soluble and functional form by disrupting the cells (by, for example, sonication or osmotic shock) to release the contents of the periplasmic space and recovering the protein, thereby obviating the need for denaturation and refolding.

Transformed or transfected host cells are cultured according to conventional procedures in a culture medium containing nutrients and other components required for the growth of the chosen host cells. A variety of suitable media, including defined media and complex media, are known in the art and generally include a carbon source, a nitrogen source, essential amino acids, vitamins and minerals. Media may also contain such components as growth factors or serum, as required. The growth medium will generally select for cells containing the exogenously added DNA by, for example, drug selection or deficiency in an essential nutrient which is complemented by the selectable marker carried on the expression vector or co-transfected into the host cell. *P. methanolica* cells are cultured in a medium comprising adequate sources of carbon, nitrogen and trace nutrients at a temperature of about 25°C to 35°C. Liquid cultures are provided with sufficient aeration by conventional means, such as shaking of small flasks or sparging of fermentors. A preferred culture medium for *P. methanolica* is YEPD (2% D-glucose, 2% Bacto™ Peptone (Difco Laboratories, Detroit, MI), 1% Bacto™ yeast extract (Difco Laboratories), 0.004% adenine and 0.006% L-leucine).

The proteins of the present invention can also comprise non-naturally occurring amino acid residues. Non-naturally occurring amino acids include, without limitation, *trans*-3-methylproline, 2,4-methanoproline, *cis*-4-hydroxyproline, *trans*-4-hydroxyproline, *N*-methylglycine, *allo*-threonine, methylthreonine, hydroxyethylcysteine, hydroxyethylhomocysteine, nitroglutamine, homoglutamine, pipercolic acid, thiazolidine carboxylic acid, dehydroproline, 3- and 4-methylproline, 3,3-dimethylproline, *tert*-leucine, norvaline, 2-azaphenylalanine, 3-azaphenylalanine, 4-azaphenylalanine, and 4-fluorophenylalanine. Several methods are known in the art for incorporating non-

naturally occurring amino acid residues into proteins. For example, an *in vitro* system can be employed wherein nonsense mutations are suppressed using chemically aminoacylated suppressor tRNAs. Methods for synthesizing amino acids and aminoacylating tRNA are known in the art. Transcription and translation of plasmids containing nonsense mutations is carried out in a cell-free system comprising an *E. coli* S30 extract and commercially available enzymes and other reagents. Proteins are purified by chromatography. See, for example, Robertson et al., J. Am. Chem. Soc. 113:2722, 1991; Ellman et al., Methods Enzymol. 202:301, 1991; Chung et al., Science 259:806-9, 1993; and Chung et al., Proc. Natl. Acad. Sci. USA 90:10145-9, 1993). In a second method, translation is carried out in *Xenopus* oocytes by microinjection of mutated mRNA and chemically aminoacylated suppressor tRNAs (Turcatti et al., J. Biol. Chem. 271:19991-8, 1996). Within a third method, *E. coli* cells are cultured in the absence of a natural amino acid that is to be replaced (e.g., phenylalanine) and in the presence of the desired non-naturally occurring amino acid(s) (e.g., 2-azaphenylalanine, 3-azaphenylalanine, 4-azaphenylalanine, or 4-fluorophenylalanine). The non-naturally occurring amino acid is incorporated into the protein in place of its natural counterpart. See, Koide et al., Biochem. 33:7470-6, 1994. Naturally occurring amino acid residues can be converted to non-naturally occurring species by *in vitro* chemical modification. Chemical modification can be combined with site-directed mutagenesis to further expand the range of substitutions (Wynn and Richards, Protein Sci. 2:395-403, 1993).

A limited number of non-conservative amino acids, amino acids that are not encoded by the genetic code, non-naturally occurring amino acids, and unnatural amino acids may be substituted for UMLR amino acid residues.

It is preferred to purify the polypeptides of the present invention to  $\geq 80\%$  purity, more preferably to  $\geq 90\%$  purity, even more preferably  $\geq 95\%$  purity, and particularly preferred is a pharmaceutically pure state, that is greater than 99.9% pure with respect to contaminating macromolecules, particularly other proteins and nucleic acids, and free of infectious and pyrogenic agents. Preferably, a purified polypeptide is substantially free of other polypeptides, particularly other polypeptides of animal origin.

Expressed recombinant UMLR proteins (including chimeric polypeptides and multimeric proteins) are purified by conventional protein purification methods,

typically by a combination of chromatographic techniques. See, in general, Affinity Chromatography: Principles & Methods, Pharmacia LKB Biotechnology, Uppsala, Sweden, 1988; and Scopes, Protein Purification: Principles and Practice, Springer-Verlag, New York, 1994. Proteins comprising a polyhistidine affinity tag (typically about 6 histidine residues) are purified by affinity chromatography on a nickel chelate resin. See, for example, Houchuli et al., *Bio/Technol.* 6: 1321-1325, 1988. Proteins comprising a glu-glu tag can be purified by immunoaffinity chromatography according to conventional procedures. See, for example, Grussenmeyer et al., *ibid.* Maltose binding protein fusions are purified on an amylose column according to methods known in the art.

The polypeptides of the present invention can be isolated by a combination of procedures including, but not limited to, anion and cation exchange chromatography, size exclusion, and affinity chromatography. For example, immobilized metal ion adsorption (IMAC) chromatography can be used to purify histidine-rich proteins, including those comprising polyhistidine tags. Briefly, a gel is first charged with divalent metal ions to form a chelate (Sulkowski, Trends in Biochem. 3:1-7, 1985). Histidine-rich proteins will be adsorbed to this matrix with differing affinities, depending upon the metal ion used, and will be eluted by competitive elution, lowering the pH, or use of strong chelating agents. Other methods of purification include purification of glycosylated proteins by lectin affinity chromatography and ion exchange chromatography (Methods in Enzymol., Vol. 182, "Guide to Protein Purification", M. Deutscher, (ed.), Acad. Press, San Diego, 1990, pp.529-39). Within additional embodiments of the invention, a fusion of the polypeptide of interest and an affinity tag (e.g., maltose-binding protein, an immunoglobulin domain) may be constructed to facilitate purification.

UMLR polypeptides can also be prepared through chemical synthesis according to methods known in the art, including exclusive solid phase synthesis, partial solid phase methods, fragment condensation or classical solution synthesis. See, for example, Merrifield, *J. Am. Chem. Soc.* 85:2149, 1963; Stewart et al., Solid Phase Peptide Synthesis (2nd edition), Pierce Chemical Co., Rockford, IL, 1984; Bayer and Rapp, *Chem. Pept. Prot.* 3:3, 1986; and Atherton et al., Solid Phase Peptide Synthesis:

A Practical Approach, IRL Press, Oxford, 1989. In vitro synthesis is particularly advantageous for the preparation of smaller polypeptides.

Using methods known in the art, UMLR proteins can be prepared as monomers or multimers; glycosylated or non-glycosylated; pegylated or non-pegylated; and may or may not include an initial methionine amino acid residue.

The activity of UMLR polypeptides can be measured using a variety of assays that measure, for example, cell-cell interactions, apoptosis, inflammation, anti-infective capabilities and other biological functions associated with TNFR family members or with TNFR interactions, such as, differentiation, and proliferation for example. Of particular interest is a change in inflammation. Assays measuring inflammation are well known in the art.

Proteins, including alternatively spliced peptides, of the present invention are useful for tumor suppression, immunologic recognition, and growth and differentiation either working in isolation, or in conjunction with other molecules (growth factors, cytokines, etc.) in uterus, as well as in melanoma, osteosarcoma, breast carcinoma, and lymphoma, and in tumors of the lung. Alternative splicing of UMLR may cell-type specific and confer activity to specific tissues.

As polynucleotides of the present invention have been identified in lung carcinoma, melanoma, osteosarcoma, breast carcinoma, and lymphoma cells, polynucleotides of UMLR can be used to detect these cell types by hybridization, or with UMLR binding proteins. Similarly antibodies to the UMLR polypeptides can detect cells expressing the surface bound UMLR receptor. Such detection can be useful to determine metastasis, disease stage, or primary diagnosis of disease. Additionally, proliferation, differentiation and/or apoptosis of these cell types can be modulated by contacting the cells with UMLR binding proteins, including a UMLR-cognate ligand, or an antibody to UMLR, for example. The proliferation, differentiation and/or apoptosis of such cells can also be mediated by UMLR polypeptides, or fragments thereof, as an antagonist of cell signaling by binding the ligand and thereby reducing the interaction between ligand and membrane-bound UMLR proteins. The effects of the modulation of UMLR molecules on proliferation, differentiation and/or apoptosis can be measured, for example, by the presence or absence of cell markers specific to these cell types, or by

measuring other manifestations of these diseases. For example, UMLR polypeptides or fragments thereof, as well as UMLR binding proteins can be administered to cells of lung carcinoma and inhibition of the carcinoma can be monitored by conventional imaging techniques.

5                   Another assay of interest measures or detects changes in proliferation, differentiation, and development. Additionally, the effects of a UMLR polypeptides on cell-cell interactions of fibroblasts, epithelial cells, tumor cells and cells of the genito-urinary tract in particular, would be of interest to measure. Yet other assays examines changes in apoptosis, and intracellular signaling.

10                   The activity of molecules of the present invention can be measured using a variety of assays that, for example, measure neogenesis or hyperplasia (i.e., proliferation) of tissues of the uterus, lung carcinoma, or breast carcinoma, as well as cells of melanoma, osteosarcoma, and lymphoma. Additional activities likely associated with the polypeptides of the present invention include proliferation of endothelial cells,  
15   fibroblasts, and lymphoid cells directly or indirectly through other growth factors; action as a chemotaxic factor for endothelial cells, fibroblasts and/or phagocytic cells; osteogenic factor; and factor for expanding mesenchymal stem cell and precursor populations.

                  The UMLR polypeptides of the present invention can be used to study  
20   proliferation or differentiation in uterus, as well as in melanoma, osteosarcoma, breast carcinoma, and lymphoma, and in tumors of the lung. Such methods of the present invention generally comprise incubating cells derived from these tissues in the presence and absence of UMLR polypeptide, monoclonal antibody, agonist or antagonist thereof and observing changes in cell proliferation or differentiation. Cell lines from these  
25   tissues are commercially available from, for example, American Type Culture Collection (Manasas, VA).

                  Proliferation can be measured using cultured uterine or lung cells or *in vivo* by administering molecules of the claimed invention to an appropriate animal model. Generally, proliferative effects are observed as an increase in cell number and  
30   therefore, may include inhibition of apoptosis, as well as mitogenesis. Cultured cells include uterine fibroblasts, lung tumors, and melanoma, osteosarcoma, breast carcinoma,



and lymphoma, as well as from primary cultures. Established cell lines are easily identifiable by one skilled in the art and are available from ATCC (Manassas, VA). Assays measuring cell proliferation are well known in the art. For example, assays measuring proliferation include such assays as chemosensitivity to neutral red dye

5 (Cavanaugh et al., Investigational New Drugs 8:347-354, 1990), incorporation of radiolabelled nucleotides (Cook et al., Analytical Biochem. 179:1-7, 1989), incorporation of 5-bromo-2'-deoxyuridine (BrdU) in the DNA of proliferating cells (Porstmann et al., J. Immunol. Methods 82:169-179, 1985), and use of tetrazolium salts (Mosmann, J. Immunol. Methods 65:55-63, 1983; Alley et al., Cancer Res. 48:589-601,

10 1988; Marshall et al., Growth Reg. 5:69-84, 1995; and Scudiero et al., Cancer Res. 48:4827-4833, 1988).

To determine if UMLR is a chemotractant *in vivo*, UMLR can be given by intradermal or intraperitoneal injection. Characterization of the accumulated leukocytes at the site of injection can be determined using lineage specific cell surface markers and

15 fluorescence immunocytometry or by immunohistochemistry (Jose, J. Exp. Med. 179:881-87, 1994). Release of specific leukocyte cell populations from bone marrow into peripheral blood can also be measured after UMLR injection.

Differentiation is a progressive and dynamic process, beginning with pluripotent stem cells and ending with terminally differentiated cells. Pluripotent stem

20 cells that can regenerate without commitment to a lineage express a set of differentiation markers that are lost when commitment to a cell lineage is made. Progenitor cells express a set of differentiation markers that may or may not continue to be expressed as the cells progress down the cell lineage pathway toward maturation. Differentiation markers that are expressed exclusively by mature cells are usually functional properties

25 such as cell products, enzymes to produce cell products and receptors and receptor-like complementary molecules. The stage of a cell population's differentiation is monitored by identification of markers present in the cell population. For example, myocytes, osteoblasts, adipocytes, chondrocytes, fibroblasts and reticular cells are believed to originate from a common mesenchymal stem cell (Owen et al., Ciba Fdn. Symp. 136:42-

30 46, 1988). Markers for mesenchymal stem cells have not been well identified (Owen et al., J. of Cell Sci. 87:731-738, 1987), so identification is usually made at the progenitor

and mature cell stages. The novel polypeptides of the present invention are useful for studies to isolate mesenchymal stem cells and uterine myocyte progenitor cells, both *in vivo* and *ex vivo*.

There is evidence to suggest that factors that stimulate specific cell types  
 5 down a pathway towards terminal differentiation or dedifferentiation affect the entire cell population originating from a common precursor or stem cell. Thus, UMLR polypeptides may stimulate inhibition or proliferation of endocrine and exocrine cells of the uterus, as well as in melanoma, osteosarcoma, breast carcinoma, and lymphoma, and in tumors of the lung.

10 Molecules of the present invention may, while stimulating proliferation or differentiation of uterine fibroblasts, inhibit proliferation or differentiation of adipocytes, by virtue of their effect on common precursor/stem cells. The novel polypeptides of the present invention are useful to study neural and epithelial stem cells and uterus, as well as in melanoma, osteosarcoma, breast carcinoma, and lymphoma, and  
 15 in tumors in lung progenitor cells, both *in vivo* and *ex vivo*.

Assays measuring differentiation include, for example, measuring cell-surface markers associated with stage-specific expression of a tissue, enzymatic activity, functional activity or morphological changes (Watt, FASEB, 5:281-284, 1991; Francis, Differentiation 57:63-75, 1994; Raes, Adv. Anim. Cell Biol. Technol. Bioprocesses,  
 20 161-171, 1989).

Proteins, including alternatively spliced peptides, and fragments, of the present invention are useful for studying cell-cell interactions, fertilization, development, immune recognition, growth control, tumor suppression, and embryo maturation. UMLR molecules, variants, and fragments can be applied in isolation, or in conjunction  
 25 with other molecules (growth factors, cytokines, etc.) in uterus, as well as in melanoma, osteosarcoma, breast carcinoma, and lymphoma, and in tumors of the lung.

Proteins of the present invention are useful for delivery of therapeutic agents such as, but not limited to, proteases, radionuclides, chemotherapy agents, and small molecules. Effects of these therapeutic agents can be measured *in vitro* using  
 30 cultured cells, *ex vivo* on tissue slices, or *in vivo* by administering molecules of the claimed invention to the appropriate animal model. An alternative *in vivo* approach for

assaying proteins of the present invention involves viral delivery systems. Exemplary viruses for this purpose include adenovirus, herpesvirus, lentivirus, vaccinia virus and adeno-associated virus (AAV). Adenovirus, a double-stranded DNA virus, is currently the best studied gene transfer vector for delivery of heterologous nucleic acid (for a review, see T.C. Becker et al., Meth. Cell Biol. 43:161-89, 1994; and J.T. Douglas and D.T. Curiel, Science & Medicine 4:44-53, 1997). The adenovirus system offers several advantages: adenovirus can (i) accommodate relatively large DNA inserts; (ii) be grown to high-titer; (iii) infect a broad range of mammalian cell types; and (iv) be used with a large number of available vectors containing different promoters. Also, because adenoviruses are stable in the bloodstream, they can be administered by intravenous injection.

By deleting portions of the adenovirus genome, larger inserts (up to 7 kb) of heterologous DNA can be accommodated. These inserts can be incorporated into the viral DNA by direct ligation or by homologous recombination with a co-transfected plasmid. In an exemplary system, the essential E1 gene has been deleted from the viral vector, and the virus will not replicate unless the E1 gene is provided by the host cell (the human 293 cell line is exemplary). When intravenously administered to intact animals, adenovirus primarily targets the liver. If the adenoviral delivery system has an E1 gene deletion, the virus cannot replicate in the host cells. However, the host's tissue (e.g., liver) will express and process (and, if a secretory signal sequence is present, secrete) the heterologous protein. Secreted proteins will enter the circulation in the highly vascularized liver, and effects on the infected animal can be determined.

Moreover, adenoviral vectors containing various deletions of viral genes can be used in an attempt to reduce or eliminate immune responses to the vector. Such adenoviruses are E1 deleted, and in addition contain deletions of E2A or E4 (Lusky, M. et al., J. Virol. 72:2022-2032, 1998; Raper, S.E. et al., Human Gene Therapy 9:671-679, 1998). In addition, deletion of E2b is reported to reduce immune responses (Amalfitano, A. et al., J. Virol. 72:926-933, 1998). Moreover, by deleting the entire adenovirus genome, very large inserts of heterologous DNA can be accommodated. Generation of so called "gutless" adenoviruses where all viral genes are deleted are particularly

advantageous for insertion of large inserts of heterologous DNA. For review, see Yeh, P. and Perricaudet, M., FASEB J. **11**:615-623, 1997.

The adenovirus system can also be used for protein production *in vitro*. By culturing adenovirus-infected non-293 cells under conditions where the cells are not rapidly dividing, the cells can produce proteins for extended periods of time. For instance, BHK cells are grown to confluence in cell factories, then exposed to the adenoviral vector encoding the secreted protein of interest. The cells are then grown under serum-free conditions, which allows infected cells to survive for several weeks without significant cell division. Alternatively, adenovirus vector infected 293S cells can be grown in suspension culture at relatively high cell density to produce significant amounts of protein (see Garnier et al., Cytotechnol. **15**:145-55, 1994). With either protocol, an expressed, secreted heterologous protein can be repeatedly isolated from the cell culture supernatant. Within the infected 293S cell production protocol, non-secreted proteins may also be effectively obtained.

As a soluble or cell-surface protein, the activity of UMLR polypeptide or a peptide to which UMLR binds, can be measured by a silicon-based biosensor microphysiometer which measures the extracellular acidification rate or proton excretion associated with cell-surface protein interactions and subsequent physiologic cellular responses. An exemplary device is the Cytosensor™ Microphysiometer manufactured by Molecular Devices, Sunnyvale, CA. A variety of cellular responses, such as cell proliferation, ion transport, energy production, inflammatory response, regulatory and receptor activation, and the like, can be measured by this method. See, for example, McConnell, H.M. et al., Science **257**:1906-1912, 1992; Pitchford, S. et al., Meth. Enzymol. **228**:84-108, 1997; Arimilli, S. et al., J. Immunol. Meth. **212**:49-59, 1998; Van Liefde, I. et al., Eur. J. Pharmacol. **346**:87-95, 1998. The microphysiometer can be used for assaying adherent or non-adherent eukaryotic or prokaryotic cells. By measuring extracellular acidification changes in cell media over time, the microphysiometer directly measures cellular responses to various stimuli, including proteins, agonists, and antagonists which affect a UMLR-mediated pathway. UMLR-responsive eukaryotic cells comprise cells into which a polynucleotide for UMLR has been transfected creating a cell that is responsive to activation of UMLR; or cells naturally responsive to activation

of UMLR . Differences, measured by a change in the response of cells exposed to UMLR activation, relative to a control not exposed to UMLR activation, are a direct measurement of UMLR-mediated cellular responses. Moreover, such UMLR-mediated responses can be assayed under a variety of stimuli. The present invention provides a method of identifying agonists and antagonists of UMLR protein, comprising providing  
 5 cells responsive to activation of UMLR polypeptide, culturing a first portion of the cells in the absence of a test compound, culturing a second portion of the cells in the presence of a test compound, and detecting a change in a cellular response of the second portion of the cells as compared to the first portion of the cells. The change in cellular response  
 10 is shown as a measurable change in extracellular acidification rate. Moreover, culturing a third portion of the cells in the presence of UMLR polypeptide and the absence of a test compound provides a positive control for the UMLR -responsive cells, and a control to compare the agonist activity of a test compound with that of the UMLR polypeptide. Antagonists of UMLR can be identified by exposing the cells to UMLR protein in the  
 15 presence and absence of the test compound, whereby a reduction in UMLR-stimulated activity is indicative of antagonist activity in the test compound.

Similarly, the microphysiometer, can be used to rapidly identify cells, tissues, or cell lines which activate a UMLR-stimulated pathway. Such tissues and cell lines can be used to identify ligands, antagonists and agonists of UMLR polypeptide as  
 20 described above. Using similar methods, cells expressing UMLR can be used to identify cells which stimulate or block a UMLR-signaling pathway.

UMLR shares homology with OPG, a soluble TNF receptor involved in the regulation of bone density (Simonet et al., *ibid.*). Well established animal models are available to test the *in vivo* efficacy of UMLR polypeptides for certain disease states,  
 25 such as bone-related disorders. For example, the hypocalcemic rat model can be used to determine the effect of UMLR on serum calcium, and the ovariectomized rat or mouse can be used as a model system for osteoporosis. Bone changes seen in these models and in humans during the early stages of estrogen deficiency are qualitatively similar.

In view of the tissue distribution (uterus, lung carcinoma, breast  
 30 carcinoma, and thymus and pituitary, as well as cells of melanoma, osteosarcoma, and lymphoma) observed for UMLR expression, agonists (including the native cysteine-rich

pseudo-repeat and cytoplasmic domains, and antagonists have enormous potential in both *in vitro* and *in vivo* applications. Compounds identified as UMLR agonists and antagonists are useful for studying cell-cell interactions, apoptosis, tumor proliferation and suppression, infection, and inflammation *in vitro* and *in vivo*. For example, UMLR and agonist compounds are useful as components of defined cell culture media, and may be used alone or in combination with cytokines and hormones to replace serum that is commonly used in cell culture. Agonists are thus useful in specifically promoting the growth and/or development of cells of the myeloid and lymphoid lineages in culture. Additionally, UMLR polypeptides and UMLR agonists, including small molecules are useful as a research reagent, such as for the expansion, differentiation, proliferation, and/or cell-cell interactions of uterus, as well as in melanoma, osteosarcoma, breast carcinoma, and lymphoma, and in tumors of the lung. UMLR polypeptides are added to tissue culture media for these cell types.

Compounds identified as UMLR agonists are useful for modifying the proliferation and development of target cells *in vitro* and *in vivo*. For example, agonist compounds are useful alone or in combination with other cytokines and hormones as components of defined cell culture media. Agonists are thus useful in specifically mediating the growth and/or development of UMLR -bearing T lymphocytes or other UMLR-bearing cells in culture. Agonists and antagonists may also prove useful in the study of effector functions of T lymphocytes, in particular T lymphocyte activation and differentiation. Antagonists are useful as research reagents for characterizing ligand-receptor interaction.

As a member of the TNFR family, UMLR polypeptides are likely to be involved in the immune response to infection. Lymphotoxin-beta receptor, another member of this receptor family, has been shown to regulate HIV-1 replication. (Marshall, W.L. et al., J. Immun. 162: 6016-6023, 1999). Further, it has been shown that cosignaling via the lymphotoxin-beta receptor and TNF- receptors is probably involved in the modulation of HIV-1 replication and the subsequent determination of HIV-1 viral burden in monocytes. UMLR polypeptides, agonists, and antagonists can be used to treat microbial infections. Such infections include bacterial, yeast, and viral infections. Anti-microbial activity of proteins is evaluated by techniques that are known in the art. For

example, anti-microbial activity can be assayed by evaluating the sensitivity of microbial cell cultures to test agents and by evaluating the protective effect of test agents on infected mice. See, for example, Musiek et al., *Antimicrob. Agents Chemother.* 3:40, 1973. Antiviral activity can also be assessed by protection of mammalian cell cultures.

5 Known techniques for evaluating anti-microbial activity include, for example, Barsum et al., *Eur. Respir. J.* 8:709-714, 1995; Sandovsky-Losica et al., *J. Med. Vet. Mycol (England)* 28:279-287, 1990; Mehentee et al., *J. Gen. Microbiol (England)* 135(2):2181-2188, 1989; and Segal and Savage, *J. Med. Vet. Mycol.* 24:477-479, 1986. Assays specific for anti-viral activity include, for example, those described by Daher et al., *J.*

10 *Virol.* 60:1068-1074, 1986. Similarly, assays measuring HIV-1 viral burden on cells can be used.

Shock is a manifestation of infection. Studies show that increased serum TNF levels are associated with high mortality rates. See Wage, A. et al., *Lancet* i:355-357, 1987, and Girardin et al., *New. Eng. J. Med.* 39: 397-400, 1988. Soluble TNF

15 receptors, including UMLR polypeptides of the present invention, may be useful to reduce serum concentrations of TNF, and minimize the effects of sepsis.

The invention also provides antagonists, which either bind to UMLR polypeptides or, alternatively, to a ligand to which UMLR polypeptides bind, thereby inhibiting or eliminating the function of UMLR. Such UMLR antagonists would include

20 antibodies; polypeptides which bind either to the UMLR polypeptide or to its ligand; natural or synthetic analogs of UMLR ligands which retain the ability to bind the receptor but do not result in either ligand or receptor signaling. Such analogs could be peptides or peptide-like compounds. Natural or synthetic small molecules which bind to UMLR polypeptides and prevent signaling are also contemplated as antagonists. Also

25 contemplated are soluble UMLR receptors. As such, UMLR antagonists would be useful as therapeutics for treating certain disorders where blocking signal from either a UMLR receptor or ligand would be beneficial. Antagonists are useful as research reagents for characterizing ligand-receptor interaction.

UMLR polypeptides may be used within diagnostic systems to detect the

30 presence of ligand polypeptides. Antibodies or other agents that specifically bind to UMLR may also be used to detect the presence of circulating or membrane bound

receptor or ligand polypeptides. Such detection methods are well known in the art and include, for example, enzyme-linked immunosorbent assay (ELISA) and radioimmunoassay. Immunohistochemically labeled UMLR antibodies can be used to detect UMLR receptor and/or ligands in tissue samples. UMLR levels can also be monitored by such methods as RT-PCR, where UMLR mRNA can be detected and quantified. The information derived from such detection methods would provide insight into the significance of UMLR polypeptides in various diseases, and as such would serve as diagnostic tools for diseases for which altered levels of UMLR are significant. Altered levels of UMLR receptor polypeptides may be indicative of pathological conditions including cancer, autoimmune disorders, bone disorders, inflammation and immunodeficiencies.

Antagonists are also useful as research reagents for characterizing sites of interactions between members of complement/anti-complement pairs as well as sites of cell-cell interactions. Inhibitors of UMLR activity (UMLR antagonists) include anti-UMLR antibodies and soluble UMLR polypeptides (such as described above), as well as other peptidic and non-peptidic agents (including ribozymes).

UMLR can also be used to identify inhibitors (antagonists) of its activity. Test compounds are added to the assays disclosed herein to identify compounds that inhibit the activity of UMLR. In addition to those assays disclosed herein, samples can be tested for inhibition of UMLR activity within a variety of assays designed to measure receptor/ligand binding or the stimulation/inhibition of UMLR-dependent cellular responses. For example, UMLR-responsive cell lines can be transfected with a reporter gene construct that is responsive to a UMLR-stimulated cellular pathway. Reporter gene constructs of this type are known in the art, and will generally comprise a DNA response element operably linked to a gene encoding an assayable protein, such as luciferase, or a metabolite, such as cyclic AMP. DNA response elements can include, but are not limited to, cyclic AMP response elements (CRE), hormone response elements (HRE), insulin response element (IRE) (Nasrin et al., Proc. Natl. Acad. Sci. USA **87**:5273-7, 1990) and serum response elements (SRE) (Shaw et al. Cell **56**: 563-72, 1989). Cyclic AMP response elements are reviewed in Roestler et al., J. Biol. Chem. **263** (19):9063-6; 1988 and Habener, Molec. Endocrinol. **4** (8):1087-94; 1990. Hormone response



elements are reviewed in Beato, Cell 56:335-44; 1989. Such a reporter gene construct would contain a cysteine-rich pseudo-repeat that, upon binding a TNF, would signal intracellularly through, for example, a SRE reporter. Candidate compounds, solutions, mixtures or extracts are tested for the ability to inhibit the activity of UMLR on the target cells, as evidenced by a decrease in UMLR stimulation of reporter gene expression. Assays of this type will detect compounds that directly block UMLR binding to a cell-surface protein, i.e., ligand, or the anti-complementary member of a complementary/anti-complementary pair, as well as compounds that block processes in the cellular pathway subsequent to complement/anti-complement binding. In the alternative, compounds or other samples can be tested for direct blocking of UMLR binding to a ligand using UMLR tagged with a detectable label (e.g., <sup>125</sup>I, biotin, horseradish peroxidase, FITC, and the like). Within assays of this type, the ability of a test sample to inhibit the binding of labeled UMLR to the TNF is indicative of inhibitory activity, which can be confirmed through secondary assays. TNFs used within binding assays may be cellular TNFs, soluble TNFs, or isolated, immobilized TNFs.

Also, UMLR polypeptides, agonists or antagonists thereof may be therapeutically useful for promoting wound healing, for example, in uterus, as well as in melanoma, osteosarcoma, breast carcinoma, and lymphoma, and in tumors of the lung. To verify the presence of this capability in UMLR polypeptides, agonists or antagonists of the present invention, such UMLR polypeptides, agonists or antagonists are evaluated with respect to their ability to facilitate wound healing according to procedures known in the art. If desired, UMLR polypeptide performance in this regard can be compared to growth factors, such as EGF, NGF, TGF- $\alpha$ , TGF- $\beta$ , insulin, IGF-I, IGF-II, fibroblast growth factor (FGF) and the like. In addition, UMLR polypeptides or agonists or antagonists thereof may be evaluated in combination with one or more growth factors to identify synergistic effects.

The activity of agonists and antagonists can be determined by activity assays which determine the potency of receptor/ligand engagement. Stably transfected cell lines, which co-express high levels of reporter gene constructs for NfKB, NFAT-1 and AP-1 can be made which express ztnfr11. A ztnfr11 ligand can be found to signal

through the reporter genes in these constructs. Soluble ztnfr11 and antibodies can be used to measure binding.

A UMLR ligand-binding polypeptide can also be used for purification of  
 5 ligand. The polypeptide is immobilized on a solid support, such as beads of agarose, cross-linked agarose, glass, cellulosic resins, silica-based resins, polystyrene, cross-linked polyacrylamide, or like materials that are stable under the conditions of use. Methods for linking polypeptides to solid supports are known in the art, and include amine chemistry, cyanogen bromide activation, N-hydroxysuccinimide activation,  
 10 epoxide activation, sulfhydryl activation, and hydrazide activation. The resulting medium will generally be configured in the form of a column, and fluids containing ligands are passed through the column one or more times to allow ligands to bind to the receptor polypeptide. The ligand is then eluted using changes in salt concentration, chaotropic agents (guanidine HCl), or pH to disrupt ligand-receptor binding.

15 An assay system that uses a ligand-binding receptor (or an antibody, one member of a complementary/ anti-complementary pair or other cell-surface binding protein) or a binding fragment thereof, and a commercially available biosensor instrument (BIAcore, Pharmacia Biosensor, Piscataway, NJ) may be advantageously employed. Such receptor, antibody, member of a complement/anti-complement pair or  
 20 fragment is immobilized onto the surface of a receptor chip. Use of this instrument is disclosed by Karlsson, J. Immunol. Methods 145:229-40, 1991 and Cunningham and Wells, J. Mol. Biol. 234:554-63, 1993. A receptor, antibody, member or fragment is covalently attached, using amine or sulfhydryl chemistry, to dextran fibers that are attached to gold film within the flow cell. A test sample is passed through the cell. If a  
 25 ligand, epitope, or opposite member of the complementary/anti-complementary pair is present in the sample, it will bind to the immobilized ligand, antibody or member, respectively, causing a change in the refractive index of the medium, which is detected as a change in surface plasmon resonance of the gold film. This system allows the determination of on- and off-rates, from which binding affinity can be calculated, and  
 30 assessment of .

Ligand binding receptor polypeptides can also be used within other assay systems known in the art. Such systems include Scatchard analysis for determination of

binding affinity (see Scatchard, Ann. NY Acad. Sci. 51: 660-72, 1949) and calorimetric assays (Cunningham et al., Science 253:545-48, 1991; Cunningham et al., Science 245:821-25, 1991).

A "soluble protein" is a protein that is not bound to a cell membrane.

- 5 Soluble proteins are most commonly ligand-binding receptor polypeptides that lack transmembrane and cytoplasmic domains. Soluble proteins can comprise additional amino acid residues, such as affinity tags that provide for purification of the polypeptide or provide sites for attachment of the polypeptide to a substrate, or immunoglobulin constant region sequences. Many cell-surface proteins have naturally occurring, soluble
- 10 counterparts that are produced by proteolysis or translated from alternatively spliced mRNAs. Proteins are said to be substantially free of transmembrane and intracellular polypeptide segments when they lack sufficient portions of these segments to provide membrane anchoring or signal transduction, respectively. The sequence as shown in SEQ ID NO:38 from residue 1 to residue 173 is a soluble protein. Fragments of the
- 15 polypeptide as shown in SEQ ID NO:38 can also be used as a soluble protein. The partial cysteine-rich domain of SEQ ID NO:2, may be a complete domain in the polypeptide as shown in SEQ ID NO:38. Thus, this cycteine-rich domain would be from residue 85 to residue 147, or from residue 85 to residue 163 as shown in SEQ ID NO:38. Additionally, there are two possible dibasic cleavage sites in SEQ ID NO:38 at residues
- 20 153-154, and residues 164-165. Thus, soluble fragments of ztnfr11 include the polypeptide from residue 1 to residue 147 fo SEQ ID NO:38; from residue 1 to 154 of SEQ ID NO:38; from residue 1 to residue 163 of SEQ ID NO:38; from residue 1 to 165 of SEQ ID NO:38; and from residue 1 to 173 of SEQ ID NO:38.

- 25 Soluble forms of UMLR polypeptides may act as antagonsits to or agonists of UMLR polypeptides, and would be useful to modulate the effects of UMLR in uterus, as well as in melanoma, osteosarcoma, breast carcinoma, and lymphoma, and in tumors of the lung. Thus, the isoform of UMLR that does not contain a transmembrane domain (i.e., the polypeptide of residues 1 to 129 of SEQ ID NO:2; residues 1 to 173 of SEQ ID NO:38; residues 1 to 147 of SEQ ID NO:38; residues 1 to
- 30 154 of SEQ ID NO:38; and residues 1 to 165 of SEQ ID NO:38;) will be soluble, and may act as an agonist or antagonist of UMLR activity. Since polypeptides of this nature

are not anchored to the membrane, they can act at sites distant from the tissues in which they are expressed. Thus, the activity of the soluble form of UMLR polypeptides can be more wide spread than its membrane-anchored counterpart. Both isoforms would be useful in studying the effects of the present invention *in vitro* an *in vivo*.

5                   Molecules of the present invention can be used to identify and isolate TNFs, or members of complement/anti-complement pairs involved in cell-cell interactions. For example, proteins and peptides of the present invention can be immobilized on a column and membrane preparations run over the column (Immobilized Affinity Ligand Techniques, Hermanson et al., eds., Academic Press, San Diego, CA, 10 1992, pp.195-202). Proteins and peptides can also be radiolabeled (Methods in Enzymol., vol. 182, "Guide to Protein Purification", M. Deutscher, ed., Acad. Press, San Diego, 1990, 721-37) or photoaffinity labeled (Brunner et al., Ann. Rev. Biochem. 62:483-514, 1993 and Fedan et al., Biochem. Pharmacol. 33:1167-80, 1984) and specific cell-surface proteins can be identified.

15                   The molecules of the present invention will be useful in modulating abnormal cell growth, inflammation, apoptosis, proliferation and differentiation. The polypeptides, nucleic acid and/or antibodies of the present invention can be used in treatment of disorders associated with infection, tumor growth, immunodeficiency, auto- immunity, and fertility. The molecules of the present invention can be used to modulate 20 apoptosis, cell adhesion, cell fusion, and signaling or to treat or prevent development of pathological conditions in such diverse tissue as uterus, lung carcinoma, and breast carcinoma, as well as in cells of melanoma, osteosarcoma, and lymphoma. In particular, certain diseases may be amenable to such diagnosis, treatment or prevention. These diseases include, but are not limited to, melanoma, osteosarcoma, lung carcinoma, breast 25 carcinoma, endometriosis, immunodeficiency, and infection of the genito-urinary tract. The molecules of the present invention can be used to modulate inhibition and proliferation of tissues in the uterus, lung carcinoma and breast carcinoma, as well as cells of melanoma, osteosarcoma, and lymphoma.

30                   Polynucleotides encoding UMLR polypeptides are useful within gene therapy applications where it is desired to increase or inhibit UMLR activity. If a mammal has a mutated or absent UMLR gene, the UMLR gene can be introduced into

the cells of the mammal. In one embodiment, a gene encoding a UMLR polypeptide is introduced *in vivo* in a viral vector. Such vectors include an attenuated or defective DNA virus, such as, but not limited to, herpes simplex virus (HSV), papillomavirus, Epstein Barr virus (EBV), adenovirus, adeno-associated virus (AAV), and the like.

5 Defective viruses, which entirely or almost entirely lack viral genes, are preferred. A defective virus is not infective after introduction into a cell. Use of defective viral vectors allows for administration to cells in a specific, localized area, without concern that the vector can infect other cells. Examples of particular vectors include, but are not limited to, a defective herpes simplex virus 1 (HSV1) vector (Kaplitt et al., Molec. Cell.

10 Neurosci. 2:320-30, 1991); an attenuated adenovirus vector, such as the vector described by Stratford-Perricaudet et al., J. Clin. Invest. 90:626-30, 1992; and a defective adeno-associated virus vector (Samulski et al., J. Virol. 61:3096-101, 1987; Samulski et al., J. Virol. 63:3822-8, 1989).

In another embodiment, a UMLR gene can be introduced in a retroviral

15 vector, e.g., as described in Anderson et al., U.S. Patent No. 5,399,346; Mann et al. Cell 33:153, 1983; Temin et al., U.S. Patent No. 4,650,764; Temin et al., U.S. Patent No. 4,980,289; Markowitz et al., J. Virol. 62:1120, 1988; Temin et al., U.S. Patent No. 5,124,263; International Patent Publication No. WO 95/07358, published March 16, 1995 by Dougherty et al.; and Kuo et al., Blood 82:845, 1993. Alternatively, the vector

20 can be introduced by lipofection *in vivo* using liposomes. Synthetic cationic lipids can be used to prepare liposomes for *in vivo* transfection of a gene encoding a marker (Felgner et al., Proc. Natl. Acad. Sci. USA 84:7413-7, 1987; Mackey et al., Proc. Natl. Acad. Sci. USA 85:8027-31, 1988). The use of lipofection to introduce exogenous genes

25 into specific organs *in vivo* has certain practical advantages. Molecular targeting of liposomes to specific cells represents one area of benefit. More particularly, directing transfection to particular cells represents one area of benefit. For instance, directing transfection to particular cell types would be particularly advantageous in a tissue with cellular heterogeneity, such as the pancreas, liver, kidney, and brain. Lipids may be chemically coupled to other molecules for the purpose of targeting. Targeted peptides

30 (e.g., hormones or neurotransmitters), proteins such as antibodies, or non-peptide molecules can be coupled to liposomes chemically.

Similarly, the UMLR polynucleotides (SEQ ID NOs:1, 32, 33, and 36) can be used to target specific tissues such as tissues of the uterus, lung carcinoma and breast carcinoma, as well as in cells of melanoma, osteosarcoma, and lymphoma. It is possible to remove the target cells from the body; to introduce the vector as a naked DNA plasmid; and then to re-implant the transformed cells into the body. Naked DNA vectors for gene therapy can be introduced into the desired host cells by methods known in the art, e.g., transfection, electroporation, microinjection, transduction, cell fusion, DEAE dextran, calcium phosphate precipitation, use of a gene gun or use of a DNA vector transporter. See, e.g., Wu et al., J. Biol. Chem. 267:963-7, 1992; Wu et al., J. Biol. Chem. 263:14621-4, 1988.

Various techniques, including antisense and ribozyme methodologies, can be used to inhibit UMLR gene transcription and translation, such as to inhibit cell proliferation in vivo. Polynucleotides that are complementary to a segment of a UMLR -encoding polynucleotide (e.g., a polynucleotide as set forth in SEQ ID NOs:1 or 3) are designed to bind to UMLR -encoding mRNA and to inhibit translation of such mRNA. Such antisense polynucleotides are used to inhibit expression of UMLR polypeptide-encoding genes in cell culture or in a subject.

Mice engineered to express the UMLR gene, referred to as "transgenic mice," and mice that exhibit a complete absence of UMLR gene function, referred to as "knockout mice," may also be generated (Snouwaert et al., Science 257:1083, 1992), may also be generated (Lowell et al., Nature 366:740-42, 1993; Capecchi, M.R., Science 244: 1288-1292, 1989; Palmiter, R.D. et al. Annu Rev Genet. 20: 465-499, 1986). For example, transgenic mice that over-express UMLR , either ubiquitously or under a tissue-specific or tissue-restricted promoter can be used to ask whether over-expression causes a phenotype. For example, over-expression of a wild-type UMLR polypeptide, polypeptide fragment or a mutant thereof may alter normal cellular processes, resulting in a phenotype that identifies a tissue in which UMLR expression is functionally relevant and may indicate a therapeutic target for the UMLR, its agonists or antagonists. For example, a transgenic mouse to engineer is one that over-expresses the soluble UMLR polypeptide (approximately amino acids 1 to 129 of SEQ ID NO:2; amino acids 1 to 173 of SEQ ID NO:38; amino acids 1 to 147 of SEQ ID NO:38; amino acids 1 to 154 of SEQ

ID NO:38; and amino acids 1 to 165 of SEQ ID NO:38;), or the membrane-bound receptor, (residues 1 to 269 of SEQ ID NO:2; residues 1 to 297 of SEQ ID NO:27; and residues 1 to 299 of SEQ ID NO:35). Moreover, such over-expression may result in a phenotype that shows similarity with human diseases. Similarly, knockout UMLR mice  
5 can be used to determine where UMLR is absolutely required *in vivo*. The phenotype of knockout mice is predictive of the *in vivo* effects of that a UMLR antagonist, such as those described herein, may have. The human UMLR cDNA can be used to isolate murine UMLR mRNA, cDNA and genomic DNA, which are subsequently used to generate knockout mice. These mice may be employed to study the UMLR gene and the  
10 protein encoded thereby in an *in vivo* system, and can be used as *in vivo* models for corresponding human diseases. Moreover, transgenic mice expression of UMLR antisense polynucleotides or ribozymes directed against UMLR, described herein, can be used analogously to transgenic mice described above.

Another use for *in vivo* models includes delivery of an antigen challenge  
15 to the animal followed by administration of soluble ztnfr11 or its ligand, and measuring the T cell response, or the proliferation or decline of ztnfr11-expressing cells.

T-cell dependent and T-cell independent immune response can be measured as described in Perez-Melgosa et al., J. Immunol. 163: 1123-7, 1999.

Pharmacokinetic studies can be used in association with radiolabeled,  
20 soluble ztnfr11 polypeptides or fusions to determine the distribution and half life of such polypeptides *in vivo*. Additionally animal models can be used to determine the effects of soluble ztnfr11 on tumors and tumor development *in vivo*.

Also provided is the use of ztnfr11 polypeptides as surrogate markers for abnormal cell growth, especially, such growth as related to melanoma, breast carcinoma,  
25 lung carcinoma, and cutaneous T-cell lymphoma. Patients having such diseases can be bleed and ztnfr11 soluble receptors and its ligand(s) can be detected in the blood.

UMLR polypeptides, variants, and fragments thereof, may be useful as replacement therapy for disorders associated with cell-cell interactions, including  
30 disorders related to, for example, immunology, fertility, egg maturation, and epithelial disorders, in general.

A less widely appreciated determinant of tissue morphogenesis is the process of cell rearrangement: Both cell motility and cell-cell adhesion are likely to play central roles in morphogenetic cell rearrangements. Cells need to be able to rapidly break and probably simultaneously remake contacts with neighboring cells. See  
 5 Gumbiner, B.M., Cell 69:385-387, 1992. As a secreted protein in tissues of the uterus, lung carcinoma and breast carcinoma, as well as in cells of melanoma, osteosarcoma, and lymphoma. UMLR can play a role in intercellular rearrangement in these and other tissues.

UMLR gene may be useful to as a probe to identify humans who have a  
 10 defective UMLR gene. The strong expression of UMLR in tissues of the uterus, lung carcinoma and breast carcinoma, as well as in cells of melanoma, osteosarcoma, and lymphoma suggests that UMLR polynucleotides or polypeptides can be used as measured as an indication of aberrant growth in these tissues. Thus, polynucleotides and polypeptides of UMLR , and mutations to them, can be used a diagnostic indicators of  
 15 cancer in these tissues.

The polypeptides of the present invention are useful in studying cell adhesion and the role thereof in metastasis and may be useful in preventing metastasis, in particular metastasis in tumors of the uterus, lung, and in melanoma, osteosarcoma, breast carcinoma, and lymphoma. Similarly, polynucleotides and polypeptides of UMLR  
 20 may be used to replace their defective counterparts in tumor or malignant tissues.

The UMLR polypeptide is expressed in tissues of the uterus, lung carcinoma and breast carcinoma as well as in cells of melanoma, osteosarcoma, and lymphoma. Thus, UMLR polypeptide pharmaceutical compositions of the present invention may be useful in prevention or treatment of disorders associated with  
 25 pathological regulation or the expansion of these tissues.

The polynucleotides of the present invention may also be used in conjunction with a regulatable promoter, thus allowing the dosage of delivered protein to be regulated.

Moreover, the activity and effect of ztnfr11 on tumor progression and  
 30 metastasis can be measured *in vivo*. Several syngeneic mouse models have been developed to study the influence of polypeptides, compounds or other treatments on



tumor progression. In these models, tumor cells passaged in culture are implanted into mice of the same strain as the tumor donor. The cells will develop into tumors having similar characteristics in the recipient mice, and metastasis will also occur in some of the models. Tumor models include the Lewis lung carcinoma (ATCC No. CRL-1642) and

5 B16 melanoma (ATCC No. CRL-6323), amongst others. These are both commonly used tumor lines, syngeneic to the C57BL6 mouse, that are readily cultured and manipulated *in vitro*. Tumors resulting from implantation of either of these cell lines are capable of metastasis to the lung in C57BL6 mice. The Lewis lung carcinoma model has recently

10 been used in mice to identify an inhibitor of angiogenesis (O'Reilly MS, et al. Cell 79: 315-328,1994). C57BL6/J mice are treated with an experimental agent either through daily injection of recombinant protein, agonist or antagonist or a one time injection of recombinant adenovirus. Three days following this treatment,  $10^5$  to  $10^6$  cells are

15 implanted under the dorsal skin. Alternatively, the cells themselves may be infected with recombinant adenovirus, such as one expressing ztnfr11, before implantation so that the protein is synthesized at the tumor site or intracellularly, rather than systemically. The mice normally develop visible tumors within 5 days. The tumors are allowed to grow for a period of up to 3 weeks, during which time they may reach a size of 1500 - 1800 mm<sup>3</sup> in the control treated group. Tumor size and body weight are carefully monitored throughout the experiment. At the time of sacrifice, the tumor is removed and weighed

20 along with the lungs and the liver. The lung weight has been shown to correlate well with metastatic tumor burden. As an additional measure, lung surface metastases are counted. The resected tumor, lungs and liver are prepared for histopathological examination, immunohistochemistry, and *in situ* hybridization, using methods known in the art and described herein. The influence of the expressed polypeptide in question,

25 e.g., ztnfr11, on the ability of the tumor to recruit vasculature and undergo metastasis can thus be assessed. In addition, aside from using adenovirus, the implanted cells can be transiently transfected with ztnfr11. Moreover, purified ztnfr11 or ztnfr11-conditioned media can be directly injected in to this mouse model, and hence be used in this system. Use of stable ztnfr11 transfectants as well as use of inducible promoters to activate

30 ztnfr11 expression *in vivo* are known in the art and can be used in this system to assess ztnfr11 induction of metastasis. For general reference see, O'Reilly MS, et al. Cell

79:315-328, 1994; and Rusciano D, et al. Murine Models of Liver Metastasis. Invasion Metastasis 14:349-361, 1995.

5 Ztnfr11 polypeptides and antibodies may be used within diagnostic systems to detect the presence of its ligand polypeptides, such as a tumor necrosis factor ligand. The information derived from such detection methods would provide insight into the significance of ztnfr11 polypeptides in various diseases, and as a would serve as diagnostic tools for diseases for which altered levels of ztnfr11 are significant. Altered levels of ztnfr11 polypeptides may be indicative of pathological conditions including cancer, autoimmune disorders and infectious diseases.

10 In a basic assay, a single-stranded probe molecule is incubated with RNA, isolated from a biological sample, under conditions of temperature and ionic strength that promote base pairing between the probe and target ztnfr11 species. After separating unbound probe from hybridized molecules, the amount of hybrids is detected.

15 Well-established hybridization methods of RNA detection include northern analysis and dot/slot blot hybridization (see, for example, Ausubel *ibid.* and Wu et al. (eds.), "Analysis of Gene Expression at the RNA Level," in Methods in Gene Biotechnology, pages 225-239 (CRC Press, Inc. 1997)). Nucleic acid probes can be detectably labeled with radioisotopes such as <sup>32</sup>P or <sup>35</sup>S. Alternatively, ztnfr11 RNA can be detected with a nonradioactive hybridization method (see, for example, Isaac (ed.), 20 Protocols for Nucleic Acid Analysis by Nonradioactive Probes, Humana Press, Inc., 1993). Typically, nonradioactive detection is achieved by enzymatic conversion of chromogenic or chemiluminescent substrates. Illustrative nonradioactive moieties include biotin, fluorescein, and digoxigenin.

25 ztnfr11 oligonucleotide probes are also useful for *in vivo* diagnosis. As an illustration, <sup>18</sup>F-labeled oligonucleotides can be administered to a subject and visualized by positron emission tomography (Tavitian *et al.*, Nature Medicine 4:467, 1998).

Numerous diagnostic procedures take advantage of the polymerase chain reaction (PCR) to increase sensitivity of detection methods. Standard techniques for performing PCR are well-known (see, generally, Mathew (ed.), Protocols in Human 30 Molecular Genetics (Humana Press, Inc. 1991), White (ed.), PCR Protocols: Current Methods and Applications (Humana Press, Inc. 1993), Cotter (ed.), Molecular Diagnosis of Cancer (Humana Press, Inc. 1996), Hanausek and Walaszek (eds.), Tumor Marker Protocols (Humana Press, Inc. 1998), Lo (ed.), Clinical Applications of PCR (Humana Press, Inc. 1998), and Meltzer (ed.), PCR in Bioanalysis (Humana Press, Inc. 1998)).

PCR primers can be designed to amplify a sequence encoding a particular ztnfr11 domain or motif, such as the ztnfr11 cysteine rich pseudo repeat.

One variation of PCR for diagnostic assays is reverse transcriptase-PCR (RT-PCR). In the RT-PCR technique, RNA is isolated from a biological sample, reverse  
 5 transcribed to cDNA, and the cDNA is incubated with ztnfr11 primers (see, for example, Wu et al. (eds.), "Rapid Isolation of Specific cDNAs or Genes by PCR," in Methods in Gene Biotechnology, CRC Press, Inc., pages 15-28, 1997). PCR is then performed and the products are analyzed using standard techniques.

As an illustration, RNA is isolated from biological sample using, for  
 10 example, the guanidinium-thiocyanate cell lysis procedure described above. Alternatively, a solid-phase technique can be used to isolate mRNA from a cell lysate. A reverse transcription reaction can be primed with the isolated RNA using random oligonucleotides, short homopolymers of dT, or ztnfr11 anti-sense oligomers. Oligo-dT  
 15 primers offer the advantage that various mRNA nucleotide sequences are amplified that can provide control target sequences. Ztnfr11 sequences are amplified by the polymerase chain reaction using two flanking oligonucleotide primers that are typically at least 5 bases in length.

PCR amplification products can be detected using a variety of approaches. For example, PCR products can be fractionated by gel electrophoresis, and  
 20 visualized by ethidium bromide staining. Alternatively, fractionated PCR products can be transferred to a membrane, hybridized with a detectably-labeled ztnfr11 probe, and examined by autoradiography. Additional alternative approaches include the use of digoxigenin-labeled deoxyribonucleic acid triphosphates to provide chemiluminescence detection, and the C-TRAK colorimetric assay.

Another approach is real time quantitative PCR (Perkin-Elmer Cetus, Norwalk, Ct.). A fluorogenic probe, consisting of an oligonucleotide with both a reporter and a quencher dye attached, anneals specifically between the forward and reverse  
 25 primers. Using the 5' endonuclease activity of Taq DNA polymerase, the reporter dye is separated from the quencher dye and a sequence-specific signal is generated and increases as amplification increases. The fluorescence intensity can be continuously  
 30 monitored and quantified during the PCR reaction.

Another approach for detection of ztnfr11 expression is cycling probe technology (CPT), in which a single-stranded DNA target binds with an excess of DNA-  
 RNA-DNA chimeric probe to form a complex, the RNA portion is cleaved with RNase  
 35 H, and the presence of cleaved chimeric probe is detected (see, for example, Beggs et al., J. Clin. Microbiol. 34:2985, 1996 and Bekkaoui et al., Biotechniques 20:240, 1996).

Alternative methods for detection of ztnfr11 sequences can utilize approaches such as nucleic acid sequence-based amplification (NASBA), cooperative amplification of templates by cross-hybridization (CATCH), and the ligase chain reaction (LCR) (see, for example, Marshall et al., U.S. Patent No. 5,686,272 (1997), Dyer et al., J. Virol. Methods 5 60:161, 1996; Ehricht et al., Eur. J. Biochem. 243:358, 1997 and Chadwick et al., J. Virol. Methods 70:59, 1998). Other standard methods are known to those of skill in the art.

Ztnfr11 probes and primers can also be used to detect and to localize ztnfr11 gene expression in tissue samples. Methods for such *in situ* hybridization are 10 well-known to those of skill in the art (see, for example, Choo (ed.), In Situ Hybridization Protocols, Humana Press, Inc., 1994; Wu et al. (eds.), "Analysis of Cellular DNA or Abundance of mRNA by Radioactive *In Situ* Hybridization (RISH)," in Methods in Gene Biotechnology, CRC Press, Inc., pages 259-278, 1997 and Wu et al. (eds.), "Localization of DNA or Abundance of mRNA by Fluorescence *In Situ* Hybridization (RISH)," in 15 Methods in Gene Biotechnology, CRC Press, Inc., pages 279-289, 1997).

Various additional diagnostic approaches are well-known to those of skill in the art (see, for example, Mathew (ed.), Protocols in Human Molecular Genetics Humana Press, Inc., 1991; Coleman and Tsongalis, Molecular Diagnostics, Humana Press, Inc., 1996 and Elles, Molecular Diagnosis of Genetic Diseases, Humana Press, 20 Inc., 1996).

In addition, such polynucleotide probes could be used to hybridize to counterpart sequences on individual chromosomes. Chromosomal identification and/or mapping of the ztnfr11 gene could provide useful information about gene function and disease association. Many mapping techniques are available to one skilled in the art, for 25 example, mapping somatic cell hybrids, and fluorescence *in situ* hybridization (FISH). A preferred method is radiation hybrid mapping. Radiation hybrid mapping is a somatic cell genetic technique developed for constructing high-resolution, contiguous maps of mammalian chromosomes (Cox et al., Science 250:245-50, 1990). Partial or full knowledge of a gene's sequence allows the designing of PCR primers suitable for use 30 with chromosomal radiation hybrid mapping panels. Commercially available radiation hybrid mapping panels which cover the entire human genome, such as the Stanford G3 RH Panel and the GeneBridge 4 RH Panel (Research Genetics, Inc., Huntsville, AL), are

available. These panels enable rapid, PCR based, chromosomal localizations and ordering of genes, sequence-tagged sites (STSs), and other non-polymorphic- and polymorphic markers within a region of interest. This includes establishing directly proportional physical distances between newly discovered genes of interest and previously mapped markers. The precise knowledge of a gene's position can be useful in a number of ways including: 1) determining if a sequence is part of an existing contig and obtaining additional surrounding genetic sequences in various forms such as YAC-, BAC- or cDNA clones, 2) providing a possible candidate gene for an inheritable disease which shows linkage to the same chromosomal region, and 3) for cross-referencing model organisms such as mouse which may be beneficial in helping to determine what function a particular gene might have.

The UMLR polynucleotides of SEQ ID NO:2 have been mapped to chromosome Xq11-q12. Thus, the present invention also provides reagents which will find use in diagnostic applications. For example, the UMLR gene, a probe comprising UMLR DNA or RNA or a subsequence thereof can be used to determine if the UMLR gene is present on chromosome Xq11-q12 or if a mutation has occurred. Detectable chromosomal aberrations at the UMLR gene locus include, but are not limited to, aneuploidy, gene copy number changes, insertions, deletions, restriction site changes and rearrangements. These aberrations can occur within the coding sequence, within introns, or within flanking sequences, including upstream promoter and regulatory regions, and may be manifested as physical alterations within a coding sequence or changes in gene expression level.

Such aberrations can be detected using polynucleotides of the present invention by employing molecular genetic techniques, such as restriction fragment length polymorphism (RFLP) analysis, short tandem repeat (STR) analysis employing PCR techniques, and other genetic linkage analysis techniques known in the art (Sambrook et al., *ibid.*; Ausubel et. al., *ibid.*; Marian, *Chest* 108:255-65, 1995).

In general, these diagnostic methods comprise the steps of (a) obtaining a genetic sample from a patient; (b) incubating the genetic sample with a polynucleotide probe or primer as disclosed above, under conditions wherein the polynucleotide will

hybridize to complementary polynucleotide sequence, to produce a first reaction product; and (iii) comparing the first reaction product to a control reaction product. A difference between the first reaction product and the control reaction product is indicative of a genetic abnormality in the patient. Genetic samples for use within the present invention

5 include genomic DNA, cDNA, and RNA. The polynucleotide probe or primer can be RNA or DNA, and will comprise a portion of SEQ ID NOs:1 or 3, the complement of SEQ ID NOs:1 or 3, or an RNA equivalent thereof. Suitable assay methods in this regard include molecular genetic techniques known to those in the art, such as restriction

10 fragment length polymorphism (RFLP) analysis, short tandem repeat (STR) analysis employing PCR techniques, ligation chain reaction (Barany, PCR Methods and Applications 1:5-16, 1991), ribonuclease protection assays, and other genetic linkage analysis techniques known in the art (Sambrook et al., ibid.; Ausubel et. al., ibid.; Marian, Chest 108:255-65, 1995). Ribonuclease protection assays (see, e.g., Ausubel et al., ibid., ch. 4) comprise the hybridization of an RNA probe to a patient RNA sample,

15 after which the reaction product (RNA-RNA hybrid) is exposed to RNase. Hybridized regions of the RNA are protected from digestion. Within PCR assays, a patient's genetic sample is incubated with a pair of polynucleotide primers, and the region between the primers is amplified and recovered. Changes in size or amount of recovered product are indicative of mutations in the patient. Another PCR-based technique that can be

20 employed is single strand conformational polymorphism (SSCP) analysis (Hayashi, PCR Methods and Applications 1:34-8, 1991).

Antisense methodology can be used to inhibit ztnfr11 gene transcription, such as to inhibit T cell development and interaction with other cells. Polynucleotides that are complementary to a segment of a ztnfr11-encoding polynucleotide (e.g., a

25 polynucleotide as set forth in SEQ ID NOs:2, 27, or 38) are designed to bind to ztnfr11-encoding mRNA and to inhibit translation of such mRNA. Such antisense polynucleotides are used to inhibit expression of ztnfr11 polypeptide-encoding genes in cell culture or in a subject.

Additionally, polynucleotides of the present invention can be used as a

30 marker for the X chromosome. Diseases which map to the Xq11-12 include, but are not limited to, adenocarcinoma of the prostate (Verma, R. S. et al., Cancer Invest. 17(6):441-

447, 1999); and dystonia-parkinsonism, sideroblastic anemia, and several forms of mental retardation (Villard, L. et al., Gene 235(1-2):43-50, 1999). Other aberrations at this locus include a deletion which is related to borderline and invasive epithelial tumors (Edelson, M.L. et al., Oncogene 16(2):192-202, 1998), and a gain in patients with adult  
 5 testicular germ cell tumors (Summersgill, B. et al., Br J Cancer 77(2):305-13, 1998).

The polypeptides of UMLR may represent an antigenic marker for melanoma, osteosarcoma, breast carcinoma, and lymphoma, as well as tumors of the lung and uterus. Thus, these polypeptides, or fragments thereof may be useful as antigens to produce humanized antibodies for treatment of these specific tumors.  
 10 Additionally, these polypeptides and polypeptide fragments can be useful to generate vaccines for use cancer therapy.

For pharmaceutical use, the proteins of the present invention can be administered intravaginally, orally, rectally, parenterally (particularly intravenous or subcutaneous), intracisternally, intraperitoneally, topically (as douches, powders,  
 15 ointments, drops or transdermal patch) buccally, or as a pulmonary or nasal inhalant. Intravenous administration will be by bolus injection or infusion over a typical period of one to several hours. In general, pharmaceutical formulations will include a UMLR protein, alone, or in conjunction with a dimeric partner, in combination with a pharmaceutically acceptable vehicle, such as saline, buffered saline, 5% dextrose in  
 20 water or the like. Formulations may further include one or more excipients, preservatives, solubilizers, buffering agents, albumin to prevent protein loss on vial surfaces, etc. Methods of formulation are well known in the art and are disclosed, for example, in Remington: The Science and Practice of Pharmacy, Gennaro, ed., Mack Publishing Co., Easton, PA, 19th ed., 1995. Therapeutic doses will generally be in the  
 25 range of 0.1 to 100 µg/kg of patient weight per day, preferably 0.5-20 mg/kg per day, with the exact dose determined by the clinician according to accepted standards, taking into account the nature and severity of the condition to be treated, patient traits, etc. Determination of dose is within the level of ordinary skill in the art. The proteins may be administered for acute treatment, over one week or less, often over a period of one to  
 30 three days or may be used in chronic treatment, over several months or years. In general, a therapeutically effective amount of UMLR is an amount sufficient to produce a

clinically significant change in, tumor suppression, apoptosis, myogenesis, inflammation, and infection in tissues of the uterus, lung carcinoma and breast carcinoma, as well as in cells of melanoma, osteosarcoma, and lymphoma. Similarly, a therapeutically effective amount of UMLR is an amount sufficient to produce a clinically significant change in disorders associated with tissues of the uterus, melanoma, osteosarcoma, breast carcinoma, and lymphoma, and tumors of the lung.

The invention is further illustrated by the following non-limiting examples.

10

## EXAMPLES

### Example 1

#### Extension of EST Sequence

The novel UMLR polypeptide-encoding polynucleotides of the present invention were initially identified by querying a database of partial sequences. A partial sequence was identified in a uterus, myometrium, leiomyoma cDNA library. The polynucleotide sequence (SEQ ID NO:1) of the insert corresponding to the cDNA clone was sequenced. The deduced amino acid sequence of the insert was determined to be full-length and is shown in SEQ ID NO:2. This polypeptide, and the polynucleotides encoding it, were identified as a novel member of the tumor necrosis factor receptor family, UMLR.

20

### Example 2

#### Tissue Distribution

Human Multiple Tissue Northern Blots (MTN I, MTN II and MTN III; Human Cancer Cell Line; Human Immune Blot; Clontech, In-house human lymphocyte subset blot; Clontech, Palo Alto, CA) were probed to determine the tissue distribution of human UMLR expression. An approximately 555 bp PCR derived probe was amplified as a template and oligonucleotide ZC25364 (SEQ ID NO:12) and oligonucleotide ZC25365 (SEQ ID NO:13) as primers. The amplification was carried out as follows: 1 cycle at 94°C for 1.5 minutes, 35 cycles of 94°C for 15 seconds and 60°C for 30 seconds, followed by one cycle at 72°C for 10 minutes. The PCR products were visualized by agarose gel electrophoresis and the 555 bp PCR product was purified using

30



a Gel Extraction Kit (Qiagen, Chatsworth, CA) according to manufacturer's instructions. The probe was radioactively labeled using the MULTIPRIME DNA labeling kit (Amersham, Arlington Heights, IL) according to the manufacturer's instructions. The probe was purified using a NUCTRAP push column (Stratagene). EXPRESSHYB (Clontech) solution was used for prehybridization and as a hybridizing solution for the Northern blots. Hybridization took place overnight at 65°C using  $1 \times 10^6$  cpm/ml of labeled probe. The blots were then washed four times in 2X SSC and 0.1% SDS at RT, followed by 2 washes in 0.1X SSC and 0.1% SDS at 55°C. A major transcript of approximately 3.5 kb and a minor transcript of ~4.7kb were seen in the lung carcinoma cell line A-549 and the melanoma cell line G-361. All other samples were negative at a 24hr exposure time.

### Example 3

#### Chromosomal Assignment and Placement of UMLR

UMLR was mapped to human chromosome X using the commercially available version of the "Stanford G3 Radiation Hybrid Mapping Panel" (Research Genetics, Inc., Huntsville, AL). The "Stanford G3 RH Panel" contains PCRable DNAs from each of 83 radiation hybrid clones of the whole human genome, plus two control DNAs (the RM donor and the A3 recipient). A publicly available WWW server (<http://shgc-www.stanford.edu>) allows chromosomal localization of markers.

For the mapping of UMLR with the "Stanford G3 RH Panel", 20  $\mu$ l reactions were set up in a 96-well microtiter plate (Stratagene, La Jolla, CA) and used in a "RoboCycler Gradient 96" thermal cycler (Stratagene). Each of the 85 PCR reactions consisted of 2  $\mu$ l 10X KlenTaq PCR reaction buffer (CLONTECH Laboratories, Inc., Palo Alto, CA), 1.6  $\mu$ l dNTPs mix (2.5 mM each, PERKIN-ELMER, Foster City, CA), 1  $\mu$ l sense primer, ZC 25352 (SEQ ID NO:14), 1  $\mu$ l antisense primer, ZC 25353 (SEQ ID NO:15), 2  $\mu$ l "RediLoad" (Research Genetics, Inc., Huntsville, AL), 0.4  $\mu$ l 50X Advantage KlenTaq Polymerase Mix (Clontech Laboratories, Inc.), 25 ng of DNA from an individual hybrid clone or control and x  $\mu$ l ddH<sub>2</sub>O for a total volume of 20  $\mu$ l. The reactions were overlaid with an equal amount of mineral oil and sealed. The PCR cycler conditions were as follows: an initial 1 cycle 5 minute denaturation at 94°C, 35 cycles of

a 45 seconds denaturation at 94°C, 45 seconds annealing at 64°C and 1 minute AND 15 seconds extension at 72°C, followed by a final 1 cycle extension of 7 minutes at 72°C. The reactions were separated by electrophoresis on a 2% agarose gel.

The results showed linkage of UMLR to the human chromosome X framework marker SHGC-1486 with a LOD score of 7.00 and at a distance of 22 cR\_10000 from the marker. The use of surrounding genes/markers positions UMLR in the Xq11-q12 chromosomal region.

#### Example 4

##### Construction of Mammalian Soluble ztnfr11 Expression Vectors: ztnfr11sR/CEE and ztnfr11sR/Fc4

An expression vector was prepared to express the soluble ztnfr11 polypeptide (ztnfr11sR, i.e., from residue 2 to residue 132 of SEQ ID NO:2) fused to a C-terminal Glu-Glu tag (SEQ ID NO:16).

A PCR generated ztnfr11 DNA fragment of about 400bp was created using oligo ZC25598 (SEQ ID NO:17) and oligo ZC25596 (SEQ ID NO:18) as PCR primers to add BamHI and BglII restriction sites at 5' and 3' ends of the soluble ztnfr11 DNA, respectively. A plasmid containing the ztnfr11 cDNA (SEQ ID NO:1) was used as a template. PCR amplification of the ztnfr11 fragment was performed as follows: One cycle at 94°C for 1 minute; 25 cycles at 94°C for 30 seconds, 68°C for 90 seconds, followed by an additional 68°C incubation for 4 minutes, and hold at 10°C. The reaction was purified by chloroform/phenol extraction and isopropanol precipitation, and digested with BamHI and Bgl2 (Boehringer Mannheim, Indianapolis, IN). A band of approximately 400bp, as visualized by 1% agarose gel electrophoresis, was excised and the DNA was purified using a QiaexII™ purification kit (Qiagen, Valencia, CA) according to the manufacturer's instruction.

The pZP9CEE plasmid is a mammalian expression vector containing an expression cassette having the mouse metallothionein-1 promoter, human tPA leader peptide, multiple restriction sites for insertion of coding sequences, a Glu-Glu tag, and a human growth hormone terminator. The plasmid also has an *E. coli* origin of replication, a mammalian selectable marker expression unit having an SV40 promoter, an enhancer

and an origin of replication, as well as a DHFR gene, and the SV40 terminator. The ztnfr11sR/pZP9 /CEE expression vector uses the human tPA leader peptide (SEQ ID NO:19) and attaches the Glu-Glu tag to the C-terminus of the extracellular portion of the ztnfr11 polypeptide sequence.

5 About 30ng of the restriction digested ztnfr11sR insert and about 10ng of the digested vector were ligated at room temperature for 2 hours. One microliter of ligation reaction was electroporated into DH10B competent cells (Gibco BRL, Rockville, MD) according to manufacturer's direction and plated onto LB plates containing 50mg/ml ampicillin, and incubated overnight. Colonies were screened by  
10 restriction analysis of DNA, which was prepared from 2 ml liquid cultures of individual colonies. The insert sequence of positive clones was verified by sequence analysis. The sequence of the soluble ztnfr11 gene fused amino-terminally to the human tPA leader and carboxy-terminally to the CEE tag is shown in SEQ ID NO:21. Thus the excised ztnfr11sR DNA was subcloned into the pZP9CEE plasmid, which had been cut with  
15 BamHI. The A large-scale plasmid preparation was done using a Qiagen® Mega prep kit (Qiagen) according to manufacturer's instruction.

The same process was used to prepare the ztnfr11 soluble receptor with a C-terminal Fc4 tag (SEQ ID NO:20), creating the ztnfr11sR/Fc4. To prepare ztnfr11sR/Fc4, the pZP9 vector has a Fc4 tag in place of the Glu-Glu tag. The sequence  
20 of the soluble ztnfr11 gene fused amino-terminally to the human tPA leader and carboxy-terminally to the Fc4 tag is shown in SEQ ID NO:22. Fc4 is the Fc region derived from human IgG, which contains a mutation so that it no longer binds the Fc receptor.

### Example 5

#### 25 Transfection and Expression of ztnfr11 Soluble Receptor Polypeptides

The day before the transfection, BHK 570 cells (ATCC No. CRL-10314; ATCC, Manassas, VA) were plated in a 10-cm plate with 50% confluence in normal BHK DMEM (Gibco/BRL High Glucose) media. The day of the transfection, the cells were washed once with Serum Free (SF) DMEM, followed by transfection with the  
30 ztnfr11sR/Fc4 or ztnfr11sR/CEE expression plasmids. Sixteen micrograms of each DNA construct were separately diluted into a total final volume of 640µl SF DMEM. A

diluted LipofectAMINE™ mixture (35µl LipofectAMINE™ in 605µl SF media) was added to the DNA mix, and incubated for 30 minutes at room temperature. Five milliliters of SF media was added to the DNA/LipofectAMINE™ mixture, which was then added to BHK cells. The cells were incubated at 37°C/5% CO<sub>2</sub> for 5 hours, after which 6.4ml of BHK media with 10% FBS was added. The cells were incubated overnight at 37°C/5% CO<sub>2</sub>.

Approximately 24 hours post-transfection, the BHK cells were split into selection media with 1µM methotrexate (MTX). The cells were repeatedly split in this manner until stable ztnfr11sR/CEE and ztnfr11sR/Fc4 cell lines were identified. To detect the expression level of the ztnfr11 soluble receptor fusion proteins, the BHK cells were washed with PBS and incubated in SF media for 72 hours. The SF condition media was collected and 20 µl of the sample was run on 10% SDS-PAGE gel under reduced conditions. The protein bands were transferred to nitrocellulose filter by Western blot, and the fusion proteins were detected using either goat-anti-human IgG/HRP conjugates for the ztnfr11sR/Fc4 fusion or mouse-anti-Glu-Glu tag/HRP conjugates for the ztnfr11sR/CEE fusion. Expression vectors containing a different soluble receptor fused to the Fc4 or the CEE tags were used as controls. The expression level is approximately 10mg/L for ztnfr11sR/Fc4 and 0.5mg/L for ztnfr11sR/CEE.

Transfected BHK cells were transferred into T-162 flasks. Once the BHK cells reached about 80% confluence, they were washed with PBS and incubated in 100ml SF media for 72 hours, and then the condition media was collected for protein purification.

### Example 6

#### Purification and Analysis of ztnfr11sR/CEE

Recombinant carboxyl terminal Glu-Glu tagged ztnfr11sR was produced from transfected BHK cells as described in Example 5 above. About six liters of conditioned media were harvested from 60 dishes after roughly 72 hours incubation. A portion of the media was sterile filtered using filtration units from different manufactures. The Nalgene 0.2µm and 0.45µm filters, and Millipore Express 0.22µm filter were compared. The amount of ztnfr11sR/CEE lost through the filter can be more

than 50% of the starting material with Nalgene's 0.2 $\mu$ m filter. The other filters performed better and gave more reasonable recovery of the protein. Additionally, the Millipore Express 0.22 $\mu$ m filter achieved a good flow rate. The level of protein expression reached the optimal concentration after about 72 hours in new media. Three  
 5 harvests of the ztnfr11sR/CEE conditioned media were collected.

Protein was purified from the filtered media by a combination of Anti-Glu-Glu (Anti-EE) peptide antibody affinity chromatography and S-100 gel exclusion chromatography. Culture medium was directly loaded onto a 20x185mm (58-ml bed volume) anti-EE antibody affinity column at a flow of about 4 ml/minute. Following  
 10 column washing with ten column volumes of PBS, bound protein was eluted with two column volumes of 0.4mg/ml EYMPTD peptide (Princeton Biomolecules, NJ). Fractions of 5 ml were collected. Samples from the anti-EE antibody affinity column were analyzed by SDS-PAGE with silver staining and western blotting for the presence of ztnfr11sR/CEE. Fractions containing the ztnfr11sR/CEE protein were pooled and  
 15 concentrated to 4 mls using Biomax-5 concentrator (Millipore), and loaded onto a 16 x 1000 mm Sephacryl S-100 HR gel filtration column (Amersham Pharmacia Biotech). The fractions containing purified ztnfr11sR/CEE were pooled, filtered through 0.2  $\mu$ m filter, aliquoted into 100  $\mu$ l each, and frozen at -80°C. The concentration of the final purified protein was determined by BCA assay (Pierce) and HPLC-amino acid analysis.

20 Recombinant ztnfr11sR/CEE was analyzed by SDS-PAGE (Nupage 4-12%), Novex) with either coomassie and silver staining method (Fast Silver, Geno Tech), and Western blotting using monoclonal anti-EE antibody. Either the conditioned media or purified protein was electrophoresed using a Novex's Xcell II mini-cell (San Diego, CA) and transferred to nitrocellulose (0.2  $\mu$ m; Bio-Rad Laboratories, Hercules,  
 25 CA) at room temperature using Novex's Xcell II blot module with stirring according to directions provided in the instrument manual. The transfer was run at 500 mA for one hour in a buffer containing 25 mM Tris base, 200 mM glycine, and 20% methanol. The filters were then blocked with 10% non-fat dry milk in PBS for 10 minutes at room temperature. The nitrocellulose was quickly rinsed, then primary antibody was added in  
 30 PBS containing 2.5% non-fat dry milk. The blots were incubated for two hours at room temperature or overnight at 4°C with gentle shaking. Following the incubation, blots

were washed three times for 10 minutes each in PBS. Secondary antibody (goat anti-mouse IgG conjugated to horseradish peroxidase; obtained from Rockland Inc., Gilbertsville, PA) diluted 1:2000 in PBS containing 2.5% non-fat dry milk was added, and the blots were incubated for two hours at room temperature with gentle shaking.

5 The blots were then washed three times, 10 minutes each, in PBS, then quickly rinsed in H<sub>2</sub>O. The blots were developed using commercially available chemiluminescent substrate reagents (SuperSignal® ULTRA reagents 1 and 2 mixed 1:1; reagents obtained from Pierce Chemical Co.), and the signal was captured using Lumi-Imager's Lumi Analyst 3.0 software (Boehringer Mannheim GmbH, Germany) for exposure times

10 ranging from 10 second to 5 minutes or as necessary.

The purified ztnfr11sR/CEE appeared as a single band with either the coomassie or silver staining at about 18 kDa under non-reducing conditions, and at about 20 kDa under reducing conditions, suggesting the monomeric form of ztnfr11sR/CEE under non-reducing conditions.

15

### Example 7

#### Purification and Analysis of ztnfr11sR/Fc4

Recombinant carboxyl terminal Fc4 tagged ztnfr11sR was produced from transfected BHK cells as described in Example 5 above. Approximately five-liters of

20 conditioned media were harvested from 60 dishes after about 72 hours of incubation. A portion of the media was sterile filtered using filtration units from different manufactures. The Nalgene 0.2µm and 0.45µm filters, Millipore Express 0.22µm filter, and Durapore 0.45µm filter were compared. The Millipore Express 0.22µm filter gave the best yield and flow rate. The level of protein expression reached the optimal

25 concentration after about 72 hours in the new media. Normally three to four harvests of the media were collected.

Protein was purified from the filtered media by a combination of Poros 50 protein A affinity chromatography (PerSeptive Biosystems, 1-5559-01, Framingham, MA) and S-200 gel exclusion chromatography column (Amersham Pharmacia Biotech).

30 Culture medium was directly loaded onto a 10x80mm (6.2-ml bed volume) protein A affinity column at a flow of about 4 ml/minute. Following column washing for ten

column volumes of PBS, bound protein was eluted by five column volumes of 0.1 M glycine, pH 3.0 at 10 ml/minute). Fractions of 1.5 ml each were collected into tubes containing 38µl of 2.0 M Tris, pH 8.8, in order to neutralize the eluted proteins. Samples from the affinity column were analyzed by SDS-PAGE with Coomassie staining and Western blotting for the presence of ztnfr11sR/Fc4 using human IgG-HRP. Ztnfr11sR/Fc4-containing fractions were pooled and concentrated to 4 mls using Biomax-30 concentrator (Millipore), and loaded onto a 16 x1000 mm Sephacryl S-200 HR gel filtration. The fractions containing purified ztnfr11sR/Fc4 were pooled, filtered through 0.2 µm filter, aliquoted into 100, 200 and 500 µl each, and frozen at -80°C. The concentration of the final purified protein was determined by BCA assay (Pierce) and HPLC-amino acid analysis.

Recombinant ztnfr11sR/Fc4 was analyzed by SDS-PAGE (Nupage 4-12%, Novex) with coomassie staining method and Western blotting using human IgG-HRP. Either the conditioned media or purified protein was electrophoresed using a Novex's Xcell II mini-cell (San Diego, CA) and transferred to nitrocellulose (0.2 µm; Bio-Rad Laboratories, Hercules, CA) at room temperature using Novex's Xcell II blot module with stirring according to directions provided in the instrument manual. The transfer was run at 500 mA for one hour in a buffer containing 25 mM Tris base, 200 mM glycine, and 20% methanol. The filters were then blocked with 10% non-fat dry milk in PBS for 10 minutes at room temperature. The nitrocellulose was quickly rinsed, then the human Ig-HRP antibody (1:2000) was added in PBS containing 2.5% non-fat dry milk. The blots were incubated for two hours at room temperature, or overnight at 4°C, with gentle shaking. Following the incubation, the blots were washed three times for 10 minutes each in PBS, then quickly rinsed in H<sub>2</sub>O. The blots were developed using commercially available chemiluminescent substrate reagents (SuperSignal® ULTRA reagents 1 and 2 mixed 1:1; reagents obtained from Pierce Chemical Co.), and the signal was captured using Lumi-Imager's Lumi Analyst 3.0 software (Boehringer Mannheim GmbH, Germany) for exposure times ranging from 10 second to 5 minutes or as necessary.

The purified ztnfr11sR/Fc4 appeared as a single band with either the coomassie or silver staining at about 100 kDa under non-reducing conditions, and at

about 55 kDa under reducing conditions, suggesting the dimeric form of ztnfr11sR/Fc4 under non-reducing conditions as expected.

### Example 8

#### 5                    Identification of Cells Expressing ztnfr11 Using *In Situ* Hybridization

Specific human tissues were isolated and screened for ztnfr11 expression by *in situ* hybridization. Various human tissues prepared, sectioned and subjected to *in situ* hybridization included normal lung, lung carcinoma and breast carcinoma. The tissues were fixed in 10% buffered formalin and blocked in paraffin using standard techniques. Tissues were sectioned at 4 to 8 microns. Tissues were prepared using a standard protocol ("Development of non-isotopic *in situ* hybridization" at <http://dir.niehs.nih.gov/dirlep/ish.html>). Briefly, tissue sections were deparaffinized with HistoClear (National Diagnostics, Atlanta, GA) and then dehydrated with ethanol. Next they were digested with Proteinase K (50 µg/ml) (Boehringer Diagnostics, Indianapolis, IN) at 37°C for 2 to 20 minutes. This step was followed by acetylation and re-hydration of the tissues.

Two *in situ* probes generated by PCR were designed against the human ztnfr11 sequence. Two sets of oligos were designed to generate probes for separate regions of the ztnfr11 cDNA: (1) Oligos ZC26463 (SEQ ID NO:23) and ZC26464 (SEQ ID NO:24) were used to generate a 439 bp probe for ztnfr11 (from nucleotide 290 to nucleotide 840 as shown in SEQ ID NO:1); and (2) ZC26470 (SEQ ID NO: 25) and ZC26471 (SEQ ID NO: 26) were used to generate a 548 bp probe for ztnfr11 (from nucleotide 560 to nucleotide 1000 as shown in SEQ ID NO:1). The antisense oligo from each set also contained the working sequence for the T7 RNA polymerase promoter to allow for easy transcription of antisense RNA probes from these PCR products. The PCR reaction conditions were as follows: 2 cycles at 95°C for 30 sec, 50°C for 1 min, 72°C for 1 min followed by 33 cycles of 95°C for 30 sec, 72°C for 2 min. Probes were subsequently labeled with digoxigenin (Boehringer) or biotin (Boehringer) using an *In Vitro* transcription System (Promega, Madison, WI) as per manufacturer's instruction.

30                    *In situ* hybridization was performed with a digoxigenin- or biotin-labeled ztnfr11 probe (above). The probe was added to the slides at a concentration of 1 to 5



pmol/ml for 12 to 16 hours at 60°C. Slides were subsequently washed in 2XSSC and 0.1XSSC at 55°C. The signals were amplified using tyramide signal amplification (TSA) (TSA, *in situ* indirect kit; NEN) and visualized with Vector Red substrate kit (Vector Lab) as per manufacturer's instructions. The slides were then counter-stained with hematoxylin (Vector Laboratories, Burlingame, CA).

Positive signals were seen in the human lung carcinoma and one of two breast carcinoma tissue samples. The positive-staining cells in the human lung carcinoma sample appeared to be plasma cells, mononuclear cells of unknown type and occasional alveolar epithelial cells. In one of the breast carcinoma tissue samples the positive staining cells appeared to be epithelial/myoepithelial cells, clusters of tumor cells within vascular lumens, mononuclear cells and occasional vascular endothelial cells. The normal human lung tissue sample and one of the breast carcinoma tissue samples were negative for ztnfr11.

### Example 9

#### Human ztnfr11 Polyclonal Antibodies

Polyclonal antibodies were prepared by immunizing 2 female New Zealand white rabbits with the purified recombinant protein ztnfr11-CEE protein expressed in BHK from Example 6. The rabbits were each given an initial intraperitoneal (ip) injection of 200 µg of purified protein in Complete Freund's Adjuvant followed by booster ip injections of 100 µg peptide in Incomplete Freund's Adjuvant every three weeks. Seven to ten days after the administration of the second booster injection (3 total injections), the animals were bled and the serum was collected. The animals were then boosted and bled every three weeks.

The ztnfr11sR-specific polyclonal antibodies were affinity purified from the rabbit serum using a CNBr-SEPHAROSE 4B protein column (Pharmacia LKB) that was prepared using 10 mg of purified recombinant huztnfr11-Fc protein (as prepared in Example 6) per gram of CNBr-SEPHAROSE, followed by 20X dialysis in PBS overnight. Ztnfr11sR-specific antibodies were characterized by ELISA using 1 µg/ml of the specific purified recombinant huztnfr11-CEE-BHK protein as antibody target. The

lower limit of detection (LLD) of the rabbit anti-ztnfr11sR affinity purified antibody was 100 pg/ml on its specific purified recombinant protein antigen ztnfr11sR/CEE.

#### Example 10

##### Human ztnfr11 Skin Northern

5 A probe of approximately 555 bp was made by PCR using plasmid DNA containing SEQ ID NO:1 as a template and oligonucleotide ZC25364 (SEQ ID NO:12 ) and ZC25365 (SEQ ID NO:13) as primers. The amplification was carried out as follows: 1 cycle at 94°C for 1 minutes, 35 cycles of 94°C for 30 seconds, 55°C for 30 seconds and 72°C for 30 seconds, followed by 1 cycle at 72°C for 10 minutes. The PCR  
10 products were visualized by agarose gel electrophoresis and the 555 bp probe was purified using a Gel Extraction Kit (Qiagen, Chatsworth, CA) according to manufacturer's instructions. The probe was radioactively labeled using the REDIPRIME DNA labeling kit (Amersham, Arlington Heights, IL) according to the manufacturer's instructions. The probe was purified using a NUCTRAP push column (Stratagene).

15 An in-house blot containing poly A+ RNA from human normal liver, human normal pancreas and the human melanoma cell lines including C32, Malme 3M, SK-MEL-2, and WM-115 was prehybridized in EXPRESSHYB (Clontech) for 3 hours at 65°C. Hybridization took place overnight at 65°C using 10<sup>6</sup> cpm/ml of labeled probe. The blots were then washed four times in 2X SSC and 0.1% SDS at room temp,  
20 followed by 2 washes in 0.1X SSC and 0.1% SDS at 50°C. A transcript of approximately 3.5 kb was observed most strongly in the Malme 3M cell line and weakly in the C32 and SK-MEL-2 cell lines. Liver, and pancreas tissues and the WM-115 cell line were negative at the 18 hour exposure time.

#### Example 11

##### Human ztnfr11 Tumor Polymerase Chain Reaction

25 A nested 5' RACE reaction was performed using marathon cDNAs from a lymphoma cell line (Burkitt's Lymphoma, DAUDI) and three osteosarcoma cell lines: MG63, HOS, and SaOS2. A 50 ul PCR reaction was run under the following conditions: 20 pmol of a sense primer corresponding to the vector sequence and 20 pmol if an  
30 antisense primer corresponding to the polynucleotide sequence as shown in SEQ ID NO:1 from nucleotide 664 to nucleotide 688 were used in a 3-step, 2-temperature

reaction including an initial one minute at 94°C followed by 5 cycles of 94°C for 30 seconds and 72°C for 5 minutes, then 5 cycles of 94°C for 30 seconds and 70°C for 5 minutes, then 25 cycles of 94°C for 30 seconds and 68°C for 5 minutes. A final extension of 72°C for 10 min completed the reaction. One ul of a 1:50 dilution of the  
 5 above reactions was used for nested 5' RACE using 20 pmol of an nested sense primer corresponding to the vector sequence and 20 pmol of an antisense primer corresponding to the polynucleotide sequence as shown in SEQ ID NO:1 from nucleotide 248 to nucleotide 272. A 50 µl reaction was set up and run under the following conditions. An  
 10 initial one minute at 94°C was followed by 4 cycles of 94°C for 30 seconds and 72°C for 5 minutes, then 4 cycles of 94°C for 30 seconds and 70°C for 5 minutes, then 20 cycles of 94°C for 30 seconds and 68°C for 5 minutes. The DAUDI, MG63, and SaOS2, were positive for a 250 bp band. The SaOS2 cell line also contained a 350 bp band. The HOS cell line contained a band of about 900 bp. All bands were sequenced and are isoforms of the ztnfr11 sequence.

15

### Example 12

#### Additional Human ztnfr11 Tumor Polymerase Chain Reaction

A PCR reaction was set up using marathon cDNAs as follows: 30 pmol of a sense primer (corresponding to the polynucleotide as shown in SEQ ID NO:1 from  
 20 nucleotide 441 to nucleotide 464) and an anti sense primer (corresponding to the polynucleotide as shown in SEQ ID NO:1 from nucleotide 856 to 879) were used in a 50 ul reaction in which an initial step of 94°C for 1 minute was run followed by 3 cycles of 94°C for 30 seconds, 63°C for 30 seconds and 72°C for 30 seconds then 32 cycles of 94°C for 30 seconds, 61°C for 30 seconds and 72°C for 30 seconds with a final  
 25 minute extension at 72°C. Two bands were seen at ~530 bp and ~430 bp in MG63, DAUDI, U2OS, cell lines and thymus and pituitary tissues. These PCR bands were excised from the U2OS lane and submitted to sequencing. The upper band was found to have a ~82 bp insertion relative to the corresponding region of the polynucleotide as shown in SEQ ID NO:1. The polynucleotide sequence of the upper PCR band can be  
 30 seen in SEQ ID NO:31. A polypeptide sequence corresponding to the variant polynucleotide sequence which contains the ~82 bp insertion can be seen in SEQ ID

NO:27. A degenerate polynucleotide sequence for this variant is shown in SEQ ID NO:28. The polynucleotide sequence of this lower PCR band can be seen in SEQ ID NO:32. The lower PCR band was found to have the same insertion and an additional ~87 bp deletion relative to the corresponding region of the polynucleotide as shown in  
 5 SEQ ID NO:1. A polypeptide sequence corresponding to this variant polynucleotide sequence which contains the ~82 bp insertion and the ~87 bp deletion can be seen in SEQ ID NO:29. A degenerate polynucleotide sequence for this variant is shown in SEQ ID NO:30.

These PCR bands were excised, purified using a Qiagen gel extraction kit  
 10 and subcloned into the pCR 2.1 TOPO vector (Invitrogen, San Diego, CA). The resulting nucleotide sequence of the clone containing the 552 bp band (upper) is shown in SEQ ID NO: 33, which corresponds to the 175 residue amino acid sequence as shown in SEQ ID NO:34. A composite of the polypeptide sequence of SEQ ID NO:2 with SEQ ID NO:34 results in a 299 residue protein shown in SEQ ID NO:35. This variant of  
 15 ztnfr11 differs from the polypeptide sequence as shown in SEQ ID NO:2 by a two amino acid insertion (Val-Ala) at residue 172 and a 28 residue insertion at residue 204 relative to the corresponding region of SEQ ID NO:2.

The resulting nucleotide sequence for the clone containing the 431 bp band (lower) is shown in SEQ ID NO:36, which corresponds to the 142 residue amino  
 20 acid sequence shown in SEQ ID NO:37. A composite of the polypeptide sequence of SEQ ID NO:2 with SEQ ID NO:37 results in a 173 residue protein shown in SEQ ID NO:38. This variant of ztnfr11 differs from the polypeptide sequence as shown in SEQ ID NO:2 by a translation stop codon resulting in a truncated soluble ztnfr11 receptor.

25

### Example 13

#### Identification of cells expressing ztnfr11 using *in situ* hybridization

Specific human tissues were isolated and screened for ztnfr11 expression by *in situ* hybridization. Various human tissues prepared, sectioned and subjected to *in situ* hybridization included normal skin, melanoma skin, squamous cell carcinoma,  
 30 psoriasis and cutaneous T-cell lymphoma skin samples, normal and adenocarcinoma colon and inflamed appendix samples. The tissues were fixed in 10% buffered formalin

and blocked in paraffin using standard techniques. Tissues were sectioned at 4 to 8 microns. Tissues were prepared using a standard protocol ("Development of non-isotopic *in situ* hybridization" at <http://dir.niehs.nih.gov/dirlep/ish.html>). Briefly, tissue sections were deparaffinized with HistoClear (National Diagnostics, Atlanta, GA) and then dehydrated with ethanol. Next they were digested with Proteinase K (50 µg/ml) (Boehringer Diagnostics, Indianapolis, IN) at 37°C for 2 to 20 minutes. This step was followed by acetylation and re-hydration of the tissues.

Two *in situ* probes generated by PCR were designed against the human ztnfr11 sequence. Two sets of oligos were designed to generate probes for separate regions of the ztnfr11 cDNA: (1) Oligos ZC26463 (SEQ ID NO:40) and ZC26464 (SEQ ID NO:41) were used to generate a 439 bp probe for ztnfr11 (from n.a. #576 to 1014); (2) ZC26470 (SEQ ID NO:42) and ZC26471 (SEQ ID NO:43) were used to generate a 548 bp probe for ztnfr11 (from n.a. #308 to 856). The antisense oligo from each set also contained the working sequence for the T7 RNA polymerase promoter to allow for easy transcription of antisense RNA probes from these PCR products. The PCR reaction conditions were as follows: 2 cycles at 95°C for 30 sec, 50°C for 1 min, 72°C for 1 min followed by 33 cycles of 95°C for 30 sec, 72°C for 2 min. Probes were subsequently labeled with digoxigenin (Boehringer) or biotin (Boehringer) using an *In Vitro* transcription System (Promega, Madison, WI) as per manufacturer's instruction.

*In situ* hybridization was performed with a digoxigenin- or biotin-labeled ztnfr11 probe (above). The probe was added to the slides at a concentration of 1 to 5 pmol/ml for 12 to 16 hours at 60°C. Slides were subsequently washed in 2XSSC and 0.1XSSC at 55°C. The signals were amplified using tyramide signal amplification (TSA) (TSA, *in situ* indirect kit; NEN) and visualized with Vector Red substrate kit (Vector Lab) as per manufacturer's instructions. The slides were then counter-stained with hematoxylin (Vector Laboratories, Burlingame, CA).

Positive signals were seen in the human cutaneous T-cell lymphoma and melanoma skin samples. The positive-staining cells appeared to be keratinocytes in the cutaneous lymphoma skin, and very weak signal in melanocytes of melanoma skin samples. All other tissue samples tested were negative for ztnfr11.

### Example 14

#### Construction of an Assay Cell Line

A BaF3 assay cell line was constructed for ztnfr11 ligand cloning. A  
 5 ztnfr11/TNFR1 chimera was built in which the transmembrane and cytoplasmic domain  
 of ztnfr11 were replaced by those of the TNF $\alpha$  receptor (TNFR1) (SEQ ID NO:44). The  
 ztnfr11/TNFR1 chimera was transfected into a BaF3 cell line that contains a  
 KZ159/mIL4 reporter gene (SEQ ID NO:45). KZ159/mIL4 responds to the activation of  
 TNFR1 receptor by TNF $\alpha$ , triggering the expression and secretion of mIL4 that leads to  
 10 the proliferation of BaF3 cells.

The extracellular domain of ztnfr11 was amplified with PCR primers  
 ZC28835 (SEQ ID NO:46) and ZC28836 (SEQ ID NO:47) using ztnfr11 full-length  
 cDNA as a template. The PCR reaction was as follows: 20 cycles of 94°C for 30 sec and  
 15 68°C for 2 min, then four more min at 68°C and soak at 10°C. The transmembrane and  
 cytoplasmic domain of mouse TNFR1 was amplified with PCR primers ZC28830 (SEQ  
 ID NO:48) and ZC28837 (SEQ ID NO:49) using mouse placenta marathon cDNA as a  
 template. The PCR reaction was as follows: 35 cycles of 94°C for 30 sec and 68°C for 2  
 min, then four more min at 68°C and soak at 10°C. The PCR products were separated on  
 20 1% Agarose, a band of 500bp (for ztnfr11) and 700bp (for TNFR1) was excised and the  
 DNA was purified using a QiaexII™ purification kit (Qiagen, Valencia, CA) according  
 to the manufacturer's instruction.

The pZP7Z plasmid is a mammalian expression vector containing an  
 25 expression cassette having the CMV promoter and a human growth hormone terminator.  
 The plasmid also has an E. coli origin of replication, a mammalian selectable marker  
 expression unit having an SV40 promoter, an enhancer and an origin of replication, as  
 well as a Zeocin resistant gene, and SV40 terminator. About 30 ng of the restriction  
 digested ztnfr11 insert (cut with EcoRI and PinA1) and TNFR1 insert (cut with PinA1  
 30 and Xba1) and about 10 ng of the digested vector (cut with EcoRI and Xba1) were  
 ligated at 11°C overnight. One microliter of ligation reaction was electroporated into

DH10B competent cells (Gibco BRL, Rockville, MD) according to manufacturer's direction and plated onto LB plates containing 50 ug/ml ampicillin, and incubated overnight. Colonies were screened by restriction analysis of DNA, which was prepared from 2ml liquid cultures of individual colonies. The insert sequence of positive clones was verified by sequencing analysis (SEQ ID NO:50).

Ten million BaF3 cells that contain KZ159/mIL4 reporter were transfected with ztnfr11/TNFR1/pZP7Z plasmid by electroporation, and selected for Zeocin resistant stable transfectants. The proliferation of this stable cell line in response to TNF $\alpha$  without mIL-3 was assayed, and the activation of ztnfr11/TNFR1 chimera was tested using cross-linked anti-ztnfr11 reagent which was prepared by incubating rabbit anti-ztnfr11 antibody at 1 ug/ml with anti-rabbit immunoglobulin G (IgG) antibody-coupled magnetic beads. The positive clones then can be used as an assay cell line to screen for ztnfr11 ligand.

#### Example 15

##### Construction of a Transgenic Plasmid

Twenty  $\mu$ g of the pzp9 plasmid containing the ztnfr11-Fc4 soluble receptor sequence from Example 5 was digested with EcoRI and XbaI. The approximately 1kb ztnfr11-Fc4 soluble receptor fragment was then isolated by running the digested vector on a 1.2% SeaPlaque GTG<sup>®</sup> gel and excising the fragment. DNA was purified using the QiaQuick<sup>™</sup> (Qiagen) gel extraction kit. The EcoRI and XbaI overhangs were then converted to blunt ends using Klenow polymerase.

The ztnfr11-Fc4 soluble receptor fragment was then ligated into a apoA1 C1-17 transgenic vector, which was previously digested with PmeI. The apoA1 C1-17 plasmid was designed for expression of a gene of interest in transgenic mice. It contains an expression cassette comprised of the apoA1 promoter, the first apoA1 exon and a portion of the first intron, the ztnfr11-Fc4 coding sequence, a polylinker for the insertion of the desired clone and the human growth hormone poly A sequence.

About one microliter of the ligation reaction was electroporated into DH10B ElectroMax<sup>®</sup> competent cells (GIBCO BRL) according to manufacturer's direction, plated onto LB plates containing 100  $\mu$ g/ml ampicillin, and incubated overnight at 37°C. Colonies were picked and grown in LB media containing 100  $\mu$ g/ml ampicillin. Miniprep DNA was prepared from the picked clones and screened for the correctly oriented ztnfr11-Fc4 soluble receptor insert by restriction digestion analysis and subsequent agarose gel electrophoresis. Maxipreps of the correct apoA1 C1-17 ztnfr11-Fc4 soluble receptor construct were performed.

A NotI fragment containing expression cassette was prepared and used for microinjection into fertilized murine oocytes.

5

From the foregoing, it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration,  
10 various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215



## CLAIMS

What is claimed is:

1. An isolated polypeptide comprising residues 2 to 129 of SEQ ID NO:2.
2. The isolated polypeptide according to claim 1, wherein the polypeptide comprises residues 2 to X of SEQ ID NO:2; wherein X is an integer between 130 and 136.
3. The isolated polypeptide according to claim 1, wherein the polypeptide comprises residues 1 to 129 of SEQ ID NO:2.
4. The isolated polypeptide according to claim 3, wherein the polypeptide comprises residues 1 to X of SEQ ID NO:2, wherein X is an integer between 130 and 136.
5. The isolated polypeptide according to claim 1, wherein the polypeptide is selected from the group consisting of:
  - a) polypeptides comprising residues 1 to 161 of SEQ ID NO:2; and
  - b) polypeptides comprising residues 1 to 269 of SEQ ID NO:2.
6. An isolated polypeptide selected from the group consisting of:
  - a) polypeptides comprising residues 1 to 297 of SEQ ID NO:27;
  - b) polypeptides comprising residues 1 to 267 of SEQ ID NO:29; and
  - c) polypeptides comprising residues 1 to 299 of SEQ ID NO:35.
7. An isolated polypeptide comprising residues 1 to 173 of SEQ ID NO:38.
8. An isolated polypeptide selected from the group consisting of:
  - a) polypeptide molecules comprising residues 1 to 147 of SEQ ID NO:38;
  - b) polypeptide molecules comprising residues 1 to 154 of SEQ ID NO:38;

- c) polypeptide molecules comprising residues 1 to 163 of SEQ ID NO:38;
- and
- d) polypeptide molecules comprising residues 1 to 165 of SEQ ID NO:38.
9. An isolated polynucleotide that encodes the polypeptide according to claim 1.
  10. A isolated polynucleotide that encodes the polypeptide according to claim 6.
  11. A isolated polynucleotide that encodes the polypeptide according to claim 7.
  12. A isolated polynucleotide that encodes the polypeptide according to claim 8.
  13. An expression vector comprising the following operably linked elements:
    - a) a transcription promoter;
    - b) a DNA segment wherein the DNA segment is a polynucleotide molecule encoding the polypeptide molecule of claim 3; and
    - c) a transcription terminator.
  14. The expression vector according to claim 13 wherein the DNA segment contains an affinity tag.
  15. A cultured cell into which has been introduced an expression vector according to claim 13, wherein said cell expresses the polypeptide encoded by the DNA segment.

16. A method of producing a polypeptide comprising culturing a cell according to claim 15, whereby said cell expresses the polypeptide encoded by the DNA segment; and recovering the polypeptide.

17. The polypeptide produced by the method of claim 16.

18. A method of detecting lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma comprising, contacting said lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma with a polynucleotide consisting of the polynucleotide sequence as shown in SEQ ID NOs:1, 31, 32, 33, or 36, wherein the polynucleotide hybridizes to the nucleic acid in the lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma.

19. The method of detecting lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma according to claim 18, wherein the polynucleotide is selected from the group consisting of:

- a) a polynucleotide consisting of at least 16 contiguous nucleotides as shown in SEQ ID NO:1;
- b) a polynucleotide consisting of from 17 to 25 contiguous nucleotides as shown in SEQ ID NO:1;
- c) a polynucleotide consisting of 40 contiguous nucleotides as shown in SEQ ID NO:1;
- d) a polynucleotide consisting of 60 contiguous nucleotides as shown in SEQ ID NO:1;
- e) a polynucleotide consisting of at least 16 contiguous nucleotides as shown in SEQ ID NO:31;
- f) a polynucleotide consisting of from 17 to 25 contiguous nucleotides as shown in SEQ ID NO:31;
- g) a polynucleotide consisting of 40 contiguous nucleotides as shown in SEQ ID NO:31;

- h) a polynucleotide consisting of 60 contiguous nucleotides as shown in SEQ ID NO:31;
- i) a polynucleotide consisting of at least 16 contiguous nucleotides as shown in SEQ ID NO:33;
- j) a polynucleotide consisting of from 17 to 25 contiguous nucleotides as shown in SEQ ID NO:33;
- k) a polynucleotide consisting of 40 contiguous nucleotides as shown in SEQ ID NO:33;
- l) a polynucleotide consisting of 60 contiguous nucleotides as shown in SEQ ID NO:33;
- m) a polynucleotide consisting of at least 16 contiguous nucleotides as shown in SEQ ID NO:36;
- n) a polynucleotide consisting of from 17 to 25 contiguous nucleotides as shown in SEQ ID NO:36;
- o) a polynucleotide consisting of 40 contiguous nucleotides as shown in SEQ ID NO:36;
- p) a polynucleotide consisting of 60 contiguous nucleotides as shown in SEQ ID NO:36;
- q) a polynucleotide encoding a polypeptide consisting of at least 16 contiguous nucleotides as shown in SEQ ID NO:38;
- r) a polynucleotide encoding a polypeptide consisting of from 17 to 25 contiguous nucleotides as shown in SEQ ID NO:38;
- s) a polynucleotide encoding a polypeptide consisting of 40 contiguous nucleotides as shown in SEQ ID NO:38;
- t) a polynucleotide encoding a polypeptide consisting of 60 contiguous nucleotides as shown in SEQ ID NO:38; and
- u) a polynucleotide encoding a polypeptide, wherein the polypeptide comprises residues 139 to 297 of SEQ ID NO:27, or residues 139 to 267 of SEQ ID NO:29.

v) A method of detecting lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma comprising, contacting said lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma with an antibody to the polypeptide according to claim 2 .

20. A method of detecting lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma according to claim 19 wherein the antibody is generated to a polypeptide selected from the group consisting of:

- a) a polypeptide comprising residues 4 to 13 of SEQ ID NO:2;
- b) a polypeptide comprising residues 17 to 39 of SEQ ID NO:2;
- c) a polypeptide comprising residues 44 to 60 of SEQ ID NO:2;
- d) a polypeptide comprising residues 69 to 104 of SEQ ID NO:2;
- e) a polypeptide comprising residues 106 to 115 of SEQ ID NO:2;
- f) a polypeptide comprising residues 128 to 136 of SEQ ID NO:2;
- g) a polypeptide comprising residues 165 to 174 of SEQ ID NO:2;
- h) a polypeptide comprising residues 179 to 204 of SEQ ID NO:2;
- i) a polypeptide comprising residues 245 to 260 of SEQ ID NO:2;
- j) a polypeptide comprising residues 1 to 13 of SEQ ID NO:2;
- k) a polypeptide comprising residues 21 to 38 of SEQ ID NO:2;
- l) a polypeptide comprising residues 44 to 59 of SEQ ID NO:2;
- m) a polypeptide comprising residues 88 to 99 of SEQ ID NO:2;
- n) a polypeptide comprising residues 165 to 173 of SEQ ID NO:2;
- o) a polypeptide comprising residues 194 to 205 of SEQ ID NO:2;
- and
- p) a polypeptide comprising residues 213 to 221 of SEQ ID NO:2.

21. A method of detecting lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma according to claim 19 wherein the antibody is generated to a polypeptide selected from the group consisting of:

- a) a polypeptide comprising residues 181 to 210 of SEQ ID NO:35;
- b) a polypeptide comprising residues 219 to 226 of SEQ ID NO:35;
- c) a polypeptide comprising residues 218 to 236 of SEQ ID NO:35;
- d) a polypeptide comprising residues 162 to 169 of SEQ ID NO:38;
- e) a polypeptide comprising residues 149 to 173 of SEQ ID NO:38;
- f) a polypeptide comprising residues 194 to 236 of SEQ ID NO:35;
- g) a polypeptide comprising residues 150 to 171 of SEQ ID NO:38;
- h) a polypeptide comprising residues 230 to 235 of SEQ ID NO:35;
- and
- i) a polypeptide comprising residues 151 to 158 of SEQ ID NO:38.

22. A method of detecting lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma according to claim 19 wherein the antibody is generated to a polypeptide selected from the group consisting of:

- a) a polypeptide consisting of residues 4 to 13 of SEQ ID NO:2;
- b) a polypeptide consisting of residues 17 to 39 of SEQ ID NO:2;
- c) a polypeptide consisting of residues 44 to 60 of SEQ ID NO:2;
- d) a polypeptide consisting of residues 69 to 104 of SEQ ID NO:2;
- e) a polypeptide consisting of residues 106 to 115 of SEQ ID NO:2;
- f) a polypeptide consisting of 128 to 136 of SEQ ID NO:2;
- g) a polypeptide consisting of residues 165 to 174 of SEQ ID NO:2;
- h) a polypeptide consisting of residues 179 to 204 of SEQ ID NO:2;
- i) a polypeptide consisting of residues 245 to 260 of SEQ ID NO:2;
- j) a polypeptide consisting of residues 1 to 13 of SEQ ID NO:2;
- k) a polypeptide consisting of residues 21 to 38 of SEQ ID NO:2;
- l) a polypeptide consisting of residues 44 to 59 of SEQ ID NO:2;
- m) a polypeptide consisting of residues 88 to 99 of SEQ ID NO:2;
- n) a polypeptide consisting of residues 165 to 173 of SEQ ID NO:2;
- o) a polypeptide consisting of residues 194 to 205 of SEQ ID NO:2;
- and
- p) a polypeptide consisting of residues 213 to 221 of SEQ ID NO:2.

23. A method of detecting lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma according to claim 19 wherein the antibody is generated to a polypeptide selected from the group consisting of:

- a) a polypeptide consisting of residues 181 to 210 of SEQ ID NO:35;
- b) a polypeptide consisting of residues 219 to 226 of SEQ ID NO:35;
- c) a polypeptide consisting of residues 218 to 236 of SEQ ID NO:35;
- d) a polypeptide consisting of residues 162 to 169 of SEQ ID NO:38;
- e) a polypeptide consisting of residues 149 to 173 of SEQ ID NO:38;
- f) a polypeptide consisting of residues 194 to 236 of SEQ ID NO:35;
- g) a polypeptide consisting of residues 150 to 171 of SEQ ID NO:38;
- h) a polypeptide consisting of residues 230 to 235 of SEQ ID NO:35;
- and
- i) a polypeptide consisting of residues 151 to 158 of SEQ ID NO:38.

24. A method of inhibiting the quantity of lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma cells expressing a polypeptide selected from the group consisting of:

- a) an isolated polypeptide as shown in SEQ ID NO:2 ;
- b) an isolated polypeptide as shown in SEQ ID NO:27;
- c) an isolated polypeptide as shown in SEQ ID NO:29; and
- d) an isolated polypeptide as shown in SEQ ID NO:35;

comprising administering to the cells an isolated polypeptide wherein the isolated polypeptide consists of residues 1 to X of SEQ ID NO:2, wherein X is an integer between 129 and 136.

25. A method of modulating apoptosis of lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma cells comprising administering to the cells an isolated polypeptide wherein the isolated polypeptide comprises the polypeptide according to claim 2.

26. The method of modulating apoptosis of lung carcinoma, breast carcinoma, melanoma, osteosarcoma, or lymphoma cells according to claim 25, wherein the isolated polypeptide comprises the polypeptide according to claim 4.

27. An isolated polypeptide consisting of residues 2 to 129 of SEQ ID NO:2.

28. The isolated polypeptide according to claim 27, wherein the polypeptide consists of residues 2 to X of SEQ ID NO:2; wherein X is an integer between 130 and 136.

29. The isolated polypeptide according to claim 27, wherein the polypeptide consists of residues 1 to 129 of SEQ ID NO:2.

30. The isolated polypeptide according to claim 29, wherein the polypeptide consists of residues 1 to X of SEQ ID NO:2, wherein X is an integer between 130 and 136.

31. The isolated polypeptide according to claim 27, wherein the polypeptide is selected from the group consisting of:

- a) polypeptides consisting of residues 1 to 161 of SEQ ID NO:2; and
- b) polypeptides consisting of residues 1 to 269 of SEQ ID NO:2.

32. An isolated polypeptide selected from the group consisting of:

- a) polypeptides consisting of residues 1 to 297 of SEQ ID NO:27;
- b) polypeptides consisting of residues 1 to 267 of SEQ ID NO:29; and
- c) polypeptides consisting of residues 1 to 299 of SEQ ID NO:35.

33. An isolated polypeptide consisting of residues 1 to 173 of SEQ ID NO:38.



34. An isolated polypeptide selected from the group consisting of:
- a) polypeptide molecules consisting of residues 1 to 147 of SEQ ID NO:38;
- b) polypeptide molecules consisting of residues 1 to 154 of SEQ ID NO:38;
- c) polypeptide molecules consisting of residues 1 to 163 of SEQ ID NO:38; and
- d) polypeptide molecules consisting of residues 1 to 165 of SEQ ID NO:38.
35. An isolated polynucleotide that encodes the polypeptide according to claim 27.
36. A isolated polynucleotide that encodes the polypeptide according to claim 32.
37. A isolated polynucleotide that encodes the polypeptide according to claim 33.
38. A isolated polynucleotide that encodes the polypeptide according to claim 34.

## UMLR Polypeptides

## ABSTRACT OF THE DISCLOSURE

Novel secreted and membrane bound tumor necrosis factor receptor (TNFR) polypeptides, polynucleotides encoding the polypeptides, antibodies and related compositions and methods are disclosed. The polypeptides may be used for detecting ligands, agonists and antagonists. The polypeptides, polynucleotides and antibodies may also be used in methods that modulate inflammation, tumor growth and metastasis, infection, immunity, and cellular maturation.

## SEQUENCE LISTING

<110> Xu, Wenfeng  
 Lofton-Day, Catherine E.  
 Henne, Randall  
 Presnell, Scott R.  
 Yao, Yue  
 Novak, Julia E.  
 Foster, Donald C.  
 Yee, David P.

<120> UMLR POLYPEPTIDES

<130> 99-75

<150> US 60/160,880

<151> 1999-10-22

<150> US 60/163,215

<151> 1999-11-02

<150> US 60/218,769

<151> 2000-07-17

<150> US 60/222,221

<151> 2000-08-01

<160> 50

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1162

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (104)...(913)

<400> 1

gagggggctg ggtgagatgt gtgctctgcg ctgaggtgga tttgtaccgg agtcccattt

gggagcaaga gccatctact cgtccgttac cggccttccc acc atg gat tgc caa	115
Met Asp Cys Gln	
1	
gaa aat gag tac tgg gac caa tgg gga cgg tgt gtc acc tgc caa cgg	163
Glu Asn Glu Tyr Trp Asp Gln Trp Gly Arg Cys Val Thr Cys Gln Arg	
5 10 15 20	
tgt ggt cct gga cag gag cta tcc aag gat tgt ggt tat gga gag ggt	211
Cys Gly Pro Gly Gln Glu Leu Ser Lys Asp Cys Gly Tyr Gly Glu Gly	
25 30 35	
gga gat gcc tac tgc aca gcc tgc cct cct cgc agg tac aaa agc agc	259
Gly Asp Ala Tyr Cys Thr Ala Cys Pro Pro Arg Arg Tyr Lys Ser Ser	
40 45 50	
tgg ggc cac cac aaa tgt cag agt tgc atc acc tgt gct gtc atc aat	307
Trp Gly His His Lys Cys Gln Ser Cys Ile Thr Cys Ala Val Ile Asn	
55 60 65	
cgt gtt cag aag gtc aac tgc aca gct acc tct aat gct gtc tgt ggg	355
Arg Val Gln Lys Val Asn Cys Thr Ala Thr Ser Asn Ala Val Cys Gly	
70 75 80	
gac tgt ttg ccc agg ttc tac cga aag aca cgc att gga ggc ctg cag	403
Asp Cys Leu Pro Arg Phe Tyr Arg Lys Thr Arg Ile Gly Gly Leu Gln	
85 90 95 100	
gac caa gag tgc atc ccg tgc acg aag cag acc ccc acc tct gag gtt	451
Asp Gln Glu Cys Ile Pro Cys Thr Lys Gln Thr Pro Thr Ser Glu Val	
105 110 115	
caa tgt gcc ttc cag ttg agc tta gtg gag gca gat gca ccc aca gtg	499
Gln Cys Ala Phe Gln Leu Ser Leu Val Glu Ala Asp Ala Pro Thr Val	
120 125 130	
ccc cct cag gag gcc aca ctt gtt gca ctg gtg agc agc ctg cta gtg	547
Pro Pro Gln Glu Ala Thr Leu Val Ala Leu Val Ser Ser Leu Leu Val	
135 140 145	
gtg ttt acc ctg gcc ttc ctg ggg ctc ttc ttc ctc tac tgc aag cag	595
Val Phe Thr Leu Ala Phe Leu Gly Leu Phe Phe Leu Tyr Cys Lys Gln	
150 155 160	

ttc ttc aac aga cat tgc cag cgt gga ggt ttg ctg cag ttt gag gct	643
Phe Phe Asn Arg His Cys Gln Arg Gly Gly Leu Leu Gln Phe Glu Ala	
165 170 175 180	
gat aaa aca gca aag gag gaa tct ctc ttc ccc gtg cca ccc agc aag	691
Asp Lys Thr Ala Lys Glu Glu Ser Leu Phe Pro Val Pro Pro Ser Lys	
185 190 195	
gag acc agt gct gag tcc caa gag tcc ttt acc atg gcc tcc tgc acc	739
Glu Thr Ser Ala Glu Ser Gln Glu Ser Phe Thr Met Ala Ser Cys Thr	
200 205 210	
tca gag agc cac tcc cac tgg gtc cac agc ccc atc gaa tgc aca gag	787
Ser Glu Ser His Ser His Trp Val His Ser Pro Ile Glu Cys Thr Glu	
215 220 225	
ctg gac ctg caa aag ttt tcc agc tct gcc tcc tat act gga gct gag	835
Leu Asp Leu Gln Lys Phe Ser Ser Ser Ala Ser Tyr Thr Gly Ala Glu	
230 235 240	
acc ttg ggg gga aac aca gtc gaa agc act gga gac agg ctg gag ctc	883
Thr Leu Gly Gly Asn Thr Val Glu Ser Thr Gly Asp Arg Leu Glu Leu	
245 250 255 260	
aat gtg ccc ttt gaa gtt ccc agc cct taa ctctaagag gtctcttggg	933
Asn Val Pro Phe Glu Val Pro Ser Pro *	
265	
cccctggcag ccttgcccag ttgttctctc tggactctgt tcctatacca caacagcagc	993
aggggcctga aatgtgatgt ccacaagagc taatacccta cagatggggc atatcctatc	1053
ccatcccacc agaggattga ttctccatct cacaaggact gatctggagc atttcttgct	1113
tccctgttgt agtctgggga gccagattcc acatgcatgg ggcggccgc	1162

&lt;210&gt; 2

&lt;211&gt; 269

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 2

Met Asp Cys Gln Glu Asn Glu Tyr Trp Asp Gln Trp Gly Arg Cys Val
1 5 10 15

```

Thr Cys Gln Arg Cys Gly Pro Gly Gln Glu Leu Ser Lys Asp Cys Gly
      20              25              30
Tyr Gly Glu Gly Gly Asp Ala Tyr Cys Thr Ala Cys Pro Pro Arg Arg
      35              40              45
Tyr Lys Ser Ser Trp Gly His His Lys Cys Gln Ser Cys Ile Thr Cys
      50              55              60
Ala Val Ile Asn Arg Val Gln Lys Val Asn Cys Thr Ala Thr Ser Asn
      65              70              75              80
Ala Val Cys Gly Asp Cys Leu Pro Arg Phe Tyr Arg Lys Thr Arg Ile
      85              90              95
Gly Gly Leu Gln Asp Gln Glu Cys Ile Pro Cys Thr Lys Gln Thr Pro
      100             105             110
Thr Ser Glu Val Gln Cys Ala Phe Gln Leu Ser Leu Val Glu Ala Asp
      115             120             125
Ala Pro Thr Val Pro Pro Gln Glu Ala Thr Leu Val Ala Leu Val Ser
      130             135             140
Ser Leu Leu Val Val Phe Thr Leu Ala Phe Leu Gly Leu Phe Phe Leu
      145             150             155             160
Tyr Cys Lys Gln Phe Phe Asn Arg His Cys Gln Arg Gly Gly Leu Leu
      165             170             175
Gln Phe Glu Ala Asp Lys Thr Ala Lys Glu Glu Ser Leu Phe Pro Val
      180             185             190
Pro Pro Ser Lys Glu Thr Ser Ala Glu Ser Gln Glu Ser Phe Thr Met
      195             200             205
Ala Ser Cys Thr Ser Glu Ser His Ser His Trp Val His Ser Pro Ile
      210             215             220
Glu Cys Thr Glu Leu Asp Leu Gln Lys Phe Ser Ser Ser Ala Ser Tyr
      225             230             235             240
Thr Gly Ala Glu Thr Leu Gly Gly Asn Thr Val Glu Ser Thr Gly Asp
      245             250             255
Arg Leu Glu Leu Asn Val Pro Phe Glu Val Pro Ser Pro
      260             265

```

<210> 3  
 <211> 807  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> degenerate sequence

<221> misc\_feature  
 <222> (1)...(807)

<223> n = A,T,C or G

<400> 3

atggaytgyc	argaraayga	rtaytggg	cartggggnm	gntgygtnac	ntgycarmgn	60
tgyggncng	gncargaryt	nwsnaargay	tgyggntayg	gngarggngg	ngaygcntay	120
tgyacngcnt	gyccnccnmg	nmgtayaar	wsnwsntggg	gncaycayaa	rtgycarwsn	180
tgyathacnt	gygcngtnat	haaymgngtn	caraargtna	aytgyacngc	nacnwsnaay	240
gcngtntgyg	gngaytgyyt	nccnmgtty	taymgnara	cnmgnathgg	nggnytnear	300
gaycargart	gyathccntg	yacnaarcar	acnccnacnw	sngargtnca	rtgygcntty	360
carytnwsny	tngtngargc	ngaygcncn	acngtnccnc	cncargargc	nacnytnngtn	420
gcnytnngtnw	snwsnytnyt	ngtngtntty	acnytnngcnt	tyytnnggny	nttyttyytn	480
taytgyaarc	arttyttaa	ymgncaytgy	carmgngngg	gnytnytnca	rttygargcn	540
gayaaracng	cnaargarga	rwsnytnntty	ccngtnccnc	cnwsnaarga	racnwsngcn	600
garwsncarg	arwsnttyac	natggcnwsn	tgyacnwsng	arwsncayws	ncaytgggtn	660
caywsnccna	thgartgyac	ngarytngay	ytncaraart	tywsnwsnws	ngcnwsntay	720
acnggngcng	aracnytnng	nggnaayacn	gtngarwsna	cnggngaymg	nytngarytn	780
aaygtncnt	tygargtncc	nwsnccn				807

<210> 4

<211> 41

<212> PRT

<213> Artificial Sequence

<220>

<223> Pseudo repeat motif #1

<221> VARIANT

<222> (1)...(1)

<223> Xaa is any amino acid residue

<221> VARIANT

<222> (3)...(12)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (13)...(16)

<223> Each Xaa is independently any amino acid residue  
or not present

<221> VARIANT

<222> (19)...(20)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (22)...(26)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (27)...(30)

<223> Each Xaa is independently any amino acid residue  
or not present

<221> VARIANT

<222> (32)...(37)

<223> Each Xaa is independently any amino acid residue  
or not present

<221> VARIANT

<222> (38)...(39)

<223> Each Xaa is independently any amino acid residue  
or not present

<221> VARIANT

<222> (41)...(41)

<223> Xaa is any amino acid residue

<400> 4

Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
1			5				10					15			
Cys	Cys	Xaa	Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Cys	Xaa	
			20				25					30			
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Cys	Xaa							
			35				40								

<210> 5

<211> 45

<212> PRT

<213> Artificial Sequence

<220>

<223> Pseudo repeat motif #2

<221> VARIANT

<222> (1)...(1)

<223> Xaa is any amino acid residue



<221> VARIANT  
 <222> (3)...(15)  
 <223> Each Xaa is independently any amino acid residue

<221> VARIANT  
 <222> (16)...(17)  
 <223> Each Xaa is independently any amino acid residue  
 or not present

<221> VARIANT  
 <222> (19)...(20)  
 <223> Each Xaa is independently any amino acid residue

<221> VARIANT  
 <222> (22)...(23)  
 <223> Each Xaa is independently any amino acid residue

<221> VARIANT  
 <222> (24)...(24)  
 <223> Each Xaa is independently any amino acid residue  
 or not present

<221> VARIANT  
 <222> (26)...(33)  
 <223> Each Xaa is independently any amino acid residue

<221> VARIANT  
 <222> (34)...(36)  
 <223> Each Xaa is independently any amino acid residue  
 or not present

<221> VARIANT  
 <222> (38)...(44)  
 <223> Each Xaa is independently any amino acid residue

<400> 5

Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
1			5				10					15			
Xaa	Cys	Xaa	Xaa	Cys	Xaa	Xaa	Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
			20				25					30			
Xaa	Xaa	Xaa	Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Cys			
			35				40					45			

<210> 6  
 <211> 49  
 <212> PRT  
 <213> Artificial Sequence

<220>  
 <223> Pseudo repeat motif #3

<221> VARIANT  
 <222> (1)...(1)  
 <223> Xaa is any amino acid residue

<221> VARIANT  
 <222> (3)...(7)  
 <223> Each Xaa is independently any amino acid residue

<221> VARIANT  
 <222> (8)...(8)  
 <223> Xaa is any amino acid residue or not present

<221> VARIANT  
 <222> (9)...(14)  
 <223> Each Xaa is independently any amino acid residue

<221> VARIANT  
 <222> (15)...(18)  
 <223> Each Xaa is independently any amino acid residue  
 or not present

<221> VARIANT  
 <222> (20)...(21)  
 <223> Each Xaa is independently any amino acid residue

<221> VARIANT  
 <222> (23)...(24)  
 <223> Each Xaa is independently any amino acid residue

<221> VARIANT  
 <222> (25)...(29)  
 <223> Each Xaa is independently any amino acid residue  
 or not present

<221> VARIANT

<222> (31)...(38)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (39)...(39)

<223> Xaa is any amino acid residue or not present

<221> VARIANT

<222> (41)...(47)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (49)...(49)

<223> Xaa is any amino acid residue or not present

<400> 6

Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
1			5					10							15	
Xaa	Xaa	Cys	Xaa	Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Cys	Xaa	Xaa	
			20					25					30			
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Cys
			35				40						45			
Xaa																

<210> 7

<211> 48

<212> PRT

<213> Artificial Sequence

<220>

<223> Pseudo repeat motif #3 alternative motif

<221> VARIANT

<222> (1)...(1)

<223> Xaa is any amino acid residue

<221> VARIANT

<222> (3)...(7)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (8)...(8)

<223> Xaa is independently any amino acid residue or not present

<221> VARIANT

<222> (10)...(13)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (14)...(18)

<223> Each Xaa is independently any amino acid residue or not present

<221> VARIANT

<222> (20)...(21)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (23)...(24)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (25)...(29)

<223> Each Xaa is independently any amino acid residue or not present

<221> VARIANT

<222> (31)...(40)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (41)...(46)

<223> Each Xaa is independently any amino acid residue or not present

<221> VARIANT

<222> (48)...(48)

<223> Xaa is any amino acid residue

<400> 7

Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
1			5				10					15		
Xaa	Xaa	Cys	Xaa	Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Cys	Xaa	Xaa
			20				25					30		

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa  
           35                                  40                                  45

<210> 8

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> Pseudo repeat motif #4

<221> VARIANT

<222> (1)...(1)

<223> Xaa is any amino acid residue

<221> VARIANT

<222> (3)...(12)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (13)...(16)

<223> Each Xaa is independently any amino acid residue  
 or not present

<221> VARIANT

<222> (18)...(19)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (21)...(22)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (24)...(27)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (28)...(33)

<223> Each Xaa is independently any amino acid residue  
 or not present

<221> VARIANT

<222> (35)...(37)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (38)...(41)

<223> Each Xaa is independently any amino acid residue  
or not present

<221> VARIANT

<222> (43)...(43)

<223> Each Xaa is independently any amino acid residue

<400> 8

Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
1			5					10						15		
Cys	Xaa	Xaa	Cys	Xaa	Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
			20					25						30		
Xaa	Cys	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Cys	Xaa						
			35					40								

<210> 9

<211> 43

<212> PRT

<213> Artificial Sequence

<220>

<223> Pseudo repeat motif #4 alternative motif

<221> VARIANT

<222> (1)...(1)

<223> Xaa is any amino acid residue

<221> VARIANT

<222> (3)...(12)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT

<222> (13)...(16)

<223> Each Xaa is independently any amino acid residue  
or not present

<221> VARIANT

<222> (18)...(22)

<223> Each Xaa is independently any amino acid residue

<221> VARIANT  
 <222> (24)...(27)  
 <223> Each Xaa is independently any amino acid residue

<221> VARIANT  
 <222> (28)...(33)  
 <223> Each Xaa is independently any amino acid residue  
 or not present

<221> VARIANT  
 <222> (34)...(37)  
 <223> Each Xaa is independently any amino acid residue

<221> VARIANT  
 <222> (38)...(41)  
 <223> Each Xaa is independently any amino acid residue  
 or not present

<221> VARIANT  
 <222> (43)...(43)  
 <223> Xaa is any amino acid residue

<400> 9  
 Xaa Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa  
 1 5 10 15  
 Cys Xaa Xaa Xaa Xaa Xaa Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa  
 20 25 30  
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa  
 35 40

<210> 10  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide primer ZC25352

<400> 10  
 ccttgcccag ttgttctc

<210> 11

<211> 18  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide primer ZC25353

<400> 11  
 tctggtggga tgggatag 18

<210> 12  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide primer ZC25364

<400> 12  
 acctgtgctg tcatcaatcg tgttca 26

<210> 13  
 <211> 24  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide primer ZC25365

<400> 13  
 cccccaaggt ctcagctcca gtat 24

<210> 14  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide primer ZC25352

<400> 14  
 ccttgcccag ttgttctc 18



<210> 15  
 <211> 18  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide primer ZC25353

<400> 15  
 tctggtggga tgggatatag 18

<210> 16  
 <211> 6  
 <212> PRT  
 <213> Artificial protein

<400> 16  
 Glu Tyr Met Pro Met Glu  
 1 5

<210> 17  
 <211> 33  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide primer ZC25598

<400> 17  
 gcggatccga ttgccaagaa aatgagtact ggg 33

<210> 18  
 <211> 37  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> oligonucleotide primer ZC25596

<400> 18  
 gcagatctgg gctccactgt ggtgcatct gcctcca 37

<210> 19  
 <211> 108  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <223> tPA leader

<400> 19  
 atggatgcaa tgaagagagg gctctgctgt gtgctgctgc tgtgtggcgc cgtcttcggt 60  
 tcgctcagcc aggaaatcca tgccgagttg agacgcttcc gtagatcc 108

<210> 20  
 <211> 693  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Fc4 tag

<400> 20  
 agatcttcag acaaaactca cacatgccca ccgtgccag cacctgaagc cgagggggca 60  
 ccgtcagtct tcctcttccc cccaaaaccc aaggacaccc tcgatgctc ccggaccctt 120  
 gaggtcacat gcgtgggtgt ggacgtgagc cacgaagacc ctgaggtcaa gttcaactgg 180  
 tacgtggacg gcgtggaggt gcataatgcc aagacaaagc cgcgaggaga gcagtacaac 240  
 agcacgtacc gtgtggtcag cgctctcacc gtcctgcacc aggactggct gaatggcaag 300  
 gagtacaagt gcaagggtctc caacaaagcc ctcccatcct ccatcgagaa aaccatctcc 360  
 aaagccaaag ggcagccccg agaaccacag gtgtacaccc tgccccatc ccgggatgag 420  
 ctgaccaaga accaggtcag cctgacctgc ctgggtcaaag gcttctatcc cagcgacatc 480  
 gccgtggagt gggagagcaa tgggcagccg gagaacaact acaagaccac gcctcccgtg 540  
 ctggactccg acggctcctt cttcctctac agcaagctca ccgtggacaa gaggcaggtg 600  
 cagcagggga acgtcttctc atgctccgtg atgcatgagg ctctgcacaa ccactacacg 660  
 cagaagagcc tctccctgtc tccgggtaaa taa 693

<210> 21  
 <211> 534  
 <212> DNA  
 <213> Artificial sequence

<400> 21  
 atggatgcaa tgaagagagg gctctgctgt gtgctgctgc tgtgtggcgc cgtcttcggt 60  
 tcgctcagcc aggaaatcca tgccgagttg agacgcttcc gtagatccga ttgccaagaa 120  
 aatgagtact gggaccaatg gggacggtgt gtcacctgcc aacggtgtgg tcctggacag 180

gagctatcca	aggattgtgg	ttatggagag	ggtggagatg	cctactgcac	agcctgccct	240
cctcgcaggt	acaaaagcag	ctggggccac	cacaaatgtc	agagttgcat	cacctgtgct	300
gtcatcaatc	gtgttcagaa	ggtcaactgc	acagctacct	ctaattgctgt	ctgtggggac	360
tgtttgccca	ggttctaccg	aaagacacgc	attggaggcc	tgcaggacca	agagtgcata	420
ccgtgcacga	agcagacccc	cacctctgag	gttcaatgtg	ccttccagtt	gagcttagtg	480
gaggcagatg	cacccacagt	ggagcccaga	tctgaatata	tgcccatgga	ataa	534

&lt;210&gt; 22

&lt;211&gt; 1200

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; construct

&lt;400&gt; 22

atggatgcaa	tgaagagagg	gctctgctgt	gtgctgctgc	tgtgtggcgc	cgtcttcggt	60
tcgctcagcc	aggaaatcca	tgccgagttg	agacgcttcc	gtagatccga	ttgccaagaa	120
aatgagtact	gggaccaatg	gggacggtgt	gtcacctgcc	aacggtgtgg	tcctggacag	180
gagctatcca	aggattgtgg	ttatggagag	ggtggagatg	cctactgcac	agcctgccct	240
cctcgcaggt	acaaaagcag	ctggggccac	cacaaatgtc	agagttgcat	cacctgtgct	300
gtcatcaatc	gtgttcagaa	ggtcaactgc	acagctacct	ctaattgctgt	ctgtggggac	360
tgtttgccca	ggttctaccg	aaagacacgc	attggaggcc	tgcaggacca	agagtgcata	420
ccgtgcacga	agcagacccc	cacctctgag	gttcaatgtg	ccttccagtt	gagcttagtg	480
gaggcagatg	cacccacagt	ggagcccaga	tcttcagaca	aaactcacac	atgccaccg	540
tgcccagcac	ctgaagccga	gggggcaccg	tcagtcttcc	tcttcccccc	aaaacccaag	600
gacaccctca	tgatctcccg	gaccctgag	gtcacatgcg	tggtggtgga	cgtgagccac	660
gaagaccctg	aggtcaagtt	caactggtac	gtggacggcg	tggaggtgca	taatgccaag	720
acaaagccgc	gggaggagca	gtacaacagc	acgtaccgtg	tggtcagcgt	cctcaccgtc	780
ctgcaccagg	actggctgaa	tggcaaggag	tacaagtgca	aggtctccaa	caaagccctc	840
ccatcctcca	tcgagaaaac	catctccaaa	gccaaagggc	agccccgaga	accacaggtg	900
tacaccctgc	ccccatcccg	ggatgagctg	accaagaacc	aggtcagcct	gacctgcctg	960
gtcaaaggct	tctatcccag	cgacatcgcc	gtggagtggg	agagcaatgg	gcagccggag	1020
aacaactaca	agaccacgcc	tcccgtgctg	gactccgacg	gctccttctt	cctctacagc	1080
aagctcaccg	tggacaagag	caggtggcag	caggggaacg	tcttctcatg	ctccgtgatg	1140
catgaggctc	tgcaacaacca	ctacacgcag	aagagcctct	ccctgtctcc	gggtaaataa	1200

&lt;210&gt; 23

&lt;211&gt; 47

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

<223> Oligonucleotide ZC26463

<400> 23

atgcattaac cctcactaaa gggccttcct ggggctcttc ttcctct

47

<210> 24

<211> 46

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide ZC26464

<400> 24

taatacgact cactataggg aggggcccct gctgctgttg tggat

46

<210> 25

<211> 49

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide ZC26470

<400> 25

atgcattaac cctcactaaa gggacctgtg ctgtcatcaa tcgtgttca

49

<210> 26

<211> 47

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide ZC26471

<400> 26

taatacgact cactataggg agggccccc aa ggtctcagct ccagtat

47

<210> 27

<211> 297

atgcattaac cctcactaaa gggccttcct ggggctcttc ttcctct



<210> 28  
 <211> 891  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> degenerate sequence

<221> misc\_feature  
 <222> (1)...(891)  
 <223> n = A,T,C or G

<400> 28  
 atggaytgyc argaraayga rtaytgggag cartgggggm gntgygtnac ntgyccarmgn 60  
 tgyggncnng gncargaryt nwsnaargay tgyggntayg gngarggngg ngaygcntay 120  
 tgyacngcnt gyccnccnmg nmgtayaar wsnwsntggg gncaycayaa rtgycarwsn 180  
 tgyathacnt gygcngtnat haaymgngtn caraargtna aytgyacngc nacnwsnaay 240  
 gcngtntgyg gngaytgyyt nccnmgtty taymgnaraa cnmgcnathgg nggnytnear 300  
 gaycargart gyathccntg yacnaarcar acnccnaccw sngargtnca rtgygcntty 360  
 carytnwsny tngtngargc ngaygcncn acngtnccnc cncargargc nacnytnngtn 420  
 gcnytnngtnw snwsnytnyt ngtngtntty acnytnngnt tyytnggnyt nttyttyytn 480  
 taytgyaarc arttyttaa ymgncaytgy carmgngngg gnytnytnca rtytgargcn 540  
 gyaaracng cnaargarga rwsnytnntty ccngtnccnc cnwsnaarga racnwsngcn 600  
 garwsncarg tnwsngaraa yathttycar acncarccny tnaayccnat hytngargay 660  
 gaytgywsnw snacnwsngg nttyccnaccn cargarwsnt tyacnatggc nwsntgyacn 720  
 wsnarwsnc aywsncaytg ggtncaywsn ccnathgart gyacngaryt ngayytnear 780  
 aarttywsnw snwsngcnws ntayacnggn gcngaracny tnggnggnaa yacngtnear 840  
 wsnacngngg aymgnytna rytnaaygtm ccnttygarg tnccnwsncc n 891

<210> 29  
 <211> 267  
 <212> PRT  
 <213> Homo sapiens

<400> 29  
 Met Asp Cys Gln Glu Asn Glu Tyr Trp Asp Gln Trp Gly Arg Cys Val  
 1 5 10 15  
 Thr Cys Gln Arg Cys Gly Pro Gly Gln Glu Leu Ser Lys Asp Cys Gly  
 20 25 30  
 Tyr Gly Glu Gly Gly Asp Ala Tyr Cys Thr Ala Cys Pro Pro Arg Arg  
 35 40 45  
 Tyr Lys Ser Ser Trp Gly His His Lys Cys Gln Ser Cys Ile Thr Cys  
 50 55 60

Ala Val Ile Asn Arg Val Gln Lys Val Asn Cys Thr Ala Thr Ser Asn  
 65 70 75 80  
 Ala Val Cys Gly Asp Cys Leu Pro Arg Phe Tyr Arg Lys Thr Arg Ile  
 85 90 95  
 Gly Gly Leu Gln Asp Gln Glu Cys Ile Pro Cys Thr Lys Gln Thr Pro  
 100 105 110  
 Thr Ser Glu Val Gln Cys Ala Phe Gln Leu Ser Leu Val Glu Ala Asp  
 115 120 125  
 Ala Pro Thr Val Pro Pro Gln Glu Ala Thr Leu Val Ala Leu Gly Gly  
 130 135 140  
 Leu Leu Gln Phe Glu Ala Asp Lys Thr Ala Lys Glu Glu Ser Leu Phe  
 145 150 155 160  
 Pro Val Pro Pro Ser Lys Glu Thr Ser Ala Glu Ser Gln Val Ser Glu  
 165 170 175  
 Asn Ile Phe Gln Thr Gln Pro Leu Asn Pro Ile Leu Glu Asp Asp Cys  
 180 185 190  
 Ser Ser Thr Ser Gly Phe Pro Thr Gln Glu Ser Phe Thr Met Ala Ser  
 195 200 205  
 Cys Thr Ser Glu Ser His Ser His Trp Val His Ser Pro Ile Glu Cys  
 210 215 220  
 Thr Glu Leu Asp Leu Gln Lys Phe Ser Ser Ser Ala Ser Tyr Thr Gly  
 225 230 235 240  
 Ala Glu Thr Leu Gly Gly Asn Thr Val Glu Ser Thr Gly Asp Arg Leu  
 245 250 255  
 Glu Leu Asn Val Pro Phe Glu Val Pro Ser Pro  
 260 265

<210> 30

<211> 801

<212> DNA

<213> Artificial Sequence

<220>

<223> degenerate sequence

<221> misc\_feature

<222> (1)...(801)

<223> n = A,T,C or G

<400> 30

atggaytgyc	argaraayga	rtaytgggay	cartggggnm	gntgygtnac	ntgycarmgn	60
tyggngccng	gncargaryt	nwsnaargay	tygggntayg	gngarggngg	ngaygcntay	120
tgyacngcnt	gyccnccnmg	nmgntayaar	wsnwsntggg	gncaycayaa	rtgycarwsn	180

tgathacnt	gygcngtnat	haaymgngtn	caraargtna	aytgyacngc	nacnwsnaay	240
gcngtntgyg	gngaytgyyt	nccnmgnatty	taymgnaara	cnmgcnathgg	nggnytnicar	300
gaycargart	gyathccntg	yacnaarcar	acnccnacnw	sngargtnca	rtgygcntty	360
carytnwsny	tngtngargc	ngaygcncn	acngtnccnc	cncargargc	nacnytngt	420
gcnytnggng	gnytnytnc	rttygargcn	gayaaracng	cnaargarga	rwsnytnntty	480
ccngtnccnc	cnwsnaarga	racnwsngcn	garwsncarg	tnwsngaraa	yathttycar	540
acncarceny	tnaayccnat	hytnargay	gaytgywsnw	snacnwsngg	nttyccnacn	600
cargarwsnt	tyacnatggc	nwsntgyacn	wsngarwsnc	aywsncaytg	ggtncayswn	660
ccnathgart	gyacngaryt	ngayytncar	aarttywsnw	snwsngcnws	ntayacnggn	720
gcngaracny	tnggnggnaa	yacngtngar	wsnacnggng	aymgnytnga	rytnaaygt	780
ccnttygarg	tnccnwsncc	n				801

&lt;210&gt; 31

&lt;211&gt; 529

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(529)

&lt;223&gt; n = A,T,C or G

&lt;400&gt; 31

ggattcnatn	nctgaggntg	natggcnttc	nagttnwgas	tkagtggagg	cagatgcasc	60
cacagtcccc	gcctcaggag	gycacacttg	ttgcrmtggt	gagcagcstg	ctagtgggtg	120
ttrccctggc	cttcctgggg	ctcttcttcc	tcwacygcaa	gcagttcttc	aacagacatt	180
gycagcsgng	gaggtttgct	gcagtttgag	gctgatraaa	cagcaaagga	ggaatctstm	240
ttycycgtgc	cacccagcaa	ggagaccagt	gctgagtccc	aagtgagtga	gaacatyttt	300
cakaccagm	cacttaaccc	tatcctyrag	gacgactgca	rctcgactag	tgggttcccc	360
acacaggart	mctttaccat	ggcctyctgc	acctyagaga	gccactscca	ctgggwccac	420
arcccatcg	aatgcacaka	gctggacctg	caaaagtttt	ccagctctgc	ctcctatact	480
ggagctgara	ccttgggggg	aaacacagnc	aaaagcactg	ganacaggg		529

&lt;210&gt; 32

&lt;211&gt; 401

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(401)

&lt;223&gt; n = A,T,C or G



&lt;400&gt; 32

```

cagttgagct tagtggaggc agatgcaccc acagtgcccc ctcaggaggc cacacttggt      60
gsactggagg tttgctgcag tttgaggctg ataaaacagc aaaggaggaa tctctcttins    120
ccgtgccacc cagcaaggag accagtgctg agtcccaagt gagtgagaac atctttcaga      180
cccagccact taaccctatc ctcgaggacg actgcagctc gactagtggc ttccccacac      240
aggagtccct taccatggcc tcctgcacct cagagagcca ctcccactgg gtccacagcc      300
ccatcgaatg cacagagctg gacctgcaaa agttttccag ctctgcctcc tatactggag      360
ctgagacctt ggggggaaac acagtcgaaa gcactggaga c                                401

```

&lt;210&gt; 33

&lt;211&gt; 528

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 33

```

ctctgagggt caatgtgcct tccagttgag cttagtggag gcagatgcac ccacagtgcc      60
ccctcaggag gccacacttg ttgcactggt gaggagcctg ctagtgggtg ttaccctggc      120
cttcctgggg ctcttcttcc tctactgcaa gcagttcttc aacagacatt gccagcgtgt      180
tgcaggaggt ttgctgcagt ttgaggctga taaaacagca aaggagggaat ctctcttccc      240
cgtgccaccc agcaaggaga ccagtgtctg gtcccaagtg agtgagaaca tctttcagac      300
ccagccactt aaccctatcc tcgaggacga ctgcagctcg actagtggct tccccacaca      360
ggagtccttt accatggcct cctgcacctc agagagccac tcccactggg tccacagccc      420
catcgaatgc acagagctgg acctgcaaaa gttttccagc tctgcctcct atactggagc      480
tgagaccttg gggggaaaca cagtcgaaag cactggagac aggctgga                    528

```

&lt;210&gt; 34

&lt;211&gt; 175

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

&lt;400&gt; 34

```

Ser Glu Val Gln Cys Ala Phe Gln Leu Ser Leu Val Glu Ala Asp Ala
 1              5              10              15
Pro Thr Val Pro Pro Gln Glu Ala Thr Leu Val Ala Leu Val Ser Ser
      20              25              30
Leu Leu Val Val Phe Thr Leu Ala Phe Leu Gly Leu Phe Phe Leu Tyr
      35              40              45
Cys Lys Gln Phe Phe Asn Arg His Cys Gln Arg Val Ala Gly Gly Leu
 50              55              60
Leu Gln Phe Glu Ala Asp Lys Thr Ala Lys Glu Ser Leu Phe Pro
65              70              75              80
Val Pro Pro Ser Lys Glu Thr Ser Ala Glu Ser Gln Val Ser Glu Asn
      85              90              95

```

Ile Phe Gln Thr Gln Pro Leu Asn Pro Ile Leu Glu Asp Asp Cys Ser  
 100 105 110  
 Ser Thr Ser Gly Phe Pro Thr Gln Glu Ser Phe Thr Met Ala Ser Cys  
 115 120 125  
 Thr Ser Glu Ser His Ser His Trp Val His Ser Pro Ile Glu Cys Thr  
 130 135 140  
 Glu Leu Asp Leu Gln Lys Phe Ser Ser Ser Ala Ser Tyr Thr Gly Ala  
 145 150 155 160  
 Glu Thr Leu Gly Gly Asn Thr Val Glu Ser Thr Gly Asp Arg Leu  
 165 170 175

<210> 35  
 <211> 299  
 <212> PRT  
 <213> Homo sapiens

<400> 35  
 Met Asp Cys Gln Glu Asn Glu Tyr Trp Asp Gln Trp Gly Arg Cys Val  
 1 5 10 15  
 Thr Cys Gln Arg Cys Gly Pro Gly Gln Glu Leu Ser Lys Asp Cys Gly  
 20 25 30  
 Tyr Gly Glu Gly Gly Asp Ala Tyr Cys Thr Ala Cys Pro Pro Arg Arg  
 35 40 45  
 Tyr Lys Ser Ser Trp Gly His His Lys Cys Gln Ser Cys Ile Thr Cys  
 50 55 60  
 Ala Val Ile Asn Arg Val Gln Lys Val Asn Cys Thr Ala Thr Ser Asn  
 65 70 75 80  
 Ala Val Cys Gly Asp Cys Leu Pro Arg Phe Tyr Arg Lys Thr Arg Ile  
 85 90 95  
 Gly Gly Leu Gln Asp Gln Glu Cys Ile Pro Cys Thr Lys Gln Thr Pro  
 100 105 110  
 Thr Ser Glu Val Gln Cys Ala Phe Gln Leu Ser Leu Val Glu Ala Asp  
 115 120 125  
 Ala Pro Thr Val Pro Pro Gln Glu Ala Thr Leu Val Ala Leu Val Ser  
 130 135 140  
 Ser Leu Leu Val Val Phe Thr Leu Ala Phe Leu Gly Leu Phe Phe Leu  
 145 150 155 160  
 Tyr Cys Lys Gln Phe Phe Asn Arg His Cys Gln Arg Val Ala Gly Gly  
 165 170 175  
 Leu Leu Gln Phe Glu Ala Asp Lys Thr Ala Lys Glu Glu Ser Leu Phe  
 180 185 190  
 Pro Val Pro Pro Ser Lys Glu Thr Ser Ala Glu Ser Gln Val Ser Glu  
 195 200 205

Asn Ile Phe Gln Thr Gln Pro Leu Asn Pro Ile Leu Glu Asp Asp Cys  
 210 215 220  
 Ser Ser Thr Ser Gly Phe Pro Thr Gln Glu Ser Phe Thr Met Ala Ser  
 225 230 235 240  
 Cys Thr Ser Glu Ser His Ser His Trp Val His Ser Pro Ile Glu Cys  
 245 250 255  
 Thr Glu Leu Asp Leu Gln Lys Phe Ser Ser Ser Ala Ser Tyr Thr Gly  
 260 265 270  
 Ala Glu Thr Leu Gly Gly Asn Thr Val Glu Ser Thr Gly Asp Arg Leu  
 275 280 285  
 Glu Leu Asn Val Pro Phe Glu Val Pro Ser Pro  
 290 295

<210> 36  
 <211> 431  
 <212> DNA  
 <213> Homo sapiens

<400> 36  
 ctc tga ggt tca atg tgc ctt cca gtt gag ctt agt gga ggc aga tgc 48  
 acc cac agt gcc ccc tca gga ggc cac act tgt tgc act gga ggt ttg 96  
 ctg cag ttt gag gct gat aaa aca gca aag gag gaa tct ctc ttc ccc 144  
 gtg cca ccc agc aag gag acc agt gct gag tcc caa gtg agt gag aac 192  
 atc ttt cag acc cag cca ctt aac cct atc ctc gag gac gac tgc agc 240  
 tcg act agt ggc ttc ccc aca cag gag tcc ttt acc atg gcc tcc tgc 288  
 acc tca gag agc cac tcc cac tgg gtc cac agc ccc atc gaa tgc aca 336  
 gag ctg gac ctg caa aag ttt tcc agc tct gcc tcc tat act gga gct 384  
 gag acc ttg ggg gga aac aca gtc gaa agc act gga gac agg ctg ga 431

<210> 37  
 <211> 142  
 <212> PRT  
 <213> Homo sapiens

<400> 37  
 Leu Gly Ser Met Cys Leu Pro Val Glu Leu Ser Gly Gly Arg Cys Thr  
 1 5 10 15  
 His Ser Ala Pro Ser Gly Gly His Thr Cys Cys Thr Gly Gly Leu Leu  
 20 25 30  
 Gln Phe Glu Ala Asp Lys Thr Ala Lys Glu Glu Ser Leu Phe Pro Val  
 35 40 45  
 Pro Pro Ser Lys Glu Thr Ser Ala Glu Ser Gln Val Ser Glu Asn Ile  
 50 55 60

Phe Gln Thr Gln Pro Leu Asn Pro Ile Leu Glu Asp Asp Cys Ser Ser  
 65 70 75 80  
 Thr Ser Gly Phe Pro Thr Gln Glu Ser Phe Thr Met Ala Ser Cys Thr  
 85 90 95  
 Ser Glu Ser His Ser His Trp Val His Ser Pro Ile Glu Cys Thr Glu  
 100 105 110  
 Leu Asp Leu Gln Lys Phe Ser Ser Ala Ser Tyr Thr Gly Ala Glu  
 115 120 125  
 Thr Leu Gly Gly Asn Thr Val Glu Ser Thr Gly Asp Arg Leu  
 130 135 140

<210> 38  
 <211> 173  
 <212> PRT  
 <213> Homo sapiens

<400> 38  
 Met Asp Cys Gln Glu Asn Glu Tyr Trp Asp Gln Trp Gly Arg Cys Val  
 1 5 10 15  
 Thr Cys Gln Arg Cys Gly Pro Gly Gln Glu Leu Ser Lys Asp Cys Gly  
 20 25 30  
 Tyr Gly Glu Gly Gly Asp Ala Tyr Cys Thr Ala Cys Pro Pro Arg Arg  
 35 40 45  
 Tyr Lys Ser Ser Trp Gly His Lys Cys Gln Ser Cys Ile Thr Cys  
 50 55 60  
 Ala Val Ile Asn Arg Val Gln Lys Val Asn Cys Thr Ala Thr Ser Asn  
 65 70 75 80  
 Ala Val Cys Gly Asp Cys Leu Pro Arg Phe Tyr Arg Lys Thr Arg Ile  
 85 90 95  
 Gly Gly Leu Gln Asp Gln Glu Cys Ile Pro Cys Thr Lys Gln Thr Pro  
 100 105 110  
 Thr Ser Glu Val Gln Cys Ala Phe Gln Leu Ser Leu Val Glu Ala Asp  
 115 120 125  
 Ala Pro Thr Val Pro Pro Gln Glu Ala Thr Leu Val Ala Leu Glu Val  
 130 135 140  
 Cys Cys Ser Leu Arg Leu Ile Lys Gln Gln Arg Arg Asn Leu Ser Ser  
 145 150 155 160  
 Pro Cys His Pro Ala Arg Arg Pro Val Leu Ser Pro Lys  
 165 170

<210> 39  
 <211> 519  
 <212> DNA

<213> Artificial Sequence

<220>

<223> degenerate polynucleotide sequence

<221> misc\_feature

<222> (1)...(519)

<223> n = A,T,C or G

<400> 39

atggaytgyc	argaraayga	rtaytgggay	cartggggnm	gntgygtnac	ntgycarmgn	60
tgyggncng	gncargaryt	nwsnaargay	tgyggntayg	gngarggngg	ngaygcntay	120
tgyacngcnt	gyccnccnmg	nmgntayaar	wsnwsntggg	gncaycayaa	rtgycarwsn	180
tgyathacnt	gygcngtnat	haaymgngtn	caraargtna	aytgyacngc	nacnwsnaay	240
gcngtntgyg	gngaytgyyt	nccnmgntty	taymgnaara	cnmgcnathgg	nggnytnear	300
gaycargart	gyathccntg	yacnaarcac	acnccnacnw	sngargtnca	rtgygcntty	360
carytnwsny	tngtngargc	ngaygcncn	acngtnccnc	cncargargc	nacnytngt	420
gcnytnarg	tntgytgyws	nytnmgnytn	athaarcac	armgmnmgnaa	yytnwsnwsn	480
cntgycayc	cngcnmgmng	nccngtnytn	wsnccnaar			519

<210> 40

<211> 47

<212> DNA

<213> Artificial Sequence

<220>

<223> oligonucleotide ZC26463

<400> 40

atgcattaac	cctcactaaa	gggccttcct	ggggctcttc	ttcctct	47
------------	------------	------------	------------	---------	----

<210> 41

<211> 46

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide sequence ZC 26464

<400> 41

taatacgact	cactataggg	aggggcccct	gctgctgttg	tggtat	46
------------	------------	------------	------------	--------	----

<210> 42

<211> 49  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Oligonucleotide ZC24670

<400> 42  
 atgcattaac cctcactaaa gggacctgtg ctgtcatcaa tcgtgttca 49

<210> 43  
 <211> 47  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Oligonucleotide sequence ZC 26471

<400> 43  
 taatacgact cactataggg agggccccaa ggtctcagct ccagtat 47

<210> 44  
 <211> 657  
 <212> DNA  
 <213> murine

<400> 44  
 ggtggcatct ctcttccaat tggctctgatt gttggagtga catcactggg tctgctgatg 60  
 ttaggactgg tgaactgcat catcctgggtg cagaggaaaa agaagccctc ctgcctacaa 120  
 agagatgccca aggtgcctca tgtgcctgat gagaaatccc aggatgcagt aggccttgag 180  
 cagcagcacc tgttgaccac agcaccagct tccagcagca gctccctaga gagctcagcc 240  
 agcgtctgggg accgaagggc gccccctggg ggccatcccc aagcaagagt catggcggag 300  
 gcccaagggg ttcaggaggc ccgtgccagc tccaggattt cagattcttc ccacggaagc 360  
 cacgggacct acgtcaacgt cacctgcac gtgaacgtct gtagcagctc tgaccacagt 420  
 tctcagtgtc cttcccaagc cagcgccaca gtgggagacc cagatgccaa gccctcagcg 480  
 tccccaaagg atgagcaggt ccccttctct caggaggagt gtccgtctca gtccccgtgt 540  
 gagactacag agacactgca gagccatgag aagcccttgc cccttggtgt gccggatatg 600  
 ggcatgaagc ccagccaagc tggctgggtt gatcagattg cagtcaaagt ggcctga 657

<210> 45  
 <211> 824  
 <212> DNA

<213> Artificial Sequence

<220>

<223> artificial cDNA sequence

<400> 45

ggtaccgaat	tgtacgcgta	tggggacttc	ccatatcaat	cagggacttt	ccgctgggga	60
ctttccggtc	tgactcatgc	ttctgactca	tgcttggtg	acatcatctc	gactagtcgt	120
accttcccgt	aaatccctcc	ccttcccgga	attacacacg	cgtatttccc	agaaaaggaa	180
ctgtagattt	ctaggaattc	aatccttggc	cacgcgttta	caccggaagt	tttccatatt	240
aggaattcct	tccggtttcc	tttctcgagg	ccaccgtggt	tgagcccgac	actcattcat	300
aaaacgcttg	ttataaaagc	agtggctgcg	gcgccttcgt	actccaaccg	catctgcagc	360
gagcaactga	gaagccaagg	atccaggctg	aattcatggg	tctcaacccc	cagctagtgt	420
tcatcctgct	cttctttctc	gaatgtacca	ggagccatat	ccacggatgc	gacaaaaatc	480
acttgagaga	gatcatcggc	attttgaacg	aggtcacagg	agaagggacg	ccatgcacgg	540
agatggatgt	gccaaacgtc	ctcacagcaa	cgaagaacac	cacagagagt	gagctcgtct	600
gtagggcttc	caaggtgctt	cgcataat	attttaaaca	tgggaaaact	ccatgcttga	660
agaagaactc	tagtgttctc	atggagctgc	agagactcct	tggggctttt	cgatgcctgg	720
attcatcgat	aagctgcacc	atgaatgagt	ccaagtccac	atcactgaaa	gacttcctgg	780
aaagcctaaa	gagcatcatg	caaatggatt	actcgtagtc	taga		824

<210> 46

<211> 47

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide sequence ZC28835

<400> 46

taatacgact	cactataggg	aggcccccaa	ggtctcagct	ccagtat	47
------------	------------	------------	------------	---------	----

<210> 47

<211> 27

<212> DNA

<213> Artificial Sequence

<220>

<223> Oligonucleotide sequence ZC 28836

<400> 47

gcaccggtgg cctcctgagg gggcact 27

<210> 48  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Oligonucleotide ZC 28830

<400> 48  
 gcaccggtgg catctctctt ccaattgggt 29

<210> 49  
 <211> 29  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> Oligonucleotide ZC 28837

<400> 49  
 gctctagagg ggtcaggcca ctttgactg 29

<210> 50  
 <211> 1081  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <223> DNA construct

<400> 50  
 atggattgcc aagaaaatga gtactgggac caatggggac ggtgtgtcac ctgccaacgg 60  
 tgtggtcctg gacaggagct atccaaggat tgtggttatg gagagggtgg agatgcctac 120  
 tgcacagcct gccctcctcg caggtacaaa agcagctggg gccaccacaa atgtcagagt 180  
 tgcatacct gtgctgtcat caatcgtgtt cagaaggcca actgcacagc tacctctaata 240  
 gctgtctgtg gggactgttt gccaggttc taccgaaaga cacgcattgg aggctgcag 300  
 gaccaagagt gcatcccggt cacgaagcag acccccacct ctgagggttca atgtgccttc 360  
 cagttgagct tagtgaggc agatgcaccc acagtgtccc ctcaggaggc caccgttggc 420  
 atctctcttc caattggtct gattgttggg gtgacatcac tgggtctgct gatgttagga 480  
 ctggtgaact gcatcatcct ggtgcagagg aaaaagaagc cctcctgcct acaaagagat 540  
 gccaaagggtg ctcattgtgc tgatgagaaa tcccaggatg cagtaggcct tgagcagcag 600



a

<b>COMBINED DECLARATION FOR PATENT AND POWER OF ATTORNEY</b> (Includes Reference to PCT International Applications)			File No. 99-75		
<p>As a below named inventor, I hereby declare that:  My residence, post office address and citizenship are as stated below next to my name; I believe that I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:</p> <p><b>UMLR POLYPEPTIDES</b></p> <p>the specification of which (check only one item below):</p> <p><input checked="" type="checkbox"/> is attached hereto    <input type="checkbox"/> was filed as United States application Serial No.    on October 23, 2000</p> <p>and was amended on _____</p> <p><input type="checkbox"/> was filed as PCT international application Number _____ on _____</p> <p>I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, 1.56. I hereby claim foreign priority benefits under Title 35, United States Code, 119 of any foreign application(s) for patent or inventor's certificate(s) or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application(s) for patent or inventor's certificate(s) or any PCT international application(s) designating at least one country other than the United States of America filed by me on the same subject matter having a filing date before that of the application(s) of which priority is claimed:</p>					
<b>PRIOR FOREIGN/PCT APPLICATION(S) AND ANY PRIORITY CLAIMS UNDER 35 U.S.C. 119:</b>					
COUNTRY	APPLICATION NUMBER	DATE OF FILING	PRIORITY CLAIMED		
			<input type="checkbox"/> YES <input type="checkbox"/> NO		
			<input type="checkbox"/> YES <input type="checkbox"/> NO		
			<input type="checkbox"/> YES <input type="checkbox"/> NO		
I hereby claim the benefit under Title 35 United States Code 119(e) of any United States provisional application(s) listed below.					
U.S. APPLICATION NUMBER			U.S. FILING DATE		
60/160,880			October 22, 1999		
60/163,215			November 2, 1999		
60/218,769			July 17, 2000		
60/222,221			August 1, 2000		
I hereby claim the benefit under Title 35, United States Code 120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in that/those prior application(s) in the manner provided by the first paragraph of Title 35, United States Code, 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, 1.56 which occurred between the filing date of the prior application(s) and the national or PCT international filing date of this application:					
<b>PRIOR U.S. APPLICATIONS OR PCT INTERNATIONAL APPLICATIONS DESIGNATING THE U.S. FOR BENEFIT</b>					
<i>U.S. APPLICATIONS</i>				STATUS (check one)	
U.S. APPLICATION NUMBER	U.S. FILING DATE	Patented	Pending	Abandoned	
<i>PCT APPLICATIONS DESIGNATING THE U.S.</i>					
APPLICATION	FILING DATE	U.S. SERIAL NUMBERS ASSIGNED (if any)			

**POWER OF ATTORNEY:** As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

Robyn Adams  
Reg. No. 44,495

Jennifer K. Johnson  
Reg. No. 43,696

Phillip B.C. Jones  
Reg. No. 38,195

Debra K. Leith  
Reg. No. 32,619

Susan E. Lingenfelter  
Reg. No. 41,156

Paul G. Lunn  
Reg. No. 32,743

Gary E. Parker  
Reg. No. 31,648

Deborah A. Sawislak  
Reg. No. 37,438

**Send Correspondence To:** Robyn Adams  
ZymoGenetics, Inc.  
1201 Eastlake Avenue East  
Seattle, WA 98102

**Direct Telephone Calls To:**  
Robyn Adams  
(206) 442-6752

1	Full Name	Family Name Xu	First Given Name Wenfeng	Second Given Name
	Residence	City Mukilteo	State or Foreign Country WA	Country of Citizenship CN
	Post Office Address	Post Office Address 12432 54th Ave W	City Mukilteo	State & Zip Code/Country WA 98275/US
2	Full Name	Family Name Lofton-Day	First Given Name Catherine	Second Given Name E.
	Residence	City Brier	State or Foreign Country WA	Country of Citizenship US
	Post Office Address	Post Office Address 23908 35th Ave W	City Brier	State & Zip Code/Country WA 98036/US
3	Full Name	Family Name Henne	First Given Name Randall	Second Given Name
	Residence	City	State or Foreign Country	Country of Citizenship
	Post Office Address	Post Office Address	City	State & Zip Code/Country
4	Full Name	Family Name Presnell	First Given Name Scott	Second Given Name R.
	Residence	City Tacoma	State or Foreign Country WA	Country of Citizenship US
	Post Office Address	Post Office Address 2902 N. Puget Sound Ave	City Tacoma	State & Zip Code/Country WA 98407/US
5	Full Name	Family Name Yao	First Given Name Yue	Second Given Name
	Residence	City Kenmore	State or Foreign Country WA	Country of Citizenship US
	Post Office Address	Post Office Address 18216 61 <sup>st</sup> Ave. N.E.	City Kenmore	State & Zip Code/Country WA 98028/US
6	Full Name	Family Name Novak	First Given Name Julia	Second Given Name E.
	Residence	City Bainbridge Island	State or Foreign Country WA	Country of Citizenship US
	Post Office Address	Post Office Address 10699 Battle Point Dr. NE	City Bainbridge Island	State & Zip Code/Country WA 98110/US

I hereby declare that all statement made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application any patent issuing thereon.

Signature of Inventor 1	Signature of Inventor 2	Signature of Inventor 3
Date	Date	Date
Signature of Inventor 4	Signature of Inventor 5	Signature of Inventor 6
Date	Date	Date

**POWER OF ATTORNEY:** As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

Robyn Adams  
Reg. No. 44,495

Jennifer K. Johnson  
Reg. No. 43,696

Phillip B.C. Jones  
Reg. No. 38,195

Debra K. Leith  
Reg. No. 32,619

Susan E. Lingenfelter  
Reg. No. 41,156

Paul G. Lunn  
Reg. No. 32,743

Gary E. Parker  
Reg. No. 31,648

Deborah A. Sawislak  
Reg. No. 37,438

**Send Correspondence To:** Robyn Adams  
ZymoGenetics, Inc.  
1201 Eastlake Avenue East  
Seattle, WA 98102

**Direct Telephone Calls To:**  
Robyn Adams  
(206) 442-6752

7	Full Name	Family Name Foster	First Given Name Donald	Second Given Name C.
	Residence	City Lake Forest Park	State or Foreign Country WA	Country of Citizenship US
	Post Office Address	Post Office Address 3002 NE 181st St	City Lake Forest Park	State & Zip Code/Country WA 98155/US
8	Full Name	Family Name Yee	First Given Name David	Second Given Name P.
	Residence	City Cambridge	State or Foreign Country MA	Country of Citizenship US
	Post Office Address	Post Office Address 116 Henry Street	City Cambridge	State & Zip Code/Country MA 02139/US
9	Full Name	Family Name	First Given Name	Second Given Name
	Residence	City	State or Foreign Country	Country of Citizenship
	Post Office Address	Post Office Address	City	State & Zip Code/Country
10	Full Name	Family Name	First Given Name	Second Given Name
	Residence	City	State or Foreign Country	Country of Citizenship
	Post Office Address	Post Office Address	City	State & Zip Code/Country
11	Full Name	Family Name	First Given Name	Second Given Name
	Residence	City	State or Foreign Country	Country of Citizenship
	Post Office Address	Post Office Address	City	State & Zip Code/Country
12	Full Name	Family Name	First Given Name	Second Given Name
	Residence	City	State or Foreign Country	Country of Citizenship
	Post Office Address	Post Office Address	City	State & Zip Code/Country

I hereby declare that all statement made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application any patent issuing thereon.

Signature of Inventor 7	Signature of Inventor 8	Signature of Inventor 9
Date	Date	Date
Signature of Inventor 10	Signature of Inventor 11	Signature of Inventor 12
Date	Date	Date